

CMS Draft Analysis Note

The content of this note is intended for CMS internal use and distribution only

2019/05/07

Head Id: 494152

Archive Id: 494391

Archive Date: 2019/04/29

Archive Tag: trunk

Search for SM tttt in the same-sign dilepton and multi-lepton final states at $\sqrt{s} = 13$ TeV with the full Run 2 dataset

L. Bauerdick¹, K. Burkett¹, O. Gutsche¹, S. Jindariani¹, H. Weber¹, M. Derdzinski², B. Hashemi², D. Gilbert², D. Klein², S. May², D. Olivito², I. Suarez², F. Würthwein², A. Yagil², G. Zevi Della Porta², N. Amin³, C. Campagnari³, B. Marsh³, S. Wang³, C. Fangmeier⁴, and F. Golf⁴

¹ Fermi National Accelerator Laboratory, Batavia, IL, USA

² University of California, San Diego, CA, USA

³ University of California, Santa Barbara, CA, USA

⁴ University of Nebraska, Lincoln, NE, USA

Abstract

This is the AN supporting the Run 2 tttt analysis. It is based on the 2016 tttt AN (AN-17-115) and the 2016 same-sign AN (AN-16-386).

This box is only visible in draft mode. Please make sure the values below make sense.

PDFAuthor: N. Amin, C. Campagnari, C. Fangmeier, F. Golf, G. Zevi Della Porta

PDFTitle: Search for SM tttt in the same-sign dilepton final state at 13 TeV with the full Run 2 dataset

PDFSubject: CMS

PDFKeywords: CMS, physics, software, computing

Please also verify that the abstract does not use any user defined symbols

Contents

1			
2	1	Introduction	4
3	2	Samples	7
4	2.1	Collision data	7
5	2.2	Monte Carlo Simulation	7
6	3	Triggers	11
7	3.1	2016 trigger efficiency	11
8	3.2	2017 and 2018 trigger efficiency	11
9	4	Object Definition and Selection	13
10	4.1	Electron identification	13
11	4.2	Muon identification	13
12	4.3	Lepton isolation	14
13	4.4	Trigger emulation selection	15
14	4.5	Lepton definitions	16
15	4.6	Jets	16
16	4.7	Jets originating from b quarks	18
17	4.8	Missing transverse energy	18
18	5	Signal Extraction Strategy: cut-based	19
19	5.1	Baseline selection	19
20	5.2	Control regions selection	19
21	5.3	Signal regions selection	19
22	6	Signal Extraction Strategy: BDT	22
23	6.1	Intro	22
24	6.2	Selection and variables	22
25	6.3	Hyperparameters	24
26	6.4	Comparison with cut-based	26
27	6.5	Misc studies, all numbers and text is from 2016	28
28	7	Validation of kinematic variables	29
29	7.1	Opposite-sign dilepton events	29
30	7.2	Same-sign tight+fail dilepton events	29
31	7.3	Fake-enriched validation region in data	42
32	8	Background Estimations	43
33	8.1	Fake leptons	43
34	8.2	Charge misidentification	52
35	8.3	Rare SM processes	55
36	8.4	Theoretical uncertainties on TTW, TTZ and TTH	56
37	9	Systematic Uncertainties	58
38	9.1	Correlation model for 2016+2017+2018	58
39	9.2	Experimental sources of uncertainties	59
40	9.3	Uncertainties from data-driven background estimations	61
41	9.4	Systematic uncertainties from statistical sources	61
42	9.5	Signal uncertainties	61
43	9.6	Summary of theoretical uncertainties on MC-based processes	64

44	10	Kinematics	65
45	11	Results	79
46	12	Results: interpretations	85
47	12.1	Type-II 2HDM	85
48	12.2	Top Yukawa coupling	92
49	12.3	Off-shell mediators decaying to top quark pairs	98
50	12.4	Oblique Higgs parameter	103
51	12.5	Dark matter	108
52	A	Statistical checks	116
53	A.1	Impacts	116
54	A.2	Nuisance forms	119
55	A.3	Nuisances	120
56	A.4	Nuisance correlation matrix	125
57	A.5	Goodness of fits	126
58	B	Shape variations	127
59	B.1	BDT discriminator	127
60	C	Studies which did not result in analysis changes	130
61	C.1	Jet and b-jet thresholds	130
62	C.2	Taus	130
63	C.3	Top-tagging	133
64	D	Impact of the L1 ECAL prefiring(2016/2017) and HEM15/16 loss (2018) on the results	135
65	D.1	L1 prefiring(2016/2017)	135
66	D.2	HEM15/16 loss in 2018	137
68	E	Analysis changes	139
69	E.1	Changes from ANv7	139
70	F	Unblinding of 2016 dataset	142
71	F.1	Yields and results	142
72	F.2	Cut-based and BDT consistency checks	142
73	F.3	Nuisances	146
74	F.4	Impacts	150
75	G	Unblinding of 2017 dataset	153
76	G.1	Yields and results	153
77	G.2	Cut-based and BDT consistency checks	153
78	G.3	Nuisances	157
79	G.4	Impacts	161
80	G.5	Goodness of fits	164
81	G.6	Combination with 2016	165
82	H	Unblinding of 2018 dataset, BDT retraining	167
83	H.1	BDT retraining	167
84	H.2	Yields and results	167
85	H.3	Nuisances	172
86	H.4	Impacts	177

87	H.5	Goodness of fits	180
88	H.6	Tail BDT events	181
89	I	Checks from ARC review	182
90	I.1	Additional uncertainty from $tt+bb$	182
91	I.2	Distributions of BDT inputs for signal region events	183
92	I.3	More jets at higher η	183
93	I.4	Effect of $t\bar{t}H$ and $ttbb/ttjj$ measurement updates	184

DRAFT

1 Introduction

The same-sign dilepton final state is often used in searches for new physics due to its ability to stifle all but some rare standard model backgrounds. One of these rare backgrounds is the production of four top quarks, $pp \rightarrow t\bar{t}t\bar{t}$, primarily through the two processes depicted in Fig. 1. The lepton multiplicity composition in $t\bar{t}t\bar{t}$ events is shown in Fig. 2. With four W bosons, the same-sign dilepton and tri-lepton final state, considering electrons and muons (and those from arising from leptonic tau decays), occurs in approximately 11.9% of $t\bar{t}t\bar{t}$ events.

This Analysis Note describes a $t\bar{t}t\bar{t}$ analysis using the full Run 2 data, which is based on two analyses of the 2016 data: the SUSY same-sign dilepton analysis and the corresponding $t\bar{t}t\bar{t}$ analysis. The SUSY same-sign dilepton analysis used 35.9 fb^{-1} of 2016 collision data, is documented in [1], and published in [2]. The observed (expected) 95% confidence limit on cross section of four top quark production was found to be $42 (27_{-8}^{+13}) \text{ fb}$, and the cross section of the process calculated at NLO is $9.2_{-2.4}^{+2.9} \text{ fb}$ [3].

The 2016 $t\bar{t}t\bar{t}$ analysis, based on the same dataset as the SUSY search, was developed by modifying the selection of the SUSY analysis to take advantage of the jet and b-jet multiplicity associated with four top quark decays, as well as creating a separate category of events with 3 or more leptons. The resulting 2016 $t\bar{t}t\bar{t}$ analysis is document in [4], and [5]. The observed (expected) 95% confidence limit on cross section of four top quark production was found to be $41.7 (20.8_{-6.9}^{+11.2}) \text{ fb}$. The expected significance, based on an NLO cross section for $t\bar{t}t\bar{t}$ of $9.2_{-2.4}^{+2.9} \text{ fb}$ [3], was 1.05 standard deviations, corresponding to a measured signal strength parameter of $1.0_{-0.94}^{+1.2}$. The observed significance was 1.56 standard deviations (p-value of 6%), corresponding to a measured signal strength parameter of $1.83_{-1.23}^{+1.42}$. The results were also used to constrain the top quark Yukawa coupling of the Higgs boson, based on an LO*k-factor cross section of $12.2_{-4.4}^{+5.0} \text{ fb}$ and its dependence on $|y_T/y_T^{SM}|$. The central (upper,lower) value of the theoretical cross section band, combined with our 95% observed limit on $t\bar{t}t\bar{t}$ production, resulted in a 95% CL limit $|y_T/y_T^{SM}| < 2.27 (2.03,2.56)$. Additionally, an extension to the interpretation of heavy (pseudo)scalar bosons in the context of 2HDM in the manner of the SUSY analysis was studied, but it was not included in the $t\bar{t}t\bar{t}$ publication due to the very limited improvements with respect to the SUSY analysis, as discussed in Section 9 of Ref. [4].

All details of the 2016 analysis can be found in the SUSY AN [1] (Fake and Flip backgrounds) and the TOP AN [4] (everything else).




Figure 1: Leading order diagrams for $t\bar{t}t\bar{t}$ production.




Figure 2: Lepton multiplicity for decay of 4 W bosons.

125 With respect of the 2016 tttt analysis, the Run2 analysis includes the following changes and
 126 improvements:

- 127 • Increased luminosity of $35.922 + 41.53 + 58.83 = 136.3 \text{ fb}^{-1}$
- 128 • Update the cut-and-count signal regions from 8 to 18 bins in $N(\text{lep})/N(\text{jets})/N(\text{b-}$
 129 $\text{jets})$
- 130 • Note that this was the case for ANv7 and previous versions Update the correction
 131 to ttW+bb and ttZ+bb based on the Data/MC discrepancy in the ttbb/ttj ratio from
 132 1.7 ± 0.6 (TOP-16-010) to 1.0 ± 0.35 . This is based on the preliminary results of TOP-
 133 17-021, which finds a ttbb/ttj ratio of 1.0 ± 0.35 .
- 134 • This is the current prescription for ANv8 and on Maintain the correction to ttW+bb
 135 and ttZ+bb based on the Data/MC discrepancy in the ttbb/ttj ratio of 1.7 ± 0.6
 136 (TOP-16-010).
- 137 • Update $t\bar{t}t\bar{t}$ cross section to the latest and most precise estimate, from 9.2 fb [3] to
 138 11.97 fb [6].
- 139 • Reduced normalization uncertainty on $t\bar{t}H$ from 50% to 25% motivated by HIG-17-
 140 035.
- 141 • Studied two BDTs, one with a long list and one with a short list of variables. See
 142 Section 6.2.
- 143 • Optimized hyperparameter choice for BDT. See Section 6.3.

144 Upon looking at the 2017 and 2018 datasets, several more changes were adopted in order to
 145 deal with data-taking issues:

- 146 • Since the dilepton+HT triggers were not active for the first period of 2017 data taking
 147 (RunB, corresponding to 12% of the data), we decided to use the dilepton triggers
 148 throughout the 2017 run. The difference resulting from this change is expected to be
 149 at the few % level, although we will only know the exact number after all Trigger
 150 Scale Factors will be computed.
- 151 • Since an increase in non-prompt leptons was observed in our control regions, we
 152 tightened the isolation requirements with respect to 2016, as described in Section 4.3.

153 Issues relating to the missing HEM15/16 sectors starting in Run 319077 of 2018 data-collection
 154 were found to impact the fake background only negligibly, and no action was taken to specifi-
 155 cally deal with this as the estimation method is data-driven. Additionally, the L1 preferring issue
 156 affecting the efficiency of EGamma objects at high $|\eta|$ was found to have a small effect (%-level)
 157 which can be accounted for by a scale factor applied to simulation. See appendix D for checks,

158 and also see the 2LSS+multi-lepton SUSY analysis presentation at the SUSY Leptonic Meeting
 159 (slides 21-25 for HEM and slides 26-28 for L1 preferring at <https://indico.cern.ch/event/770079/>).

160 Several changes were considered, but it was decided not to implement them, as motivated
 161 below:

- 162 • Many additional variables were considered for the BDT, including resolved and
 163 boosted top-tagging variables, but they were not found to bring significant ($> 2\%$)
 164 gain. See Section 6.2.
- 165 • Separate BDT trainings for different background categories were considered, but
 166 since all backgrounds were found to have a similar BDT shape, it was decided to
 167 continue with a single training based on the sum of all backgrounds (slide 14 from
 168 Nick at <https://indico.cern.ch/event/709496/>).
- 169 • The LeptonMVA selection (combining ID, isolation and nearby jet information) was
 170 studied, but ultimately not adopted. Due to the different kinematics, it was found
 171 to increase $t\bar{t}V$ backgrounds by more than the $t\bar{t}t$ signal, resulting in no gain in
 172 significance. See Nick's presentation at the Top Cross Section meeting (slides 4-8 at
 173 <https://indico.cern.ch/event/709496/>).
- 174 • Jet and b-jet p_T thresholds were studied in the range of 20-40 GeV. Based on ex-
 175 pected significance, it was decided to stay with 40 GeV threshold for jets, and with
 176 25 GeV threshold for b-jets. See Appendix C.
- 177 • Including events with tau leptons. See Appendix C.
- 178 • Events with a 3rd lepton, with $10 < p_T < 20$ GeV and not forming a Z mass, were
 179 previously included in the 2-lepton signal regions. We considered removing these
 180 events, or moving them to a separate categories, but found that both treatments
 181 actually worsened (by a few %) the significance, as a non-negligible fraction of $t\bar{t}t$
 182 events falls into this category.

183 An extensive changelog with respect to ANv7 can be found in Appendix E.

184 Changelog with respect to ANv9

- 185 • Incorporation of lepton scale factors for 2017, 2018 (muons are incomplete for 2017,
 186 2018, and electrons are incomplete for 2018)
- 187 • Switched to preliminary Autumn18V3 JECs applied to 2018 data, MC (with no resid-
 188 uals for data)
- 189 • Added Appendix F and Appendix G showing the unblinding of 2016 and 2017
 190 datasets, respectively.
- 191 • For convenience, significance numbers will also be tracked in <https://docs.google.com/spreadsheets/d/140yqrUwEsJtOJ8OmDdOun-J8iUrV49aAdUkvfNjpn8k/edit#gid=0>

194 Changelog with respect to ANv10

- 195 • Updated to Autumn18V8 JECs applied to 2018 data, MC (includes residuals for 2018,
 196 split into eras A, B, C, and D)
- 197 • Updated to latest recommended b-tag SFs (V1) for 2018 MC
- 198 • Retrained the BDT with final samples and these updated corrections
- 199 • Added Appendix H which shows the results of unblinding 2018 and the full Run2
 200 dataset, with a retrained BDT

- Updated documentation to reflect the full unblinding
- For convenience, significance numbers are tracked in <https://docs.google.com/spreadsheets/d/140yqrUwEsJtOJ8OmDdOun-J8iUrV49aAdUkvfNjpn8k/edit#gid=0>

Changelog with respect to ANv12

- Included final lepton SFs (electrons, muons), finalized 2018 muon scale factors
- Included 50ipb from new re-reco golden JSON in 2018
- Updated to latest luminosity recommendation for 2018 (59.6 fb^{-1}), with a total of 137.2 fb^{-1}
- Switched to final luminosity uncertainty of 2.5% for 2018
- Included latest 2018 samples for heavy (pseudo)scalar boson interpretation

2 Samples

2.1 Collision data

The full Run 2 analysis uses an amount of pp collision data corresponding to an integrated luminosity of 35.9 fb^{-1} , 41.5 fb^{-1} , and 59.6 fb^{-1} , for 2016, 2017, and 2018, totaling 137.2 fb^{-1} .

The following datasets are used:

- DoubleEG/EGamma: two-electron channel,
- DoubleMuon: two-muon channel,
- MuonEG: crossed channel targeting events with one muon and one electron.
- JetHT: used to recover efficiency for dilepton+ H_T triggers in period 2016H (although no additional events are found in this sample).

The Reco campaigns used in this analysis are listed in table 1, together with the corresponding JSON file. Events present in multiple datasets are properly handled.

Table 1

Year	campaign	JSON file
2016	23Sep2016-v1	Cert.271036-284044.13TeV_23Sep2016ReReco_Collisions16
2017	31March2018-v1	Cert.294927-306462.13TeV_EOY2017ReReco_Collisions17
2018	17Sep2018-v1(2018A,B,C)-PromptReco(2018D)	Cert.314472-325175.13TeV_PromptReco_Collisions18_JSON.txt

In order to remove detector noise and unphysical events such beam-halo events, the recommended JetMET filters [7] have been applied.

2.2 Monte Carlo Simulation

The simulation samples used in this analysis have been produced in the Summer16 campaign for comparing to 2016 data, in the RunIIFall17MiniAOD-94X campaign for comparing to 2017 data, and in the RunIIAutumn18MiniAOD-102X campaign for 2018 data. The samples, almost exclusively at NLO, are listed in Table 2.

To improve on the MADGRAPH5_AMC@NLO modeling of the multiplicity of additional jets from initial-state radiation (ISR) and final-state radiation (FSR), $t\bar{t}W$ and $t\bar{t}Z$ MC events are

233 reweighted based on the number of ISR or FSR jets ($N_j^{\text{ISR/FSR}}$). The reweighting is based on the
 234 Data/MC ratio in the light-flavor jet multiplicity in dilepton $t\bar{t}$ events (using MADGRAPH5_AMC@NLO
 235 MC), as shown in Figure 3 and described in Ref. [8] for 2016 Data and MC. The method requires
 236 exactly two b-tagged jets, and assumes that all other jets are ISR or FSR. The reweighting factors
 237 vary between 0.86 and 0.77 for $N_j^{\text{ISR/FSR}}$ between 1 and 4 (where dilepton $t\bar{t}V$ plus 4 ISR/FSR
 238 jets results in $N_{\text{jets}}=8$, corresponding to our highest SR bin). For 2017 MC, new weights were
 239 derived since the MC uses a new Pythia tune. The corresponding Data/MC ratios are shown
 240 in Figure 4, where the different plots are based on different number of partons simulated at
 241 the matrix element. Reweighting factors are applied to $t\bar{t}W$, $t\bar{t}Z$, and $t\bar{t}$ (from inclusive sam-
 242 ple reweighting). Following the procedure used in the SUSY group, we take one half of the
 243 deviation from unity as the systematic uncertainty in these reweighting factors.




Figure 3: Distribution of the number of light jets (labeled ISR jets in the plot, but referring to ISR and FSR jets) in a dilepton $t\bar{t}$ sample in 2016 data, compared to 2016 MC.

244 To improve the modeling of the flavor of additional jets, the $t\bar{t}W$, $t\bar{t}Z$ and $t\bar{t}H$ simulation is also
 245 corrected to account for the measured ratio of $t\bar{t}b\bar{b}/t\bar{t}j$ cross sections reported in TOP-16-010.
 246 More details on these corrections and their uncertainties are provided in Section 9.




Figure 4: Distribution of the number of light jets in a dilepton $t\bar{t}$ sample with 0 additional partons (left) or 1 additional parton (right) for 2017 data (top) and 2018 data (bottom) compared to their respective MC samples. In the case of 0 additional partons, the reweighting factor in the last bin is taken instead from the previous bin.

Table 2: Signal and background samples from Summer16 and Fall17 campaigns. For each sample, the name has to be completed with the Summer16 string RunIISummer16MiniAODv2-PUMoriond17_80X_mcRun2_asymptotic_2016_TrancheIV_v6_v* for 2016, and with the Fall17 string RunIIFall17MiniAODv2-PU2017_12Apr2018_94X_mc2017_realist for 2017. For 2018, the string is RunIIAutumn18MiniAOD-102X_upgrade2018. Starting in 2017, the new Pythia tune "TuneCP5" was used instead of the CUETP8M2T4 and CUETP8M1 tunes. The $t\bar{t}t$ signal samples are highlighted in red. The ISR/FSR variation samples for $t\bar{t}t$ are not necessary starting in 2017, since the Parton Shower variation weights are already included in the nominal sample. **Remaining missing samples QCD samples are only used as a crosscheck, and the analysis does not depend on them.**

sample name	σ (pb)	2016	2017	2018
/DYjetsToLL.M-50.TuneCUETP8M1.13TeV-madgraphMLM-pythia8	6020.85	X	X	X
/DYjetsToLL.M-10to50.TuneCUETP8M1.13TeV-madgraphMLM-pythia8	18610	X	X	X
/WjetsToLNu.TuneCUETP8M1.13TeV-madgraphMLM-pythia8	61334.9	X	X	X
/TT.TuneCUETP8M2T4.13TeV-powheg-pythia8	831.762	X	X	X
/TTJets.TuneCP5.13TeV-amcatnloFXFX-pythia8	831.762	X	X	X
/TTWjetsToLNu.TuneCUETP8M1.13TeV-amcatnloFXFX-madspin-pythia8	0.2043	X	X	X
/TTZToLLNuNu.M-10.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.2529	X	X	X
/TTZToLL.M-1to10.TuneCUETP8M1.13TeV-madgraphMLM-pythia8	0.0493	X	X	X
/ttHTtoNonbb.M125.TuneCUETP8M2.ttHtranche3.13TeV-powheg-pythia8	0.2710	X	X	X
/tZq.ll.4f.13TeV-amcatnlo-pythia8.TuneCUETP8M1(ext1)	0.0758	X	X	X
/TTTT.TuneCP5.P5weights.13TeV-amcatnlo-pythia8	0.01197	-	X	X
/TTTT.TuneCUETP8M2T4.13TeV-amcatnlo-isrup-pythia8	0.01197	X	-	X
/TTTT.TuneCUETP8M2T4.13TeV-amcatnlo-fsrdown-pythia8	0.01197	X	-	X
/TTTT.TuneCUETP8M2T4.13TeV-amcatnlo-isrdown-pythia8	0.01197	X	-	X
/TTTT.TuneCUETP8M2T4.13TeV-amcatnlo-fsrdown-pythia8	0.01197	X	-	X
/TGjets.TuneCUETP8M1.13TeV-amcatnlo-madspin-pythia8	2.967	X	X	X
/TTGamma.SingleLeptFromT.TuneCUETP8M2T4.13TeV-amcatnlo-pythia8	0.77	X	X	X
/TTGamma.SingleLeptFromTbar.TuneCUETP8M2T4.13TeV-amcatnlo-pythia8	0.769	X	X	X
/TTGamma.Dilept.TuneCUETP8M2T4.13TeV-amcatnlo-pythia8	0.632	X	X	X
/WGToLNuG.TuneCUETP8M1.13TeV-amcatnloFXFX-pythia8	405.271	X	-	X
/WGToLNuG.TuneCP5.13TeV-madgraphMLM-pythia8	405.271	-	X	-
/ZGto2LG.TuneCUETP8M1.13TeV-amcatnloFXFX-pythia8	123.9	X	X	X
/WZto3LNu.TuneCUETP8M1.13TeV-powheg-pythia8	4.4297	X	X	X
/ZZto4L.13TeV-powheg-pythia8	1.256	X	X	X
/ZZZ.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.01398	X	X	X
/WZZ.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.05565	X	X	X
/WWZ.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.1651	X	X	X
/WZG.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.04123	X	X	X
/WWG.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.2147	X	X	X
/WWW_4F.TuneCUETP8M1.13TeV-amcatnlo-pythia8	0.2086	X	X	X
/WWto2L2Nu.DoubleScattering.13TeV-pythia8	0.16975	X	X	X
/ST.tWil5f.LO.13TeV-MadGraph-pythia8	0.01123	X	X	X
/WpWpJJ_EWK-QCD.TuneCUETP8M1.13TeV-madgraph-pythia8	0.03711	X	X	X
/GluGluHTtoZZto4L.M125.13TeV-powheg2.JHUGenV6.pythia8	0.01181	X	X	X
/VHTtoNonbb.M125.13TeV-amcatnloFXFX-madspin-pythia8	0.9561	X	X	X
/TTHH.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.000757	X	X	X
/TTZH.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.001535	X	X	X
/TTZZ.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.001982	X	X	X
/TTWZ.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.003884	X	X	X
/TTTW.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.000788	X	X	X
/TTTJ.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.000474	X	X	X
/TTWH.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.001582	X	X	X
/TTWW.TuneCUETP8M2T4.13TeV-madgraph-pythia8	0.01150	X	X	X
/QCD.Pt-20toInf.MuEnrichedPt15.TuneCUETP8M1.13TeV.pythia8	720648000 × 0.00042	X	X	TODO
/QCD.Pt-15to20.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	1273190000 × 0.003	X	X	TODO
/QCD.Pt-20to30.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	558528000 × 0.0053	X	X	TODO
/QCD.Pt-30to50.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	139803000 × 0.01182	X	X	TODO
/QCD.Pt-50to80.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	19222500 × 0.02276	X	X	TODO
/QCD.Pt-80to120.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	2758420 × 0.03844	X	X	TODO
/QCD.Pt-120to170.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	469797 × 0.05362	X	X	TODO
/QCD.Pt-170to300.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	117989 × 0.07335	X	X	TODO
/QCD.Pt-470to600.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	645.528 × 0.12242	X	X	TODO
/QCD.Pt-600to800.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	187.109 × 0.13412	X	X	TODO
/QCD.Pt-1000toInf.MuEnrichedPt5.TuneCUETP8M1.13TeV.pythia8	10.4305 × 0.15544	X	X	TODO
/QCD.Pt-20to30.EMEnriched.TuneCUETP8M1.13TeV.pythia8	557600000 × 0.0096	X	X	TODO
/QCD.Pt-30to50.EMEnriched.TuneCUETP8M1.13TeV.pythia8	136000000 × 0.073	X	X	TODO
/QCD.Pt-50to80.EMEnriched.TuneCUETP8M1.13TeV.pythia8	19800000 × 0.146	X	X	TODO
/QCD.Pt-80to120.EMEnriched.TuneCUETP8M1.13TeV.pythia8	2800000 × 0.125	X	X	TODO
/QCD.Pt-120to170.EMEnriched.TuneCUETP8M1.13TeV.pythia8	477000 × 0.132	X	X	TODO
/QCD.Pt-170to300.EMEnriched.TuneCUETP8M1.13TeV.pythia8	114000 × 0.165	X	X	TODO
/QCD.Pt-300toInf.EMEnriched.TuneCUETP8M1.13TeV.pythia8	9000 × 0.15	X	X	TODO
/QCD.Pt.20to30.bcToE.TuneCUETP8M1.13TeV.pythia8	557627000 × 0.00059	X	X	TODO
/QCD.Pt.30to80.bcToE.TuneCUETP8M1.13TeV.pythia8	159068000 × 0.00255	X	X	TODO
/QCD.Pt.80to170.bcToE.TuneCUETP8M1.13TeV.pythia8	3221000 × 0.01183	X	X	TODO
/QCD.Pt.170to250.bcToE.TuneCUETP8M1.13TeV.pythia8	105771 × 0.02492	X	X	TODO
/QCD.Pt.250toInf.bcToE.TuneCUETP8M1.13TeV.pythia8	21094.1 × 0.03375	X	X	TODO

3 Triggers

The analysis uses two slightly different trigger strategies for 2016 and 2017/2018 data. For 2016 data, the dilepton + H_T triggers are used, since they have slightly looser lepton requirements (they do not apply isolation), and are therefore slightly more efficient than the pure dilepton triggers. For 2017, these triggers were not available in RunB, which represent 12% of the 2017 dataset, and therefore the pure dilepton triggers are used for the entirety of 2017 and 2018. We considered also using the pure dilepton triggers for 2016, to further simplify this treatment, but we prefer to avoid unnecessary changes to the published result, wherever possible. The full set of triggers used for selecting events in the signal regions are listed in Table 3.

Table 3: Summary of the signal triggers

2016		
$H_{T,off}$	Channel	Trigger Name
> 300 GeV	$\mu\mu$	HLT_DoubleMu8_Mass8_PFHT300
	ee	HLT_DoubleEle8_CaloIdM_TrackIdM_Mass8_PFHT300
	$e\mu$	HLT_Mu8_Ele8_CaloIdM_TrackIdM_Mass8_PFHT300
2017		
$H_{T,off}$	Channel	Trigger Name
all	$\mu\mu$	HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL_DZ_Mass8
	ee	HLT_Ele23_Ele12_CaloIdL_TrackIdL_IsoVL_
	$e\mu$	HLT_Mu23_TrkIsoVVL_Ele12_CaloIdL_TrackIdL_IsoVL_DZ HLT_Mu8_TrkIsoVVL_Ele23_CaloIdL_TrackIdL_IsoVL_DZ
2018		
$H_{T,off}$	Channel	Trigger Name
all	$\mu\mu$	HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL_DZ_Mass3p8
	ee	HLT_Ele23_Ele12_CaloIdL_TrackIdL_IsoVL_
	$e\mu$	HLT_Mu23_TrkIsoVVL_Ele12_CaloIdL_TrackIdL_IsoVL_DZ HLT_Mu8_TrkIsoVVL_Ele23_CaloIdL_TrackIdL_IsoVL_DZ

An additional set of triggers is developed for a control sample selection for fake rate measurement. They require either a single lepton, or a single lepton and a jet in order to increase a purity of the collected data sample. The full list can be found in Table 4. Two sets of the auxiliary triggers are introduced in order to collect data with different online lepton IDs corresponding to the two sets of signal dilepton triggers.

3.1 2016 trigger efficiency

Computation of trigger efficiencies and scale factors for 2016 data is described in Section 4 of [1]. Summarizing these results, the dilepton+ H_T trigger efficiency is found to be the product of lepton leg efficiencies (93-98% for electrons, 85-95% for muons), and the HT efficiency (98-99%). As a result the trigger efficiency for events passing the baseline selection in the 2016 data is greater than 95% for ee and $e\mu$ events, and about 92% for $\mu\mu$ events.

3.2 2017 and 2018 trigger efficiency

This section is identical to same-sign SUS analysis (AN-18-280), which provides trigger efficiency maps. These are currently being applied for all 3 years.

Table 4: Summary of the control triggers ordered by lepton flavor and p_T .

2016		
$H_{T,off}$	Channel	Trigger Name
> 300 GeV	μ	HLT_Mu8 HLT_Mu17
	e	HLT_Ele8_CaloIdM_TrackIdM_PFJet30 HLT_Ele17_CaloIdM_TrackIdM_PFJet30
2017		
$H_{T,off}$	Channel	Trigger Name
all	μ	HLT_Mu8_TrkIsoVVL HLT_Mu17_TrkIsoVVL
	e	HLT_Ele8_CaloIdL_TrackIdL_IsoVL_PFJet30 HLT_Ele23_CaloIdL_TrackIdL_IsoVL_PFJet30
2018		
$H_{T,off}$	Channel	Trigger Name
all	μ	HLT_Mu8_TrkIsoVVL HLT_Mu17_TrkIsoVVL
	e	HLT_Ele8_CaloIdL_TrackIdL_IsoVL_PFJet30 HLT_Ele23_CaloIdL_TrackIdL_IsoVL_PFJet30

4 Object Definition and Selection

The object selection is based on the one used in the SUSY 2016 analysis documented in [1]. In particular, the same lepton ID is used for same-sign and trilepton events.

The somewhat complicated definitions of isolation and lepton IDs were arrived through dedicated studies within the "SUSY fake lepton working group" before data taking started in 2015. The goal was to have a fairly efficient lepton selection, with low fake rate, and with a reliable method to estimate nonprompt lepton backgrounds with minimal sample dependence starting from "fakeable objects" (FO). The studies are documented in Ref. [9].

4.1 Electron identification

The electrons are reconstructed by associating ECAL-clusters and GSF tracks. Only electrons within the tracker and ECAL acceptance are considered: $|\eta| < 2.5$.

The electron identification is performed using a multivariate (MVA) discriminant built with shower-shape variables ($\sigma_{i\eta i\eta}$, $\sigma_{i\phi i\phi}$, the cluster circularity, widths along η and ϕ , R_9 , H/E , E_{inES}/E_{raw}), track-cluster matching variables (E_{tot}/p_{in} , E_{Ele}/p_{out} , $\Delta\eta_{in}$, $\Delta\eta_{out}$, $\Delta\phi_{in}$, $1/E - 1/p$) and track quality variables (χ^2 of the KF and GSF tracks, the number of hits used by the KF/GSF filters, fbrem). A complete description of the discriminant and training used can be found in [10–12]. Three identification working points, summarized in the Table 5, are used in this analysis, following the prescriptions of the SUSY lepton scale factors group [13]. Signal events are selected using the tight working point while the loose working points are used in the estimate of background arising from mistakenly identified or non prompt leptons. Separate loose working points are derived for regions where isolated and non isolated dilepton triggers are used.

Different cuts are used in 2016, 2017, and 2018 based on the Spring16.GeneralPurpose.V1, Fall17NoIso, and Fall17V2NoIso trainings, respectively. Note that the cuts for 2018 are with respect to the "raw" BDT output (i.e., the values are not forced to lie between -1 and 1 by squashing the output with a sigmoid function). Raw values can be obtained from squashed output via $raw = \frac{1}{2} \ln \left(\frac{1+squashed}{1-squashed} \right)$. The working points depend of the momentum and pseudorapidity of the electrons: for 2016 (resp. 2017/2018) in the region between 15 (resp. 10) and 25 GeV, the working point is given by a linear interpolation between the higher and lower value, following $\min[A, \max[B, A + C * (p_T - 15(10))]]$, where A is the cut at 15(10) GeV, B is the cut at 25 GeV, and $C = (B - A)/10(15)$. Electrons with $p_T < 10$ GeV are only not used to select signal events, so the Tight WP is not assigned for them.

As the electron charge can be determined with three different techniques [14], and as the mis-measurement of the lepton charge can lead to accept more background after the selection, the requirement to get a consistent charge measurement with the three methods is considered in some of the electron definitions used in the analysis. To reject electrons originating from photon conversion, two variables are considered: the number of missing pixel hits and a conversion veto based on the vertex fit probability. Finally, impact parameter variables are also considered: impact parameter in the transverse plane d_0 , impact parameter along the z axis d_z , and the impact parameter significance in the detector space SIP_{3D} .

4.2 Muon identification

Two working points are considered for the muon identification. The loose working point follows the "muon POG Loose ID" described in [15], while the tight working point is given by the list of requirements known as the "muon Medium Id", defined in [16]. Only muons within

Table 5: Lower cut on the electron MVA discriminant for the various electron ID used in this analysis.

2016				
pseudorapidity region	momentum [GeV]	loose WP (non iso trigger)	loose WP (iso trigger)	tight WP
$0 < \eta < 0.8$	$5 < p_T < 10$	-0.30	-0.46	N/A
$0 < \eta < 0.8$	$10 < p_T < 15$	-0.86	-0.48	0.77
$0 < \eta < 0.8$	$p_T > 25$	-0.96	-0.85	0.52
$0.8 < \eta < 1.479$	$5 < p_T < 10$	-0.36	-0.03	N/A
$0.8 < \eta < 1.479$	$10 < p_T < 15$	-0.85	-0.67	0.56
$0.8 < \eta < 1.479$	$p_T > 25$	-0.96	-0.91	0.11
$1.479 < \eta < 2.5$	$5 < p_T < 10$	-0.63	0.06	N/A
$1.479 < \eta < 2.5$	$10 < p_T < 15$	-0.81	-0.49	0.48
$1.479 < \eta < 2.5$	$p_T > 25$	-0.95	-0.83	-0.01

2017				
pseudorapidity region	momentum [GeV]	loose WP (non iso trigger)	loose WP (iso trigger)	tight WP
$0 < \eta < 0.8$	$5 < p_T < 10$	-0.135	0.488	N/A
$0 < \eta < 0.8$	$10 < p_T < 25$	$-0.930 + \frac{0.043}{15} \times (p_T - 10)$	$-0.788 + \frac{0.148}{15} \times (p_T - 10)$	$0.2 + 0.032 \times (p_T - 10)$
$0 < \eta < 0.8$	$p_T > 25$	-0.887	-0.64	0.68
$0.8 < \eta < 1.479$	$5 < p_T < 10$	-0.417	-0.045	N/A
$0.8 < \eta < 1.479$	$10 < p_T < 25$	$-0.930 + \frac{0.04}{15} \times (p_T - 10)$	$-0.850 + \frac{0.073}{15} \times (p_T - 10)$	$0.1 + 0.025 \times (p_T - 10)$
$0.8 < \eta < 1.479$	$p_T > 25$	-0.890	-0.775	0.475
$1.479 < \eta < 2.5$	$5 < p_T < 10$	-0.470	0.176	N/A
$1.479 < \eta < 2.5$	$10 < p_T < 25$	$-0.942 + \frac{0.032}{15} \times (p_T - 10)$	$-0.810 + \frac{0.077}{15} \times (p_T - 10)$	$-0.1 + 0.028 \times (p_T - 10)$
$1.479 < \eta < 2.5$	$p_T > 25$	-0.910	-0.733	0.320

2018				
pseudorapidity region	momentum [GeV]	loose WP (non iso trigger)	loose WP (iso trigger)	tight WP
$0 < \eta < 0.8$	$5 < p_T < 10$	0.053	1.320	N/A
$0 < \eta < 0.8$	$10 < p_T < 25$	$-0.106 + 0.062 \times (p_T - 25)$	$1.204 + 0.066 \times (p_T - 25)$	$4.277 + 0.112 \times (p_T - 25)$
$0 < \eta < 0.8$	$p_T > 25$	-0.106	1.204	4.277
$0.8 < \eta < 1.479$	$5 < p_T < 10$	-0.434	0.192	N/A
$0.8 < \eta < 1.479$	$10 < p_T < 25$	$-0.769 + 0.038 \times (p_T - 25)$	$0.084 + 0.033 \times (p_T - 25)$	$3.152 + 0.060 \times (p_T - 25)$
$0.8 < \eta < 1.479$	$p_T > 25$	-0.769	0.084	3.152
$1.479 < \eta < 2.5$	$5 < p_T < 10$	-0.956	0.362	N/A
$1.479 < \eta < 2.5$	$10 < p_T < 25$	$-1.461 + 0.042 \times (p_T - 25)$	$-0.123 + 0.053 \times (p_T - 25)$	$2.359 + 0.087 \times (p_T - 25)$
$1.479 < \eta < 2.5$	$p_T > 25$	-1.461	-0.123	2.359

314 the muon system acceptance $|\eta| < 2.4$ are considered. Impact parameter selection is also ap-
315 plied on muons, and use the variables already defined in the previous section. Contrary to the
316 electrons, only one charge can be reconstructed for a muon track. The quality of the charge
317 reconstruction is then given for the muons by a quality criteria on the track reconstruction :
318 $\delta p_T(\mu) / p_T(\mu) < 0.2$.

319 4.3 Lepton isolation

320 The lepton isolation is constructed using three different variables:

- the mini isolation I_{mini} . Requiring I_{mini} below a given threshold ensures that the lep-
ton is locally isolated, even in boosted topologies. The impact of pileup is mitigated
using the so-called effective area correction:

$$I_{\text{mini}} = \frac{\sum_R p_T(h^\pm) - \max(0, \sum_R p_T(h^0) + p_T(\gamma) - \rho \mathcal{A} \left(\frac{R}{0.3}\right)^2)}{p_T(\ell)}. \quad (1)$$

where ρ is the pileup energy density, where $\sum_R p_T(h^\pm)$, $\sum_R p_T(h^0)$ and $\sum_R p_T(\gamma)$ refers to the sum of the transverse momentum of the charged hadrons, neutral hadrons and photons, respectively, within a cone R , dependent of the lepton p_T :

$$R = \frac{10}{\min(\max(p_T(\ell), 50), 200)} \quad (2)$$

321 The effective areas \mathcal{A} used are listed in Table 6.

322 • The second variable is the ratio of the lepton p_T and of the jet matched to the lepton
 323 p_T : $p_T^{\text{ratio}} = \frac{p_T(\ell)}{p_T(\text{jet})}$. This jet is matched geometrically to the lepton, and in most of
 324 the case is the jet containing the lepton. If no jet is clustering the lepton, then the
 325 closest one is chosen. The use of p_T^{ratio} is a simple way to identify leptons in quite
 326 boosted topologies, without any jet reclustering. In order to avoid an over-correction
 327 on prompt leptons, the application of the jet energy correction is only applied on the
 328 hadronic part of the jet, using the following formula $\text{jet} = \ell + (\text{jet-PU}-\ell) * JEC - PU$,
 329 where ℓ is the lepton, PU the pileup energy clustered into the jet, and JEC the jet
 330 energy scale correction to be applied. This formula is applied at the Lorentz vector
 331 level. This approach is commonly used in the B2G group and is blessed by the
 332 JetMET POG.

• The last variable used is the p_T^{rel} variable:

$$p_T^{\text{rel}} = \frac{|(\vec{p}(\text{jet}) - \vec{p}(\ell)) \times \vec{p}(\ell)|}{|\vec{p}(\text{jet}) - \vec{p}(\ell)|} \quad (3)$$

333 This variable allows to recover leptons from accidental with jets. Similarly to p_T^{ratio} ,
 334 the jet energy scale correction are only applied on the hadronic part of the considered
 335 jet.

Using those three variables, a lepton is considered isolated if the following condition is re-
 spected:

$$I_{\text{mini}} < I_1 \wedge (p_T^{\text{ratio}} > I_2 \vee p_T^{\text{rel}} > I_3) \quad (4)$$

336 The $I_i, i = 1, 2, 3$ values depends of the flavor of the lepton: as the probability to misidentify a
 337 jet is higher for the electrons, tighter isolation values are used. The loose lepton isolation is sig-
 338 nificantly relaxed, as well as an extra definition (fakeable) used for the fake lepton background
 339 estimation. The different values are summarized in the Table 7. The logic beyond that isolation
 340 is a relaxing of the local isolation, compensated by the requirement that the lepton carries
 341 the major part of the energy of the corresponding jet, or if not, if the lepton is considered as
 342 accidentally overlapping with a jet.

343 More details and figures about the lepton isolation can be found in the previous version of the
 344 analysis described in [17].

345 4.4 Trigger emulation selection

346 Due to the differences between online and offline reconstruction and selection, some leptons
 347 can be selected offline but not online. Even though we simulate this effect by applying trigger
 348 selection in the simulation, we still choose to reduce online/offline differences by applying a set
 349 of offline requirements to ensure that electrons will pass the trigger. Effectively this increases
 350 the efficiency of our trigger selection, reducing the role of trigger Data/MC scale factors. The
 351 requirements are summarized in Table 8.

Table 6: Effective areas for muons and electrons.
2016

Muons		Electrons	
$ \eta $ range	$\mathcal{A}(\mu)$ neutral	$ \eta $ range	$\mathcal{A}(e)$ neutral
0.0 – 0.8	0.0735	0.0 – 1.0	0.1752
0.8 – 1.3	0.0619	1.0 – 1.479	0.1862
1.3 – 2.0	0.0465	1.479 – 2.0	0.1411
2.0 – 2.2	0.0433	2.0 – 2.2	0.1534
2.2 – 2.5	0.0577	2.2 – 2.3	0.1903
		2.3 – 2.4	0.2243
		2.4 – 2.5	0.2687

2017/2018

Muons		Electrons	
$ \eta $ range	$\mathcal{A}(\mu)$ neutral	$ \eta $ range	$\mathcal{A}(e)$ neutral
0.0 – 0.8	0.0566	0.0 – 1.0	0.1440
0.8 – 1.3	0.0562	1.0 – 1.479	0.1562
1.3 – 2.0	0.0363	1.479 – 2.0	0.1032
2.0 – 2.2	0.0119	2.0 – 2.2	0.0859
2.2 – 2.5	0.0064	2.2 – 2.3	0.1116
		2.3 – 2.4	0.1321
		2.4 – 2.5	0.1654

Table 7: Isolation working points

2016

isolation value	loose WP (e/μ)	μ (Medium) WP	e (Tight) WP
I_1	0.4	0.16	0.12
I_2	0	0.76	0.80
I_3	0	7.2	7.2

2017/2018

isolation value	loose WP (e/μ)	μ (Medium) WP	e (Tight) WP
I_1	0.4	0.11	0.07
I_2	0	0.74	0.78
I_3	0	6.8	8.0

352 4.5 Lepton definitions

353 The loose, tight and fakeable lepton definition, based the different identification and isolation
354 definitions described in the previous sections, are summarized in the Table 9.

355 4.6 Jets

356 Jets are reconstructed from particle flow candidates, clustered with the anti-kt algorithm and
357 with a cone size of $\Delta R < 0.4$. Only jets above a transverse momentum $p_T > 40$ GeV and
358 within the tracker acceptance $|\eta| < 2.4$ are considered. To reject noise and mis-measured jets,
359 the selected jets have to fulfill several identification criteria.

360 For the 2016 data analysis the following criteria, corresponding to the loose JetID selection, are
361 applied:

- 362 • neutral hadronic energy fraction < 0.99

Table 8: Electron selection criteria used for the emulation of the high level trigger selection .

variable	barrel, $ \eta < 1.4442$	endcaps, $1.566 < \eta < 2.4$
Identification criteria		
$ \Delta\eta _{In} <$	0.01	0.01
$ \Delta\phi _{In} <$	0.04	0.08
$\sigma I\eta I\eta <$	0.011	0.031
$H/E <$	0.08	0.08
$ 1/E - 1/p <$	0.01	0.01
Isolation criteria (Not used with Dilepton+ H_T triggers, since those triggers do not apply isolation.)		
relative ecal PFCluster isolation (dR=0.3)		0.45
relative hcal PFCluster isolation (dR=0.3)		0.25
relative track p_T isolation		0.2

variable	muons			electrons		
	loose	fakable	tight	loose	fakable	tight
identification	loose ID	medium ID	medium ID	loose WP	loose WP	tight WP
isolation	loose WP	loose WP	μ WP	loose WP	loose WP	e WP
HLT emulation	-	-	-	-	- /yes	- /yes
d_0 (cm)	0.05	0.05	0.05	0.05	0.05	0.05
d_z (cm)	0.1	0.1	0.1	0.1	0.1	0.1
SIP _{3D}	-	< 4	< 4	-	< 4	< 4
missing inner hits	-	-	-	≤ 1	= 0	= 0
conversion veto	-	-	-	yes	yes	yes
tight charge	-	yes	yes	-	yes	yes

Table 9: Summary of the lepton selection.

- 363 • neutral electromagnetic energy fraction < 0.99
- 364 • number of constituents > 1
- 365 • charged hadron fraction > 0
- 366 • charged multiplicity > 0
- 367 • charged EM fraction < 0.99

368 For the 2017 and 2018 data analysis we followed the JetMet recommendation to switch to the
369 Tight JetID selection defined as the following set of conditions:

- 370 • neutral hadronic energy fraction < 0.9
- 371 • neutral electromagnetic energy fraction < 0.9
- 372 • number of constituents > 1
- 373 • charged hadron fraction > 0
- 374 • charged multiplicity > 0

375 The most recently recommended set of jet energy corrections (JEC) is applied:

- 376 • 2016: Summer16_23Sep2016 (corresponding Global Tag: 80X_dataRun2_2016SeptRepro.v7)
- 377 • 2017: Fall17_17Nov2017B_V32
- 378 • 2018: Autumn18_RunX_V8

To avoid double counting due to jets matched geometrically with a lepton, the jet the closest matched to a fakeable or tight lepton within $\Delta R < 0.4$ is not considered as a jet in the event.

From those selected jets, the key variable H_T is defined by

$$H_T = \sum_{jets} p_T.$$

379 **4.7 Jets originating from b quarks**

380 Jets defined in the previous section - with the only difference of a lower p_T threshold down
381 to 25 GeV - can be promoted as jets originating from b quark. The “deepCSV” discrimina-
382 tor is used for all years. The discriminator value for each jet is obtained by summing the
383 “ $x:probb$ ” and “ $x:probbb$ ” discriminators. In 2016, $x = \text{deepFlavourJetTags}$, in 2017, $x =$
384 pfDeepCSVJetTags , and in 2018, $x = \text{pfDeepCSVJetTags}$ [18]. The medium working point
385 requires the discriminant value to be greater than 0.6324 in 2016, 0.4941 in 2017, and 0.4184 in
386 2018.

387 When the jet multiplicity and H_T are computed, b-jets with $p_T > 40$ GeV are counted along
388 with standard jets.

389 **4.8 Missing transverse energy**

390 The missing transverse energy \cancel{E}_T is defined as the negative sum of the transverse momentum
391 of all particle flow particles reconstructed in an event. The jet energy corrections are propagated
392 to the \cancel{E}_T following the procedure described in [19].

393 For the analysis of 2017 data we follow the recommendation from the MET group [20] and ex-
394 clude from the MET calculation low p_T jets and unclustered PF candidates in the most forward
395 region of the ECAL (MET “v2” recipe).

5 Signal Extraction Strategy: cut-based

5.1 Baseline selection

With respect to Ref. [1], the baseline region was modified for the 2016 analysis to take advantage of the four top kinematics. The baseline region is designed to reject the majority of background (while preserving signal) before splitting kinematic phase space to form the signal regions.

With respect to the 2016 data analysis, the definition of individual signal regions has been modified (5.3), while the control regions (5.2) and the overall baseline selection (below) have remained the same.

Selected events must contain at least two tight (as defined in Table 9) same-sign leptons. To reject low-mass resonances, events with a pair of same-sign electrons with an invariant mass $m_{\ell\ell} < 12$ GeV are rejected. Events with three leptons of the same charge are also rejected.

In total, the baseline selection consists of the following:

- tight same-sign dileptons,
- $N_{\text{jets}} \geq 2$,
- $N_{\text{b jets}} \geq 2$,
- $\cancel{E}_T > 50$ GeV,
- $H_T > 300$ GeV,
- $p_{T,\text{lep1}} \geq 25$ GeV, $p_{T,\text{lep2}} \geq 20$ GeV.

Events containing a third loose (as defined in Table 9) lepton that makes a DY candidate (opposite charge, same flavor pair with $|m_{\ell\ell} - m_Z| < 15$ GeV or $m_{\ell\ell} < 12$ GeV) with one of the two SS dileptons are rejected from the signal regions, and a subset of them is assigned to the $t\bar{t}Z$ control region mentioned below. The transverse momentum of the third lepton is required to be larger than 5(7) GeV for muons (electrons).

If a Z candidate is not found, a tight third lepton present in the event with $p_T > 20$ GeV contributes to the N_{leps} count.

5.2 Control regions selection

Two control regions have been introduced to simultaneously constrain two dominant SM backgrounds: $t\bar{t}W$ and $t\bar{t}Z$.

The control region for $t\bar{t}Z$ (“CRZ”) consists of events passing the baseline selection with three leptons (where the third lepton has $p_T > 20$ GeV and passes the tight selection), with a Z boson candidate as described above.

The control region for $t\bar{t}W$ (“CRW”) consists of events with a same-sign lepton pair passing the tight lepton requirement, fewer than 6 jets, and exactly 2 b-tagged jets.

5.3 Signal regions selection

Signal regions are formed by subdividing the baseline region by number of leptons (N_{leps}), number of b-jets ($N_{\text{b jets}}$), and number of jets (N_{jets}).

The 2016+2017+2018 analysis signal regions and CRW/CRZ are tabulated in Table 10, and shown graphically in Figure 5, while the old 2016 signal regions are shown in Table 11 for reference.

Table 10: Definition of the 14 SRs and two CRs, CRW and CRZ, for the full Run2 analysis.

N_{leps}	$N_{\text{b jets}}$	N_{jets}	Region
2	2	≤ 5	CRW
		6	SR1
		7	SR2
		≥ 8	SR3
	3	5	SR4
		6	SR5
		7	SR6
		≥ 8	SR7
	≥ 4	≥ 5	SR8
	≥ 3	2	5
6			SR10
≥ 7			SR11
≥ 3		4	SR12
		5	SR13
		≥ 6	SR14
inverted Z-veto			CRZ

Table 11: SR definitions and CRW definition for the 2016 analysis.

N_{leps}	$N_{\text{b jets}}$	N_{jets}	Region
2	2	≤ 5	CRW
		6	SR1
		7	SR2
		≥ 8	SR3
	3	5, 6	SR4
		≥ 7	SR5
	≥ 4	≥ 5	SR6
≥ 3	2	≥ 5	SR7
	≥ 3	≥ 4	SR8
inverted Z-veto			CRZ




Figure 5: Signal regions and CRW (15 regions) with CRZ not displayed. Dotted edges indicate no explicit boundary.

6 Signal Extraction Strategy: BDT

6.1 Intro

A BDT approach was studied for the 2016 analysis, indicating that a 10-20% improvement in the expected significance could be reached with respect to the cut-based approach, in an optimistic scenario without additional systematics on the BDT input variables. For the 2016 analysis, the cut-based approach was chosen, resulting in a faster publication and a slightly lower performance. For the full Run 2 analysis, we have revisited the BDT, with an extended study of the input variables and hyperparameters, as discussed below. Some of the studies below are based on 75 fb^{-1} of integrated luminosity, as they were performed with a 2016+2017 analysis in mind, but their conclusions also apply to the full Run 2 analysis.

A BDT was explored using a well-known python package *xgboost* for a gradient boosted decision tree. Both ROOT-based TMVA, and python-based *xgboost* were explored, with the latter being used in the end with appropriate binning since it shows slightly higher performance. The sections will be structured as follows. A list of variables used as inputs to the BDT will be motivated and described, followed by the hyperparameters used in the training of the decision tree. The end-result comparison with the nominal cut-based analysis will be presented, and studies contributing to this result (e.g., training sets/parameters, etc) will follow.

6.2 Selection and variables

BDT training is susceptible to low statistics, so in order to increase statistics, a relaxed baseline selection for training was created, consisting of

- Lepton 1 $p_T > 15$
- Lepton 2 $p_T > 15$
- $H_T > 250$
- MET > 30
- Njets ≥ 2
- Nhtags ≥ 1

Signal was defined to be from four top MC and background was taken as all of the SM backgrounds previously described and taken in this analysis. Variations of this are discussed in the last section of miscellaneous studies.

Over 40 variables were considered. Ranked in approximate descending order of discriminative power, as reported by TMVA, they are

- Njets
- Nhtags
- Nleps
- $m_T(\ell_2)$
- $m(\ell_1, \ell_2)$
- MET
- $m_T(\ell_1)$
- H_T^b : H_T made from b jets
- $m(j_1, j_2)$: invariant mass of leading two jets
- $m(\ell_1, j_2)$

- 476 • $\Delta\phi(j_1, j_2)$
- 477 • $\Delta\phi(\ell_1, j_1)$
- 478 • $p_T(j_i)$ for $i = 1 - 8$
- 479 • q_1 : sign of the same-sign lepton pair
- 480 • $\Delta\eta(\ell_1, \ell_2)$
- 481 • H_T
- 482 • H_T^{ratio} : Ratio of H_T of first four leading jets to rest
- 483 • $m(\ell_1, j_1)$
- 484 • Nlooseb: number of btags passing loose threshold
- 485 • Ntightb: number of btags passing tight threshold
- 486 • $\max(m(j)/p_T)$: ratio of jet mass to momentum to discriminate merged jets
- 487 • Wcands: number of jet pairs with invariant mass within 30GeV of the W mass

488 Other more general variables like $m(\ell_i, j_j)$ were also considered but showed little to no discrim-
 489 ination power, where i and j encompass the first two leading objects. The same generalization
 490 applied to other indexed variables. Note that this ranking takes into account the correlation
 491 between variables, explaining why H_T is not as highly ranked, since discriminative power
 492 is first taken from Njets. Past approximately 22 variables, no extra AUC (area under curve)
 493 was gained, so it was determined to proceed with this smaller set of variables for simplicity.
 494 The AUC metric is the area under the signal versus background efficiency Receiver Operating
 495 Characteristic (ROC) curve, bounded by 0 and 1, where 1 is equivalent to perfect discrimina-
 496 tion between signal and background, and 0.5 represents discrimination no better than random
 497 guessing.

498 The 19 approximately most performant variables were then selected to continue optimization.
 499 They are

- 500 • (a) Nbtags
- 501 • (b) Njets
- 502 • (c) Nlooseb
- 503 • (d) MET
- 504 • (e) Ntightb
- 505 • (f) $p_T(\ell_2)$
- 506 • (g) $m(\ell_1, j_1)$
- 507 • (h) $p_T(j_1)$
- 508 • (i) $p_T(j_7)$
- 509 • (j) $\Delta\phi(\ell_1, \ell_2)$
- 510 • (k) $p_T(j_6)$
- 511 • (l) $\max(m(j)/p_T(j))$
- 512 • (m) Nleps
- 513 • (n) $p_T(\ell_1)$
- 514 • (o) $\Delta\eta(\ell_1, \ell_2)$
- 515 • (p) $p_T(j_8)$
- 516 • (q) H_T^b

- 517 • (r) $p_T(\ell_3)$
- 518 • (s) q_1

519 Kinematic distributions for these input variables are shown in Figure 6.

520 6.3 Hyperparameters

521 Hyper-parameter tuning was performed in order to maximize discrimination ($s/\sqrt{s+b}$ and
522 AUC) while avoiding over-training from limited statistics. The selected set of TMVA hyper-
523 parameters is given by the strings

- 524 • NTrees=500
- 525 • nEventsMin=150
- 526 • MaxDepth=5
- 527 • BoostType=AdaBoost
- 528 • AdaBoostBeta=0.25
- 529 • SeparationType=GiniIndex
- 530 • nCuts=20
- 531 • PruneMethod=NoPruning

532 In total, the complete set of combinations considered in the hyperparameter scan were

- 533 • NTrees=200,500,1000
- 534 • nEventsMin=50,150,300
- 535 • MaxDepth=4,5,6
- 536 • BoostType=AdaBoost,Bagging,Grad
- 537 • AdaBoostBeta=0.1,0.25,0.8
- 538 • SeparationType=GiniIndex,CrossEntropy
- 539 • nCuts=5,20,100

540 A similar hyperparameter scan for xgboost yielded the parameters

- 541 • n_estimators = 500
- 542 • eta = 0.07
- 543 • max_depth = 5
- 544 • subsample = 0.6
- 545 • alpha = 8.0
- 546 • gamma = 2.0
- 547 • lambda = 1.0
- 548 • min_child_weight = 1.0
- 549 • colsample_bytree = 1.0

550 Note that “n_estimators” represents the number of trees, and “eta” is the learning rate. In
551 particular, for a given learning rate, the number of trees and the depth of each tree are the most
552 impactful hyper-parameters.

553 Using these optimal hyperparameters, Figure 7 compares ROC curves and $s/\sqrt{s+b}$ curves for
554 TMVA and xgboost, showing that xgboost yields a significance performance than TMVA.




Figure 6: Distributions of kinematic inputs to the BDT. Integrals have been normalized so only shapes are relevant here.

555 Figure 8 shows that xgboost discriminator shapes for train and test sets separately for back-
 556 ground and signal. There appears to be no signs of overtraining, given the above hyperparam-
 557 eters.

558 The raw xgboost output discriminant using these 19 variables is shown in Figure 9, lumping
 559 rare MC processes into "Rares", and fakes/flips into "Others".




Figure 7: Left: ROC curves for TMVA and xgboost with AUC metric displayed in the legend. Right: Cumulative significance vs threshold value on the discriminant for TMVA and xgboost. The maximum of this curve is marked by a dotted line.





Figure 8: Raw xgboost discriminator shape for train and test sets separately for background and signal.

560 6.4 Comparison with cut-based

561 For 18 bins in total, there are 17 bins along the discriminator output from 0 to 1 corresponding
 562 to the bin edges of 0.0000, 0.0362, 0.0659, 0.1055, 0.1573, 0.2190, 0.2905,
 563 0.3704, 0.4741, 0.6054, 0.7260, 0.8357, 0.9076, 0.9506, 0.9749, 0.9884,




(a)

Figure 9: Raw xgboost BDT discriminator shape with a panel showing the cumulative $s/\sqrt{s+b}$ with MC scaled to 75 fb^{-1} .

564 0.9956, 1.0000 . The initial bin contains events matching the CRZ selection to help con-
 565 strain the $t\bar{t}Z$ background. Increasing the binning beyond this does not result in a gain in
 566 significance, and also gives issues with sample statistics.

567 Note that the binning was chosen to create a shape different from the raw xgboost discriminator
 568 output. The resulting shape is similar to the output of an AdaBoost algorithm from TMVA, and
 569 has the effect of distributing signal events normally contained within 1-2 bins into a few more,
 570 allowing for a better constraint in the fit.

571 In order to compare properly with the cut-based, the region in which the BDT discriminator is
 572 calculated and applied corresponds to the nominal baseline selection. The 18-bin BDT shown
 573 in Figure 10 yields an expected significance of 2.68σ , with the 16-bin cut-based result yielding
 574 an expected significance of 2.46σ , which is a gain slightly larger than 8%.



(a)

Figure 10: BDT discriminant in discrete signal bins, including CRZ in the first bin.

575 6.5 Misc studies, all numbers and text is from 2016

576 6.5.1 Definition of background

577 Training is performed on the full mix of backgrounds. Backgrounds modeled with simulation
 578 are normalized to a common luminosity, i.e. weighted events are used. For comparison, we
 579 also trained separate BDTs for each of the $t\bar{t}V$ backgrounds, but saw almost no gain in expected
 580 sensitivity. Note that signal-background have similar separation for each of the individual
 581 background processes, as shown in Figure 11.




Figure 11: Unstacked BDT discriminator shape for signal and background process

7 Validation of kinematic variables

In this section we describe the control-regions to validate with data the kinematic variables we use in defining our cut-based signal regions: H_T , \cancel{E}_T , N_{jets} , N_{bjets} . In addition, the variables used for the BDT are also studied: H_T^b , N_{looseb} , N_{tightb} , $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8.

The control regions considered are opposite-sign dilepton (Section 7.1), tight/loose same-sign dilepton (Section 7.2). CRW and CRZ, which are closest to the signal region, will be presented in Section 10.

7.1 Opposite-sign dilepton events

In this control region, the same requirements on H_T , \cancel{E}_T and N_{jets} as in our inclusive *baseline* selection are applied (see Section 5), but we require two opposite sign tight leptons and we remove the Z/γ^* veto. This control region coincides with the application region we use for the data-driven method to estimate the *Charge flips* background (see Section 8).

Distributions are shown in Figs. 12,13,14 (2017, 2018, 2016+2017+2018) for the main variables, and Figs. 15,16,17 for additional variables used in the BDT. The overall agreement is very good, but there are some discrepancies in the N_{bjets} and BDT distributions that we will monitor as we include additional corrections to the MC. The ISR/FSR corrections described in Section 2, corresponding to Figure 4, is applied to the $t\bar{t}$ sample here. Note that quantities like the p_T for jet 7,8 correspond to high (5+) ISR/FSR jet multiplicities in OS $t\bar{t}$. These are anyway not directly probed in the signal regions, consisting mainly of $t\bar{t}W$ and $t\bar{t}Z$ for which the matrix element provides more partons.

7.2 Same-sign tight+fail dilepton events

In this control region, the same requirements on H_T , \cancel{E}_T and N_{jets} as in our inclusive *baseline* selection are applied (see Section 5), but we require one tight lepton and one same-sign lepton failing the tight requirement. The control region is enriched in events with one fake lepton. It corresponds to the application region for the *fake-rate* method, the data-driven estimate of backgrounds with fake-leptons (see Section 8). The only difference is that in the fake-rate application we also allow events where the two leptons fail the tight requirements.

Distributions are shown in Figs. 18,19,20 (2017, 2018, 2016+2017+2018) for the main variables, and Figs. 21,22,23 for additional variables used in the BDT. There is an overall underestimate, consistent with what was seen in 2016. The underestimate is flavor-independent, and does not show large trends in the main kinematics. Such an underestimate in loose-not-tight leptons, if paired with a data FakeRate equal or larger to the MC one, can indicate that the Nonprompt lepton background is underestimated by a pure MC prediction. Since this background is predicted from data, that is not an issue with the analysis strategy.




Figure 12: Data to simulation comparisons for 2017. From left top to right bottom the H_T , E_T , N_{jets} , N_{btags} , lepton flavor and raw BDT discriminant distributions are shown for the opposite-sign dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations




Figure 13: Data to simulation comparisons for 2018. From left top to right bottom the H_T , E_T , N_{jets} , N_{btags} , lepton flavor and raw BDT discriminant distributions are shown for the opposite-sign dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations.




Figure 14: Data to simulation comparisons for 2016+2017+2018. From left top to right bottom the H_T , E_T , N_{jets} , N_{bjets} , lepton flavor and raw BDT discriminant distributions are shown for the opposite-sign dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations




Figure 15: Data to simulation comparisons for 2017, for the additional variables used by the BDT. From left top to right bottom, H_T^b , Nlooseb, Ntightb, $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the opposite-sign dilepton baseline region.




Figure 16: Data to simulation comparisons for 2018, for the additional variables used by the BDT. From left top to right bottom, H_T , Nlooseb, Ntightb, $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, \ell_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the opposite-sign dilepton baseline region.




Figure 17: Data to simulation comparisons for 2016+2017+2018, for the additional variables used by the BDT. From left top to right bottom, H_T^b , Nlooseb, Ntightb, $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the opposite-sign dilepton baseline region.




Figure 18: Data to simulation comparisons for 2017. From left top to right bottom the H_T , E_T , N_{jets} , N_{btags} , lepton flavor and raw BDT discriminant distributions are shown for the same-sign tight+fail dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations




Figure 19: Data to simulation comparisons for 2018. From left top to right bottom the H_T , E_T , N_{jets} , N_{btags} , lepton flavor and raw BDT discriminant distributions are shown for the same-sign tight+fail dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations




Figure 20: Data to simulation comparisons for 2016+2017+2018. From left top to right bottom the H_T , E_T , N_{jets} , N_{btags} , lepton flavor and raw BDT discriminant distributions are shown for the same-sign tight+fail dilepton baseline region. Shaded band shows MC stat uncertainty, except in the case of the discriminator distribution, where it has been added in quadrature with scale, btag, JEC, and JER variations




Figure 21: Data to simulation comparisons for 2017, for the additional variables used by the BDT. From left top to right bottom, H_T , $N_{\text{loose b}}$, $N_{\text{tight b}}$, $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, \ell_2)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the same-sign tight+fail dilepton baseline region.




Figure 22: Data to simulation comparisons for 2018, for the additional variables used by the BDT. From left top to right bottom, H_T , N_{looseb} , N_{tightb} , $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the same-sign tight+fail dilepton baseline region.




Figure 23: Data to simulation comparisons for 2016+2017+2018, for the additional variables used by the BDT. From left top to right bottom, H_T^b , N_{looseb} , N_{tightb} , $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1,6,7, and 8; shown for the same-sign tight+fail dilepton baseline region.

617 7.3 Fake-enriched validation region in data

618 In this control region, the same requirements on H_T , E_T and N_{jets} as in our inclusive *baseline* se-
 619 lection are applied (see Section 5), except we relax the H_T requirement and require $N_{\text{bjets}} = 1$.
 620 This region has a significant non-prompt component and allows us to check the overall clo-
 621 sure of the method in data. In the plots, the fake background is data-driven. Distributions are
 622 shown in Figs. 24. The overall data/MC normalization factor in this region is 1.06. If fakes are
 623 entirely responsible for this discrepancy, and given that fakes constitute half of the background,
 624 this represents a 12% normalization increase of fakes, well within the 30% normalization un-
 625 certainty taken on this process.




Figure 24: Data to simulation comparisons for 2016+2017+2018. From left top to right bottom the H_T , E_T , N_{jets} , N_{bjets} , lepton flavor and raw BDT discriminant distributions are shown for the same-sign dilepton region with $N_{\text{bjets}} = 1$, $N_{\text{jets}} \geq 2$, $E_T^{\text{miss}} \geq 50$. Shaded band shows MC stat uncertainty.

8 Background Estimations

Backgrounds for the same-sign dilepton final state can be divided in three categories:

- **Fake leptons:** “Non-Prompt” or “Fake” leptons are leptons from heavy-flavour decays, misidentified hadrons, muons from light-meson decays in flight, or electrons from unidentified photon conversions. Depending on the signal regions, this background is dominated by $t\bar{t}$ or W +jets processes.
- **Charge flips:** Charge misidentification, i.e. events with opposite-sign isolated leptons where the charge of one of the leptons is misidentified because of severe bremsstrahlung in the tracker material. This background, is relevant only for electrons and is negligible for muons.
- **Rare SM processes:** Rare SM processes yield same-sign leptons, mostly from $t\bar{t}W$ and $t\bar{t}Z$. We also include the contribution from the SM Higgs boson produced in association with a vector boson or a pair of top quarks in this category of background. With the exception of $t\bar{t}Z$ and $t\bar{t}W$, rares are estimated from simulation.

8.1 Fake leptons

The fake lepton prediction is determined as described in the SUSY AN [1] using the SUSY same-sign analysis baseline selection. The description is also included below, for convenience.

The uncertainties related to this estimate are discussed in Section 9.3. Note that in the case of 0 events in the application region as an input to the extrapolation into the signal regions, the prompt-nonprompt events in simulation are multiplied by the data fake rate to obtain a prediction with associated error. This error is then set as the statistical upper limit on the data-driven prediction of 0.

8.1.1 The fake rate method

Background from fake leptons is estimated with the “fake rate” method. The number of events in the sample with at least one lepton that passes a loose selection but fails the full set of tight identification and isolation requirements (application region) is weighted using the “tight-to-loose” ratio, i.e. the probability that a loosely identified non-prompt lepton also passes the full set of requirements. This probability is measured as a function of lepton p_T and η , as well as event kinematics, in a control sample of QCD multijet events that are enriched in non-prompt leptons (measurement region). Such region is triggered by the auxiliary triggers defined in Table 4 and requires only one denominator lepton in the event, one recoiling jet with $\Delta R(jet, lep) > 1.0$ and low MET and MT to suppress the contribution from W and Z . The main systematic effects are the non-universality of the “tight-to-loose” ratio, particularly due to the dependency from the mother parton p_T and the flavor composition of the sample, and the prompt contribution in the measurement region.

In the SUSY same-sign 13 TeV analyses, developments were deployed [9][17] in the fake rate estimation in order to reduce the dependency to the mother parton p_T by using a new proxy of the lepton p_T :

$$\begin{aligned} \text{if } p_T^{rel} > I_3 : \quad p_T &\rightarrow p_T \cdot (1 + \max(0, I_m - I_1)) \\ \text{else : } \quad p_T &\rightarrow \max(p_T, p_T(jet)) \cdot I_2 \end{aligned} \quad (5)$$

In addition to the corrected lepton p_T definition, another development has been introduced since [17]: for the electrons, the flavor dependency of the tight-to-loose ratio is reduced by

666 extrapolating on both isolation and lepton MVA ID. These improvement are included in the
667 2016 and in the current (2016+2017) $t\bar{t}$ analyses.

668 8.1.2 Fake Rate measurement

669 We derive different versions of the fake rate for muons and electrons, collected with triggers
670 with an isolation requirement. The numerator and denominator selections are defined as in
671 section 4.5.

672 Events for fake rate measurement need to pass the following requirements:

- 673 • pass a specific auxiliary trigger, described in Table 4
- 674 • only one denominator lepton (FO)
- 675 • at least one jet with $\Delta R(\text{jet}, \text{FO}) > 1$
- 676 • $E_T^{\text{miss}} < 20 \text{ GeV}$, $M_T < 20 \text{ GeV}$.

677 The requirements on E_T^{miss} and M_T are intended to suppress the contribution from prompt
678 leptons in the measurement region. Such contribution is subtracted from the fake rate using
679 DY, WJets, and $t\bar{t}$ Monte Carlo samples. For the 2016 data, these samples were normalized in
680 the control region defined with $E_T^{\text{miss}} > 20 \text{ GeV}$, $70 < M_T < 120 \text{ GeV}$ and a tight lepton. For
681 the 2017 data, due to the increased PU contribution, an improved normalization technique is
682 used, based on template fits of the full M_T distribution. The M_T distribution in data with
683 $E_T^{\text{miss}} > 30 \text{ GeV}$ and lepton $p_T > 20 \text{ GeV}$ is fitted with the sum of two templates derived
684 from MC, one for QCD and one for the sum of the electroweak processes (DY, WJets and $t\bar{t}$).
685 The normalization of the electroweak processes is extracted from the fit and half the difference
686 between the normalization and unity is taken as an uncertainty. As the shape of the non-prompt
687 component may not be well-modeled, we repeat the fit replacing the QCD MC template with
688 a data-driven template extracted from events failing the isolation cut. The difference between
689 the two fits is taken as an additional uncertainty in the normalization, added in quadrature
690 with the uncertainty obtained from the fit above.

691 The nominal and alternative fits for electrons and muons are shown in Figure 25 (2017) and
692 Figure 26 (2018), and the resulting normalization corrections for the electroweak samples are
693 shown in Table 12. Statistical uncertainties on the measurement are assumed to be negligible.
694 Results with 35.9 fb^{-1} of 2016 data are shown in [1], while the 2017 and 2018 results with
695 41.5 fb^{-1} and 59.6 fb^{-1} of data are shown in Figs. 27-29, 28-30 below.

	template	isolated		non-isolated	
		e	μ	e	μ
2017	MC	1.215	1.222	1.208	1.202
	data	1.277	1.195	1.298	1.178
2018	MC	1.200	1.283	1.202	1.288
	data	1.268	1.252	1.297	1.250

Table 12: Normalization scale factors for electroweak samples derived with two different M_T templates for QCD: MC and data (the data template refers to the inverted isolation region).

696 8.1.3 Fake Rate closure in MC: $t\bar{t}$ and W+jets: 2017/2018 samples

697 Using these definitions, we tested the closure of the method in the baseline and signal regions
698 and, inclusively in lepton p_T , for the most relevant kinematic distributions in 31 and 32. the
699 level of closure obtained is typically at 30% or better, similar to the 2016 one. The closure in the
700 electron channel showed a potential trend at high p_T , with deviations up to 60%.




Figure 25: Isolated triggers, 2017 Data and MC: Fitted M_T distribution used to derive the normalization of electroweak samples (DY, WJets, $t\bar{t}$) in the fake rate measurement region. Electrons are shown on the left, muons on the right. From top to bottom, the results from the nominal selection ($E_T^{\text{miss}} > 30$ GeV and lepton $p_T > 20$ GeV) with the QCD MC template and alternative data QCD template are shown.




Figure 26: Isolated triggers, 2018 Data and MC : Fitted M_T distribution used to derive the normalization of electroweak samples (DY, WJets, tt) in the fake rate measurement region. Electrons are shown on the left, muons on the right. From top to bottom, the results from the nominal selection ($E_T^{\text{miss}} > 30$ GeV and lepton $p_T > 20$ GeV) with the QCD MC template and alternative data QCD template are shown.




Figure 27: Isolated triggers: Fake rate for muons (left) and electrons (right) for: 2017 data uncorrected (top), 2017 data corrected for EWK contribution (middle) and 2017 QCD MC (bottom). Uncertainties are only statistical.




Figure 28: Isolated triggers: Fake rate for muons (left) and electrons (right) for: 2018 data uncorrected (top), 2018 data corrected for EWK contribution (middle) and 2017 QCD MC (bottom). Uncertainties are only statistical.




Figure 29: EWK-corrected data fake rate projected vs p_T for 2017 data (black) and 2017 QCD MC (red), for muons (left) and electrons (right). The shaded band in the projection is the systematic uncertainty related to the EWK contamination.




Figure 30: EWK-corrected data fake rate projected vs p_T for 2018 data (black) and 2017 QCD MC (red), for muons (left) and electrons (right). The shaded band in the projection is the systematic uncertainty related to the EWK contamination.




Figure 31: 2017 MC: Electron+muon fake rate closure for QCD measurement in MC soup.




Figure 32: 2018 MC: Electron+muon fake rate closure for QCD measurement in MC soup. **QCD Fake rate to be updated. 2018 bcToE QCD MC is not available yet**

8.2 Charge misidentification

The charge flip prediction is determined as described in the SUSY AN [1] using the SUSY same-sign analysis baseline selection. The description is also included below, for convenience. The uncertainties related to this estimate are discussed in Section 9.3.

The background due to charge flips is estimated by selecting opposite-sign ee or $e\mu$ events passing the full kinematic selection and then weighting them by the p_T and η -dependent probability of electron charge mismeasurement

This probability, shown in Fig. 33, is obtained from a soup of $T\bar{T}$ and DY simulation and is then validated with a control data sample of same-sign $Z \rightarrow ee$ events, using a $E_T^{\text{miss}} < 50$ GeV requirement to be orthogonal to the signal region. The level of agreement in this control region is used to gauge the associated systematic uncertainty and to derive a correction to the MC-based rate estimation. In the 2016 data, we find good agreement between prediction and data in the control region [1]. In the 2017 and 2018 data, the MC Flip Rate is significantly lower than the 2016 one due to the upgraded pixel detector. However, the prediction in the same-sign $Z \rightarrow ee$ region is found to only be about 30% lower than the observed number of events in this region, as shown in Figure 35. Consequently, the 2017 and 2018 charge misidentification prediction are scaled by nearly 50%, as seen in Figure 34. The yearly scaling is given in Table 13. Since we do not find significant trends in the lepton kinematics, so we do not consider p_T and η -dependent corrections. In addition to the statistical uncertainties, we apply a 20% systematic uncertainty on this background prediction for all years. In MC, the flip rate for muons is found to be $O(10^{-6})$ and is therefore neglected.

year	obs/pred
2016	1.01
2017	1.44
2018	1.41

Table 13: Ratio of observed flip rate in data to the flip rate in simulation. These are the multiplicative correction to the MC-based charge flip probabilities.




Figure 33: Electron charge flip rate for 2016, 2017, and 2018.




Figure 34: Predicted and observed lepton p_T (left) and η (middle) and $m_{\ell\ell}$ (right) in a same-sign $Z \rightarrow ee$ peak for years 2016, 2017, and 2018 from top to bottom. The prediction is normalized to the observed data.




Figure 35: Electron charge flip rate in simulation and data as a function of time

722 **8.3 Rare SM processes**

723 Rare SM processes that result in the production of same-sign leptons are estimated using MC
 724 samples. The processes with the largest contributions, $t\bar{t}W$, $t\bar{t}Z$ and $t\bar{t}H$, are treated individually
 725 and assigned separate uncertainties. Processes with smaller contributions, including diboson
 726 (WZ , ZZ) and triboson (WWW , WWZ , WZZ , ZZZ), Higgs (HZZ , VH), same-sign WW from
 727 both single ($qqWW$) and double-parton scattering (DPS WW), rare top channels (tZq), and
 728 $t\bar{t}t+W, t\bar{t}t+j$ are grouped in a single category termed "Rare SM" in all plots. Similarly, processes
 729 where one of the leptons is an electron from an unidentified photon conversion are grouped
 730 in a category called " $X+\gamma$ " in all plots: these include $W\gamma$, $Z\gamma$, $t\bar{t}\gamma$ and $t\gamma$. Both the "Rare SM"
 731 and " $X+\gamma$ " categories are assigned a large ($\pm 50\%$) theoretical normalization uncertainty. The
 732 breakdown of the individual processes of the "Rare SM" category is shown in Figure 36 The
 733 theory uncertainties considered are described in more detail in Sec. 8.4.

734 In addition to the theoretical uncertainties, all samples are assigned uncertainties based on
 735 reconstruction, as summarized in Section 9, due to JES, b-tagging, lepton and trigger scale
 736 factors uncertainties and luminosity. Finally, the statistical uncertainty of the MC samples is
 737 also taken into account.




Figure 36: Relative composition of multi-top (left) and multi-boson (right) rare backgrounds in the signal regions for all MC.

738 8.4 Theoretical uncertainties on TTW, TTZ and TTH

739 For the largest SM backgrounds, we carefully assign uncertainties on the overall normalization
740 and on the shape across signal regions.

741 8.4.1 Normalization

742 For $t\bar{t}W$ and $t\bar{t}Z$, a 40% normalization uncertainty is applied, but the results are not sensitive to
743 this initial choice, because the respective control regions included in the maximum likelihood
744 fit, discussed in more detail in Sec. 11, are used to further constrain the normalization of these
745 backgrounds.

746 For $t\bar{t}H$, the normalization uncertainty used in the 2016 analysis was 50%, to cover the 1.5
747 signal strength observed in the results of HIG-17-004. For the full Run2 analysis, we rely on the
748 updated measurement of HIG-17-035, which finds a signal strength of $1.26^{+0.31}_{-0.26}$. We therefore
749 apply a 25% normalization uncertainty.

750 8.4.2 Shape

751 To evaluate the theory uncertainties on the shape, we explore scale and PDF variations, as well
752 as differences between LO and NLO samples, where both are available. These studies were
753 performed with 2016 MC.

754 The PDF shape uncertainties for these samples are generally smaller than the MC statistical
755 uncertainties on the background yield in each region, so no additional shape uncertainty is
756 assigned to account for them.

757 The NLO scale uncertainties for these samples are fairly stable at around 15% for $t\bar{t}W$, $t\bar{t}Z$, and
758 $t\bar{t}H$, as shown in Figures 38, 37, 39. With the exception of $t\bar{t}H$, these figures also show the LO
759 scale uncertainty, which covers well the difference between LO and NLO. The LO agreement
760 with the NLO shapes, within LO scale uncertainty, gives confidence in the NLO scale uncer-
761 tainties as they are evaluated using the same variations of renormalization and factorization
762 scales.




Figure 37: Comparison of LO and NLO kinematics for $t\bar{t}Z$ with scale uncertainties in the baseline region and statistical uncertainties on NLO. The error bars on the NLO (black) are the NLO uncertainties, while the error bars on the LO (red) are the LO uncertainties. The gray band is the stat uncertainty on the NLO




Figure 38: Comparison of LO and NLO kinematics for ttW with scale uncertainties in the baseline region and statistical uncertainties on NLO. The error bars on the NLO (black) are the NLO uncertainties, while the error bars on the LO (red) are the LO uncertainties. The gray band is the stat uncertainty on the NLO




Figure 39: ttH NLO with NLO scale uncertainties in the baseline region. The gray band is the stat uncertainty on the NLO

9 Systematic Uncertainties

763

764 With respect to the 2016 SUSY same-sign analysis [1], the 2016 tttt analysis only changed un-
 765 certainties on $t\bar{t}Z$, $t\bar{t}H$, and $t\bar{t}W$, including also uncertainty for $t\bar{t}W+bb, t\bar{t}Z+bb$, JER, and adding
 766 ISR/FSR variations from dedicated samples. With respect to the 2016 tttt analysis [4], the cross-
 767 section for $t\bar{t}H$ has a 25% uncertainty taken as a systematic. Other uncertainties, which have
 768 changed as a result of new measurements, are marked in blue if the change is also applied
 769 to 2016 data. All uncertainties marked in black have remained unchanged since 2016. These
 770 changes are reflected in Table 14.

771

9.1 Correlation model for 2016+2017+2018

772

773

774

775

In order to combine the 2016+2017+2018 datasets, we have to make assumptions about the
 correlation of uncertainties in the different years. Since the analysis is statistically dominated,
 these correlations are expected to have a very small influence. The current model we have
 chosen makes the following assumptions,

776

777

- Statistical uncertainties (FO CR stat, OS CR stat, MC stat): uncorrelated
- Luminosity, JEC, JER, b-tag, PU, lepton eff., HLT, eff.: uncorrelated (derived from

source	magnitude	effect on yield
luminosity	2.5, 2.3, 2.5%	2.5, 2.3, 2.5% (2016-2018)
JES	1 – 8%	1 – 15%
JER	1 – 10%	1 – 10%
$t\bar{t}W, t\bar{t}Z$ ISR/FSR	< 15%	1 – 8%
$t\bar{t}W/Z+bb$	< 30%	< 15%
b-tag efficiency	~ 1 – 20%	1 – 15%
pileup	5%	0-5%
lepton efficiencies	2 – 5% (leg)	2 – 10%
HLT efficiencies	1 – 5% (leg)	2 – 7%
FO CR stat.	1 – 100%	1 – 100% (fake bkg. only)
FR extrapolation	30-60%	30-60% (fake bkg. only)
EWK subtraction in FR	25-50% (FR)	1 – 30% (fake bkg. only)
OS CR stat.	4 – 100%	4 – 100% (charge misId. only)
charge misId.	20%	20% (charge misId. only)
$t\bar{t}W$ norm.	40%	40 % ($t\bar{t}W$ only)
$t\bar{t}Z$ norm.	40%	40 % ($t\bar{t}Z$ only)
$t\bar{t}H$ norm.	25%	25% ($t\bar{t}H$ only)
MonteCarlo stat.	1 – 25%	1 – 25%
QCD scales and PDFs	$\times 0.5 / \times 2$	10 – 20% ($t\bar{t}W, t\bar{t}Z, t\bar{t}H$)
other bkgs.	50%	50% (Rare and $X + \gamma$)

Table 14: Summary of the sources of uncertainties, their magnitude and their effects. The second column indicates the magnitude of the yield variation. Reported uncertainties are representative for the most relevant signal regions.

778 different datasets)

- 779 • ISR/FSR: uncorrelated (derived from different datasets and with respect to different
780 MC samples)
- 781 • ttW/Z+bb, ttH normalization: correlated (based on theory predictions and CMS
782 measurements)
- 783 • ttW and ttZ normalizations: correlated (based on theory predictions and consistent
784 Control Regions)
- 785 • Rare and X+ γ normalizations: correlated (based on theory predictions)
- 786 • QCD scale and PDF uncertainties: correlated (based on theory predictions)
- 787 • EWK subtraction in FR: uncorrelated (based on differently prescaled triggers in data)
- 788 • FR extrapolation: correlated (method is the same, closure in MC shows the same
789 features)
- 790 • Charge misid.: uncorrelated (agreement in validation region is quite different, prob-
791 ably due to different Pixel detector conditions)

792 We ran limits under three correlation assumptions with 18 cut-based bins. With respect to
793 the nominal correlation model, fully uncorrelating (correlating) nuisances between the 3 years
794 decreases (increases) the expected significance by 2%.

795 9.2 Experimental sources of uncertainties

796 One of the main experimental sources of uncertainty is the knowledge of the jet energy scale
797 (JES), affecting all the simulated backgrounds and considered signals. The 13 TeV uncertainties
798 vary the jet energy scale by 1–8%, depending of the transverse momentum and pseudorapidity
799 of the jet. The impact of these uncertainties is assessed by shifting the jet energy correction
800 factors for each jet up and down by $\pm 1\sigma$ before the calculation of all kinematic quantities. The
801 variations are correlated among the different signal regions as bin-by-bin migration is allowed.
802 Variations are asymmetric in nature and are used as asymmetric nuisance parameters in the
803 result interpretation. The JES uncertainties are propagated to the missing transverse energy
804 and all jet-related variables (number of jets, H_T , number of b-jets) used in this analysis. As
805 some of the simulation samples are statistically limited, the size of the JES uncertainties can
806 reach high values in several regions not well populated by several processes. As usually those
807 variations impact mostly background samples that do not contribute much to those regions,
808 we consider large variations as they are obtained out of the box. Most populated signal regions
809 shows yield variations of 8% when the jet energy scale is varied by one standard deviation.

810 The uncertainty related to the knowledge of the jet energy resolution (JER) can be considered
811 as well for the simulated backgrounds and the signals. The effect of smearing jets on the accep-
812 tance of events is assessed in the same manner as the JES uncertainty.

813 A similar approach is used for the uncertainties associated to the corrections for the b-tagging
814 efficiencies for light and bottom flavor jets, which are parametrized as a function of p_T , η .
815 The variation of the scale factor is at maximum of the order of 1–20% per jet, and lead to an
816 overall effect on yield included between 1 and 15% depending on the signal region and on the
817 topology of the events included in those signal regions. If considering only highly populated
818 signal regions to get an overview of the main effects on the background yields, the bulk of the
819 ttW and ttH yield varies by $\sim 8\%$ and the ttZ yield by $\sim 6\%$.

820 When applying b-tagging efficiency scale factors, it is possible to apply period-dependent scale
821 factors, rather than an average scale factor for the entire 2016 dataset. This is relevant for signal

822 regions where several mistags from light jets are expected, since the mistag SFs for light jets
 823 are time-dependent. When checking the flavor composition of our main background $t\bar{t}W$, as in
 824 Figure 40, we find that none of our $N_{b \text{ jets}}$ bins are dominated by events with multiple light jets,
 825 so the period-dependent scale factors are not necessarily warranted. As expected, when we do
 826 apply them, we find negligible differences in our background predictions with respect to the
 827 nominal scale factors. For 2017, we apply period-dependent b-tagging SFs.

828 Events from processes such $t\bar{t}W$ and $t\bar{t}Z$, which have two b quarks from the decay of the two
 829 top quarks, primarily enter regions requiring three or more b-tagged jets by either the mis-tag
 830 of a charm quark from the hadronic decay of a W boson or from the production of additional
 831 heavy flavor, primarily from an ISR or FSR gluon which splits to a $b\bar{b}$ pair.

832 In 2016, events in simulation containing additional HF quarks not from top or W decays were
 833 scaled up to account for $SF_{\sigma(t\bar{t}b\bar{b})/\sigma(t\bar{t}jj)} \sim 1.7 \pm 0.6$ as measured by TOP-16-010. In the high
 834 $N_{b \text{ tag}}$ signal region bins where this effect is dominant, this resulted in a systematic uncertainty
 835 up to 20%. In 2017 and 2018, we apply the same correction as 2016 onto $t\bar{t}$, $t\bar{t}W$, $t\bar{t}Z$, and $t\bar{t}H$, as
 836 it is observed to bring agreement in the high statistics opposite-sign $t\bar{t}$ control region.




Figure 40: Flavor composition breakdown of the $t\bar{t}W$ sample, in bins of number of b-tagged jets for all MC

837 Trigger efficiency scale factors obtained from data are applied to correct the simulation.

838 Lepton efficiency scale factors [13], accounting for differences between the Data and MC for
 839 reconstruction and identification of electrons and muons, are applied to all MC events. They
 840 result in uncertainties of approximately 2-3% for muons and 3-5% for electrons for 2016.

841 The simulation is reweighted to match the expected data distribution in the number of collision
 842 per events; the uncertainty on the minimum bias cross-section is propagated to the final yields
 843 with an effect at the level of 4% or less across the three years.

844 The $t\bar{t}W$ and $t\bar{t}Z$ simulation is reweighted to match the number of additional ISR/FSR jets
 845 observed in data, as described in Section 2. An uncertainty equivalent to 50% of the difference
 846 between 1 and the reweighting factor is applied to this procedure. In 2016, since the reweighting
 847 factor can be as large as 0.77, this results in an uncertainty of 15% on the highest $N_{b \text{ jets}}$ signal
 848 regions. In 2017 and 2018, the reweighting factor is as large as 1.4, which results in a similar
 849 uncertainty to 2016 on the highest $N_{b \text{ jets}}$ signal regions.

9.3 Uncertainties from data-driven background estimations

The following uncertainties are defined for the data-driven background estimations, based on the statistics of the control region and the extrapolation from control to signal region.

For the nonprompt lepton and charge misidentified lepton backgrounds, the statistical uncertainty from the control sample varies greatly depending on the signal region considered. Both the flip rate and the fake rate are smaller than 1, so the statistical uncertainty is always smaller (or much smaller, for charge misidentification) than the size of the predicted background. In addition to the statistical uncertainty, the nonprompt lepton background is assigned an overall normalization uncertainty of 30%, based on a comparison of the non-closure of two alternative methods (nominal fake-rate and in-situ fake-rate). This uncertainty is increased to 60% for electrons with $p_T > 50 \text{ GeV}$, to account for the trends observed at high p_T in the electron closure for both methods. On top of the uncertainties mentioned above, the nonprompt lepton background prediction also includes an uncertainty related to the electroweak contamination subtraction in the region where the fake rate is computed. To evaluate this uncertainty the fake-rate is computed based on varying the size of the electroweak contamination, and the effect is propagated through the whole analysis. The overall effect on the nonprompt lepton background yield lies between 1 and 50% depending of the signal region considered. The charge misidentified lepton background is assigned a systematic uncertainty of 20%.

For $t\bar{t}W$ and $t\bar{t}Z$, the respective control regions are used only to set the overall normalization, while the shape is taken from simulation with the experimental uncertainties defined above and the theoretical uncertainties described in section 8.4. The two control regions are included in the fit with a 40% normalization nuisance parameter, and the fit constrains this parameter to approximately 20% given the statistics of the control region, where the corresponding post-fit constraint was 30% in the 2016 analysis.

9.4 Systematic uncertainties from statistical sources

The statistical precision given by the Monte Carlo samples has to be taken into account for all rare SM processes. Within the framework of the HiggsCombine tool, we use the Barlow-Beeston [21] method of handling both MC statistics and data-driven background statistics (for the nonprompt and misidentified-charge backgrounds) via the “autoMCStats” parameter in the datacard.

9.5 Signal uncertainties

The $t\bar{t}t$ signal is assigned uncertainties based on all the effects described above. The uncertainty values are summarized in Table 15. For the experimental uncertainties we report the range of their effect across SRs, while for the theoretical uncertainties we separate the correlated (acceptance*efficiency, $\mathcal{A}\epsilon$) and uncorrelated (shape) effects. The $\mathcal{A}\epsilon$ of our SR selection, including branching ratio, is $1.5^{+0.02}_{-0.03}\%$, where the uncertainties represent the QCD scale variations. The effect of QCD scale variations on the shape of $t\bar{t}t$ within the set of SRs is shown in Figure 41: is as large as 10%. PDF uncertainties on acceptance and shape are both negligible as they are smaller than 1%

Uncertainties for ISR and FSR variations during sample generation for $t\bar{t}t$ are assessed by considering 4 additional samples and are shown in Figure 42. The variation in acceptance (shape) is 1% (<8%) for ISR and 6% (<10%) for FSR.

No uncertainty is assigned to the cross-section itself, since we are measuring this process.




Figure 41: QCD scale uncertainties on the shape of the $t\bar{t}t$ signal.

source	magnitude	effect on yield
jet ES	1 – 8%	1 – 15%
JER	1 – 10%	1 – 10%
b-tag efficiency	~ 1 – 20%	1 – 15%
pileup	5%	1 – 4%
ISR	-	<8%
FSR	-	<10%
lepton efficiencies	2 – 5% (leg)	2 – 10%
HLT efficiencies	1 – 5% (leg)	2 – 7%
MonteCarlo stat.	1 – 100%	< 8%
QCD scales (acceptance)	$\times 0.5 / \times 2$	2%
QCD scales (shape)	$\times 0.5 / \times 2$	1-10%
PDFs (acceptance)	Envelope	$\leq 1\%$
PDFs (shape)		$\leq 1\%$
α_S (acceptance)		$\leq 6\%$
α_S (shape)		$\leq 1\%$

Table 15: Summary of the sources of uncertainties for $t\bar{t}t$ signal.




Figure 42: Effect of ISR (left) and FSR (right) variations on the $t\bar{t}$ signal.

9.6 Summary of theoretical uncertainties on MC-based processes

Table 16 shows a summary of the cross-section, scale, and pdf uncertainties on the MC-based processes, including signal, indicating whether the nuisance contains an acceptance and/or shape component. Associated with each row, there are 3 nuisances, parameterized as one log-normal, and two shape nuisances. Some notes about individual processes follows.

- Signal $t\bar{t}\bar{t}$ has no log-normal nuisance associated with cross-section uncertainty because it is being measured.
- Both $t\bar{t}W$ and $t\bar{t}Z$ do not have an acceptance component in the scale/pdf variations because the 40% log-normal is taken to generally cover cross-section \times acceptance as they are constrained by dedicated control regions.
- $t\bar{t}H$ has a 25% cross-section log-normal to reflect the result of HIG-17-035 as mentioned in Section 8.4.
- For $X+\text{Gamma}$ and $t\bar{t}VV$, the largest cross-section uncertainty on any constituent subprocess is taken on the whole process. Thus, $x+\text{gamma}$ obtains an 11% normalization uncertainty driven by $t\bar{t} + \gamma$ (+9.9%, -11.2%), and $t\bar{t}VV$ obtains a 11% normalization uncertainty from $t\bar{t}WW$ (+8.6% -11.3%).
- For rares, we obtain uncertainties directly from MC samples by calculating the summed yield of rares for nominal variation, scale up, scale down, pdf up, and pdf down. Adding scale and pdf variations in quadrature yields a 20% variation for 2016 and 2017 samples. We proceed with a log-normal of 20% on the sum of these processes.

	lnN	scale		pdf	
		acceptance	shape	acceptance	shape
$t\bar{t}\bar{t}$		x	x	x	x
$t\bar{t}W$	1.40		x		x
$t\bar{t}Z$	1.40		x		x
$t\bar{t}H$	1.25	x	x	x	x
$X+\gamma$	1.11	x	x	x	x
$t\bar{t}VV$	1.11	x	x	x	x
Rares	1.20	x	x	x	x

Table 16: Summary of the theory uncertainties

913 10 Kinematics

914 Prefit kinematic distributions of the 2017 and 2018 data events passing the baseline selection
 915 and falling into the $t\bar{t}W$ control region are shown in Figs. 43-45. Similarly, distributions of
 916 events falling into the $t\bar{t}Z$ control region are shown in Figs. 47-49. The background prediction
 917 for the sum of signal regions is shown in Figs. 51-53.

process	SF (2016)	SF (2017)	SF (2018)	SF (Run2)
ttz	1.58	1.005	1.207	1.258
ttw	1.347	1.35	1.156	1.299
tth	1.087	1.089	1.045	1.088
tttt	1.175	0.845	1.451	1.053
fakes	1.064	1.163	1.081	1.125
xg	1.06	1.035	1.015	1.014
rares	1.055	1.017	1.023	1.017
ttvv	1.028	1.018	1.02	1.011
flips	1.016	1.007	0.999	1.001

Table 17: Postfit/prefit scale factors with the BDT analysis. These are the ratio of the normalizations before and after the fit, and they are impacted by all nuisance parameters affecting each background. For this reason, they are not necessarily expected to be consistent between years, as are the individual nuisance parameters themselves (shown in Appendix A).




Figure 43: 2016 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}W$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and raw BDT discriminant where the last bin includes the overflow.




Figure 44: 2017 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}W$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and raw BDT discriminant where the last bin includes the overflow.




Figure 45: 2018 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}W$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and raw BDT discriminant where the last bin includes the overflow.




Figure 46: Run2 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}W$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and raw BDT discriminant where the last bin includes the overflow.




Figure 47: 2016 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}Z$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, raw BDT discriminant, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 48: 2017 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}Z$ control region: H_T , E_T^{miss} , N_{jets} , $N_{\text{b jets}}$, lepton flavor, raw BDT discriminant, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 49: 2018 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}Z$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, raw BDT discriminant, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 50: Run2 data and predictions: Prefit distributions of the main analysis variables in the $t\bar{t}Z$ control region: H_T , E_T^{miss} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, raw BDT discriminant, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 51: 2016 predictions: profit distributions of the main analysis variables in the sum of signal regions: H_T , E_T^{miss} , M_T^{min} , N_{jets} , $N_{\text{b jets}}$, lepton flavor, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 52: 2017 predictions: profit distributions of the main analysis variables in the sum of signal regions: H_T , E_T^{miss} , M_T^{min} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 53: 2018 predictions: profit distributions of the main analysis variables in the sum of signal regions: H_T , E_T^{miss} , M_T^{min} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 54: run2 predictions: profit distributions of the main analysis variables in the sum of signal regions: H_T , E_T^{miss} , M_T^{min} , N_{jets} , $N_{b \text{ jets}}$, lepton flavor, and $m_{\ell\ell}$ of the OSSF pair, where the last bin includes the overflow.




Figure 55: Same as Figure 54 but with stacked signal, assuming signal strength of 1.

11 Results

Results for the 2016+2017+2018 (“Run 2”) data are obtained with the data in “golden JSON” given in Sec.2 , for a total luminosity of 77.4 fb^{-1} . The expected yields in the signal regions as well as the $t\bar{t}W$ and $t\bar{t}Z$ control regions, and BDT regions are shown in Figure 56 for prefit for 2016/2017/2018/Run2. The postfits plot for the two analyses are shown in Figure 57. The 2016,2017,2018,Run 2 cut-based numerical yields can be found in Table 18,19,20,21 for prefit, and analogous tables for the BDT analysis can be found in Table 22,23,24,25. Postfit tables for Run 2 cut-based and BDT can be found in Table 26 and Table 27, respectively. . Note that with respect to 2016, also the theoretical cross section assumed for the $t\bar{t}t\bar{t}$ process has changed, from 9.2 to 11.96 fb.

A binned likelihood fit is performed over signal regions using the Higgs Combine tool [22]; exclusion limits at 95% CL are calculated with the Asymptotic CLs method.

For reference, the commands used to extract upper limits, significances, measurement values, and obtain the NLL vs μ , respectively, are:

```

• combine -M Asymptotic card.txt --noFitAsimov
• combine -M Significance card.txt -t -1 --expectSignal=1 --significance
• combine -M FitDiagnostics card.txt -t -1 --expectSignal=1 --robustFit=1
  --saveShapes --saveOverallShapes --saveWithUncertainties -n name
• combine -M MultiDimFit card.txt -t -1 --expectSignal=1 --algo grid
  --centeredRange=2.0 --saveFitResult --redefineSignalPOI r --robustFit=1
  --saveNLL

```

`-t -1 --expectSignal=1` and `--noFitAsimov` are omitted for observed results.

Using the Run2 data, the *cut-based* analysis sets an observed (expected) upper limit on the production cross section of 20.04 fb ($9.35^{+4.29}_{-2.88} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 1.712 (2.478) standard deviations, corresponding to a measured observed (expected) signal strength parameter of $0.784^{+0.514}_{-0.469}$ ($1.000^{+0.480}_{-0.433}$)

The *BDT* analysis sets an observed (expected) upper limit on the production cross section of 22.51 fb ($8.46^{+3.91}_{-2.57} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 2.561 (2.699) standard deviations, corresponding to a measured observed (expected) signal strength parameter of $1.052^{+0.483}_{-0.437}$ ($1.000^{+0.442}_{-0.401}$)

The likelihood scan for Run2 is shown in Fig. 58.

Several interpretations of these results are presented in the following sections

- Section 12.1: 2HDM with a heavy scalar or pseudoscalar boson decaying to on-shell $t\bar{t}$
- Section 12.2: top yukawa coupling constant
- Section 12.3: off-shell top-philic scalar or pseudoscalar boson
- Section 12.4: EFT oblique Higgs parameter \hat{H}
- Section 12.5: Simplified dark matter with scalar or pseudoscalar mediator




Figure 56: Data yields compared to prefit for cut-based (left) and BDT (right) analyses separately for data periods 2016, 2017, 2018, Run2 from top to bottom.




Figure 57: Data yields compared to postfit for cut-based (left) and BDT (right) analyses.

Table 18: Prefit event yields in SR+CR regions for 2016.

	tt̄W	tt̄Z	tt̄H	tt̄V	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	tt̄t̄
CRZ	0.50 ± 0.18	14.01 ± 5.20	0.69 ± 0.18	0.50 ± 0.06	0.06 ± 0.01	2.79 ± 0.58	0.02 ± 0.00	0.42 ± 0.17	18.98 ± 5.34	31	0.25 ± 0.04
CRW	22.56 ± 8.01	6.31 ± 2.31	8.44 ± 2.14	1.76 ± 0.21	4.41 ± 0.71	2.24 ± 0.49	4.95 ± 0.95	10.05 ± 3.82	60.72 ± 9.61	82	1.23 ± 0.10
SR1	1.87 ± 0.70	0.68 ± 0.26	1.12 ± 0.31	0.27 ± 0.04	0.18 ± 0.07	0.21 ± 0.06	0.12 ± 0.02	0.70 ± 0.32	5.16 ± 0.97	8	0.58 ± 0.07
SR2	0.61 ± 0.25	0.20 ± 0.10	0.35 ± 0.10	0.08 ± 0.02	0.05 ± 0.01	0.03 ± 0.01	0.02 ± 0.00	0.48 ± 0.23	1.83 ± 0.39	2	0.39 ± 0.02
SR3	0.18 ± 0.12	0.11 ± 0.06	0.17 ± 0.07	0.05 ± 0.01	0.00 ± 0.01	0.01 ± 0.01	0.01 ± 0.00	0.00 ± 0.11	0.53 ± 0.24	1	0.23 ± 0.05
SR4	0.63 ± 0.25	0.26 ± 0.14	0.47 ± 0.13	0.13 ± 0.03	0.17 ± 0.12	0.07 ± 0.01	0.09 ± 0.02	0.36 ± 0.21	2.19 ± 0.48	8	0.46 ± 0.05
SR5	0.40 ± 0.19	0.09 ± 0.03	0.22 ± 0.08	0.06 ± 0.02	0.03 ± 0.01	0.03 ± 0.01	0.02 ± 0.00	0.31 ± 0.29	1.17 ± 0.39	0	0.51 ± 0.05
SR6	0.09 ± 0.06	0.03 ± 0.05	0.10 ± 0.04	0.01 ± 0.01	0.07 ± 0.05	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.04	0.32 ± 0.15	2	0.34 ± 0.04
SR7	0.06 ± 0.04	0.04 ± 0.02	0.05 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.05	0.16 ± 0.10	0	0.28 ± 0.08
SR8	0.15 ± 0.08	0.04 ± 0.03	0.08 ± 0.03	0.05 ± 0.01	0.00 ± 0.00	0.02 ± 0.01	0.03 ± 0.01	0.17 ± 0.11	0.55 ± 0.17	0	0.63 ± 0.06
SR9	0.25 ± 0.10	0.49 ± 0.18	0.36 ± 0.10	0.08 ± 0.02	0.02 ± 0.00	0.02 ± 0.01	0.00 ± 0.00	0.22 ± 0.11	1.44 ± 0.28	1	0.24 ± 0.06
SR10	0.08 ± 0.03	0.16 ± 0.07	0.14 ± 0.04	0.02 ± 0.01	0.00 ± 0.00	0.01 ± 0.01	0.00 ± 0.00	0.16 ± 0.19	0.57 ± 0.22	0	0.11 ± 0.03
SR11	0.04 ± 0.02	0.09 ± 0.05	0.05 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.01 ± 0.01	0.00 ± 0.00	0.05 ± 0.04	0.26 ± 0.10	0	0.07 ± 0.02
SR12	0.04 ± 0.02	0.13 ± 0.05	0.09 ± 0.03	0.02 ± 0.00	0.00 ± 0.00	0.02 ± 0.00	0.00 ± 0.00	0.17 ± 0.18	0.46 ± 0.19	1	0.17 ± 0.02
SR13	0.04 ± 0.02	0.05 ± 0.03	0.08 ± 0.02	0.01 ± 0.00	0.02 ± 0.00	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.06	0.20 ± 0.09	0	0.21 ± 0.02
SR14	0.01 ± 0.02	0.05 ± 0.03	0.04 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.11 ± 0.05	1	0.25 ± 0.05

Table 19: Prefit event yields in SR+CR regions for 2017.

	tt̄W	tt̄Z	tt̄H	tt̄V	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	tt̄t̄
CRZ	0.53 ± 0.20	16.57 ± 6.03	0.77 ± 0.22	0.59 ± 0.08	0.08 ± 0.04	3.79 ± 0.81	0.00 ± 0.00	0.79 ± 0.42	23.12 ± 6.24	22	0.32 ± 0.02
CRW	28.09 ± 9.95	7.70 ± 2.83	9.81 ± 2.67	2.04 ± 0.25	3.77 ± 0.57	1.99 ± 0.42	4.98 ± 0.95	15.62 ± 8.52	73.99 ± 13.76	92	1.48 ± 0.07
SR1	1.97 ± 0.72	0.70 ± 0.27	1.35 ± 0.38	0.34 ± 0.05	0.15 ± 0.06	0.17 ± 0.06	0.14 ± 0.03	1.35 ± 1.05	6.17 ± 1.42	10	0.75 ± 0.04
SR2	0.64 ± 0.28	0.08 ± 0.07	0.45 ± 0.14	0.11 ± 0.02	0.01 ± 0.01	0.03 ± 0.01	0.02 ± 0.00	0.48 ± 0.37	1.83 ± 0.52	3	0.39 ± 0.04
SR3	0.13 ± 0.09	0.00 ± 0.02	0.17 ± 0.06	0.03 ± 0.01	0.00 ± 0.00	0.02 ± 0.00	0.01 ± 0.00	0.14 ± 0.16	0.49 ± 0.24	0	0.29 ± 0.05
SR4	0.65 ± 0.25	0.26 ± 0.14	0.45 ± 0.14	0.14 ± 0.02	0.06 ± 0.04	0.05 ± 0.02	0.08 ± 0.02	0.60 ± 0.49	2.28 ± 0.67	2	0.57 ± 0.05
SR5	0.36 ± 0.20	0.10 ± 0.04	0.23 ± 0.08	0.08 ± 0.01	0.01 ± 0.01	0.08 ± 0.02	0.01 ± 0.00	0.08 ± 0.09	0.96 ± 0.27	2	0.58 ± 0.04
SR6	0.14 ± 0.06	0.01 ± 0.04	0.11 ± 0.04	0.03 ± 0.00	0.01 ± 0.00	0.02 ± 0.01	0.01 ± 0.00	0.05 ± 0.06	0.37 ± 0.12	3	0.43 ± 0.03
SR7	0.05 ± 0.03	0.01 ± 0.01	0.04 ± 0.02	0.01 ± 0.00	0.01 ± 0.01	0.01 ± 0.00	0.01 ± 0.00	0.00 ± 0.06	0.14 ± 0.08	0	0.30 ± 0.04
SR8	0.07 ± 0.05	0.04 ± 0.03	0.08 ± 0.04	0.05 ± 0.01	0.02 ± 0.01	0.02 ± 0.01	0.01 ± 0.00	0.28 ± 0.28	0.58 ± 0.30	0	0.76 ± 0.08
SR9	0.32 ± 0.16	0.48 ± 0.18	0.43 ± 0.12	0.09 ± 0.01	0.03 ± 0.01	0.04 ± 0.01	0.00 ± 0.00	0.08 ± 0.07	1.47 ± 0.30	2	0.23 ± 0.01
SR10	0.00 ± 0.03	0.16 ± 0.07	0.15 ± 0.05	0.03 ± 0.01	0.01 ± 0.01	0.01 ± 0.00	0.00 ± 0.00	0.19 ± 0.21	0.55 ± 0.23	3	0.15 ± 0.02
SR11	0.02 ± 0.01	0.06 ± 0.03	0.05 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.02	0.14 ± 0.06	0	0.08 ± 0.02
SR12	0.02 ± 0.01	0.12 ± 0.05	0.08 ± 0.03	0.03 ± 0.00	0.01 ± 0.00	0.01 ± 0.00	0.00 ± 0.00	0.26 ± 0.30	0.53 ± 0.30	1	0.23 ± 0.02
SR13	0.05 ± 0.03	0.07 ± 0.03	0.06 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.06	0.21 ± 0.09	0	0.25 ± 0.02
SR14	0.06 ± 0.03	0.05 ± 0.03	0.04 ± 0.02	0.01 ± 0.00	0.00 ± 0.00	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.04	0.16 ± 0.08	0	0.29 ± 0.03

Table 20: Prefit event yields in SR+CR regions for 2018.

	tt̄W	tt̄Z	tt̄H	tt̄V	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	tt̄t̄
CRZ	0.88 ± 0.33	30.20 ± 11.07	1.34 ± 0.36	0.86 ± 0.12	0.70 ± 0.40	6.36 ± 1.28	0.01 ± 0.00	1.72 ± 0.60	42.07 ± 11.49	51	0.49 ± 0.03
CRW	49.69 ± 18.72	13.94 ± 4.93	16.64 ± 4.19	3.50 ± 0.43	5.88 ± 1.15	5.25 ± 1.10	8.80 ± 1.75	33.41 ± 9.80	137.09 ± 23.04	164	2.24 ± 0.09
SR1	3.27 ± 1.39	1.25 ± 0.47	2.31 ± 0.63	0.58 ± 0.09	0.30 ± 0.09	0.22 ± 0.05	0.24 ± 0.05	1.31 ± 0.62	9.48 ± 1.96	15	1.18 ± 0.07
SR2	1.29 ± 0.53	0.26 ± 0.16	0.71 ± 0.24	0.19 ± 0.04	0.08 ± 0.04	0.10 ± 0.03	0.06 ± 0.01	1.27 ± 0.79	3.95 ± 1.09	4	0.65 ± 0.05
SR3	0.18 ± 0.17	0.04 ± 0.08	0.26 ± 0.09	0.05 ± 0.01	0.01 ± 0.02	0.02 ± 0.00	0.03 ± 0.01	0.05 ± 0.04	0.63 ± 0.31	2	0.39 ± 0.07
SR4	1.51 ± 0.63	0.72 ± 0.32	0.87 ± 0.27	0.32 ± 0.05	0.20 ± 0.06	0.18 ± 0.05	0.15 ± 0.03	0.89 ± 0.42	4.85 ± 1.01	4	0.97 ± 0.05
SR5	0.77 ± 0.32	0.28 ± 0.18	0.46 ± 0.15	0.17 ± 0.03	0.05 ± 0.03	0.15 ± 0.05	0.07 ± 0.01	0.01 ± 0.05	1.97 ± 0.54	3	1.05 ± 0.06
SR6	0.29 ± 0.15	0.09 ± 0.04	0.18 ± 0.07	0.05 ± 0.01	0.04 ± 0.01	0.02 ± 0.00	0.02 ± 0.00	0.36 ± 0.24	1.06 ± 0.33	3	0.71 ± 0.08
SR7	0.10 ± 0.05	0.05 ± 0.03	0.09 ± 0.04	0.02 ± 0.00	0.00 ± 0.00	0.02 ± 0.01	0.00 ± 0.00	0.00 ± 0.04	0.29 ± 0.11	0	0.53 ± 0.07
SR8	0.26 ± 0.16	0.14 ± 0.09	0.22 ± 0.09	0.11 ± 0.02	0.03 ± 0.01	0.07 ± 0.01	0.04 ± 0.01	0.90 ± 0.75	1.76 ± 0.83	5	1.39 ± 0.13
SR9	0.47 ± 0.23	0.84 ± 0.36	0.68 ± 0.19	0.13 ± 0.02	0.09 ± 0.02	0.11 ± 0.09	0.00 ± 0.00	0.56 ± 0.34	2.87 ± 0.71	3	0.44 ± 0.04
SR10	0.15 ± 0.09	0.17 ± 0.08	0.23 ± 0.07	0.05 ± 0.01	0.00 ± 0.01	0.02 ± 0.01	0.00 ± 0.00	0.00 ± 0.04	0.63 ± 0.17	0	0.18 ± 0.04
SR11	0.07 ± 0.05	0.10 ± 0.06	0.09 ± 0.03	0.02 ± 0.00	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.00	0.69 ± 0.75	0.98 ± 0.75	1	0.14 ± 0.02
SR12	0.09 ± 0.06	0.23 ± 0.09	0.22 ± 0.06	0.05 ± 0.01	0.03 ± 0.01	0.03 ± 0.01	0.00 ± 0.00	0.14 ± 0.11	0.79 ± 0.20	0	0.36 ± 0.03
SR13	0.11 ± 0.06	0.16 ± 0.08	0.15 ± 0.05	0.04 ± 0.01	0.00 ± 0.00	0.02 ± 0.01	0.00 ± 0.00	0.00 ± 0.05	0.48 ± 0.16	2	0.42 ± 0.04
SR14	0.04 ± 0.02	0.09 ± 0.06	0.09 ± 0.04	0.02 ± 0.01	0.00 ± 0.00	0.02 ± 0.00	0.00 ± 0.00	0.49 ± 0.27	0.49 ± 0.27	0	0.46 ± 0.06

Table 21: Prefit event yields in SR+CR regions for 2016+2017+2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	1.92±0.70	60.77±21.92	2.80±0.72	1.94±0.22	0.84±0.40	12.94±2.74	0.03±0.00	2.94±1.10	84.18±22.57	104	1.05±0.06
CRW	100.33±36.18	27.95±10.15	34.89±8.58	7.30±0.79	14.06±1.83	9.48±2.08	18.73±2.14	59.08±21.64	271.81±46.38	338	4.94±0.17
SR1	7.11±2.70	2.63±0.96	4.78±1.29	1.20±0.15	0.63±0.14	0.60±0.15	0.49±0.06	3.36±1.41	20.80±3.77	33	2.50±0.12
SR2	2.53±0.98	0.53±0.28	1.51±0.46	0.39±0.06	0.14±0.05	0.17±0.04	0.10±0.01	2.23±1.24	7.60±1.88	9	1.43±0.07
SR3	0.49±0.33	0.16±0.11	0.60±0.22	0.12±0.02	0.02±0.02	0.04±0.01	0.04±0.01	0.19±0.18	1.65±0.61	3	0.92±0.12
SR4	2.78±1.02	1.24±0.53	1.78±0.48	0.60±0.07	0.44±0.14	0.29±0.07	0.32±0.04	1.85±0.87	9.31±1.75	14	2.00±0.09
SR5	1.54±0.64	0.47±0.21	0.92±0.29	0.31±0.04	0.10±0.04	0.26±0.07	0.10±0.01	0.40±0.28	4.10±0.97	5	2.13±0.09
SR6	0.52±0.20	0.14±0.10	0.39±0.14	0.10±0.02	0.12±0.05	0.05±0.01	0.02±0.00	0.41±0.27	1.76±0.48	8	1.49±0.09
SR7	0.22±0.09	0.10±0.04	0.18±0.07	0.04±0.01	0.01±0.01	0.03±0.01	0.01±0.00	0.00±0.08	0.59±0.19	0	1.11±0.10
SR8	0.48±0.24	0.22±0.12	0.38±0.14	0.21±0.03	0.05±0.02	0.11±0.02	0.08±0.01	1.36±0.81	2.89±0.93	5	2.77±0.16
SR9	1.03±0.44	1.80±0.66	1.47±0.39	0.30±0.04	0.15±0.02	0.16±0.09	0.00±0.00	0.86±0.45	5.78±1.16	6	0.90±0.08
SR10	0.23±0.10	0.49±0.17	0.52±0.16	0.11±0.02	0.01±0.01	0.04±0.01	0.00±0.00	0.35±0.24	1.74±0.39	3	0.44±0.06
SR11	0.13±0.07	0.25±0.11	0.19±0.06	0.04±0.01	0.01±0.01	0.02±0.01	0.00±0.00	0.74±0.74	1.38±0.77	1	0.29±0.03
SR12	0.15±0.08	0.48±0.17	0.39±0.10	0.09±0.01	0.04±0.01	0.06±0.02	0.00±0.00	0.57±0.37	1.78±0.46	2	0.75±0.04
SR13	0.20±0.10	0.28±0.11	0.28±0.09	0.07±0.01	0.02±0.01	0.04±0.01	0.00±0.00	0.00±0.09	0.89±0.24	2	0.88±0.05
SR14	0.11±0.05	0.18±0.09	0.17±0.07	0.04±0.01	0.00±0.00	0.03±0.01	0.00±0.00	0.23±0.26	0.76±0.32	1	1.00±0.09

Table 22: Prefit event yields in BDT regions for 2016.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	0.50±0.20	14.01±5.02	0.69±0.15	0.50±0.06	0.06±0.01	2.79±0.59	0.02±0.00	0.42±0.19	18.98±5.15	31	0.25±0.04
SR1	0.24±0.11	0.04±0.03	0.05±0.04	0.00±0.00	0.00±0.15	0.04±0.11	0.20±0.04	0.18±0.12	0.75±0.36	2	0.00±0.00
SR2	0.77±0.41	0.13±0.08	0.21±0.09	0.02±0.01	0.72±0.29	0.23±0.10	0.42±0.08	0.41±0.22	2.92±0.68	4	0.00±0.00
SR3	1.38±0.79	0.38±0.15	0.37±0.14	0.05±0.02	0.30±0.36	0.14±0.09	0.60±0.11	1.53±0.72	4.76±1.48	2	0.01±0.00
SR4	2.23±0.96	0.57±0.26	0.66±0.20	0.07±0.04	0.95±0.45	0.22±0.08	0.69±0.13	1.06±0.42	6.44±1.17	14	0.02±0.02
SR5	2.61±1.19	0.62±0.29	0.84±0.24	0.15±0.03	0.60±0.30	0.27±0.13	0.72±0.14	1.24±0.51	7.05±1.72	9	0.04±0.01
SR6	2.65±1.23	0.92±0.38	1.02±0.24	0.20±0.04	0.27±0.27	0.49±0.14	0.54±0.10	1.33±0.67	7.43±1.72	8	0.05±0.04
SR7	2.87±1.17	1.00±0.48	1.08±0.24	0.18±0.04	0.69±0.17	0.18±0.11	0.47±0.09	1.30±0.56	7.78±1.57	10	0.08±0.04
SR8	3.44±1.42	1.23±0.60	1.53±0.32	0.25±0.05	0.38±0.13	0.30±0.11	0.52±0.10	1.29±0.61	8.96±1.82	12	0.17±0.03
SR9	4.24±1.77	1.63±0.73	1.85±0.41	0.47±0.06	0.65±0.18	0.32±0.13	0.52±0.10	1.76±0.72	11.44±2.37	12	0.32±0.04
SR10	3.40±1.43	1.61±0.77	1.88±0.40	0.48±0.06	0.28±0.16	0.28±0.06	0.33±0.06	1.62±0.77	9.88±2.05	12	0.45±0.04
SR11	2.83±1.21	1.43±0.64	1.72±0.39	0.49±0.05	0.45±0.19	0.21±0.05	0.28±0.05	0.74±0.47	8.14±1.61	14	0.67±0.07
SR12	1.68±0.73	0.98±0.47	1.33±0.31	0.35±0.06	0.17±0.07	0.24±0.06	0.14±0.03	0.99±0.51	5.88±1.18	12	0.86±0.04
SR13	1.03±0.44	0.72±0.31	0.76±0.18	0.25±0.03	0.00±0.02	0.13±0.03	0.08±0.01	0.72±0.38	3.68±0.80	5	0.92±0.04
SR14	0.56±0.33	0.37±0.14	0.44±0.14	0.12±0.02	0.00±0.06	0.07±0.01	0.05±0.01	0.27±0.18	1.89±0.54	4	0.95±0.05
SR15	0.32±0.16	0.15±0.09	0.24±0.07	0.08±0.01	0.01±0.01	0.03±0.01	0.02±0.00	0.10±0.07	0.96±0.27	1	0.94±0.06
SR16	0.12±0.06	0.09±0.05	0.09±0.04	0.03±0.01	0.06±0.06	0.02±0.00	0.01±0.00	0.11±0.09	0.53±0.18	2	0.75±0.07
SR17	0.03±0.03	0.00±0.01	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.05±0.04	0	0.23±0.05

Table 23: Prefit event yields in BDT regions for 2017.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	0.53±0.22	16.57±5.93	0.77±0.18	0.59±0.07	0.08±0.04	3.79±0.86	0.00±0.00	0.79±0.46	23.12±6.17	22	0.32±0.02
SR1	0.28±0.22	0.06±0.04	0.06±0.04	0.01±0.01	0.16±0.12	0.03±0.02	0.17±0.03	0.33±0.39	1.10±0.47	1	0.00±0.00
SR2	0.99±0.48	0.26±0.14	0.21±0.08	0.02±0.01	0.50±0.22	0.13±0.06	0.42±0.08	0.34±0.31	2.86±0.75	7	0.00±0.00
SR3	1.87±1.00	0.53±0.25	0.48±0.15	0.06±0.01	0.33±0.20	0.23±0.18	0.59±0.11	1.63±1.11	5.72±1.81	4	0.01±0.01
SR4	3.13±1.31	0.67±0.28	0.78±0.22	0.11±0.03	0.30±0.34	0.25±0.13	0.65±0.12	2.86±1.58	8.75±2.28	7	0.02±0.01
SR5	3.42±1.49	0.88±0.39	0.92±0.23	0.13±0.02	0.49±0.18	0.33±0.12	0.69±0.13	1.94±1.20	8.80±2.18	9	0.05±0.01
SR6	3.48±1.51	0.94±0.49	1.12±0.28	0.20±0.03	0.67±0.33	0.27±0.09	0.57±0.11	2.31±1.44	9.54±2.45	11	0.07±0.02
SR7	3.34±1.47	1.23±0.57	1.22±0.29	0.21±0.03	0.60±0.14	0.32±0.07	0.52±0.10	1.23±0.86	8.68±1.99	12	0.08±0.02
SR8	4.11±1.76	1.55±0.83	1.70±0.38	0.37±0.05	0.36±0.13	0.32±0.14	0.54±0.10	2.01±1.25	10.96±2.47	13	0.21±0.02
SR9	4.48±1.96	1.89±0.91	2.24±0.52	0.54±0.07	0.54±0.07	0.36±0.11	0.51±0.10	3.20±1.82	13.76±3.12	17	0.35±0.03
SR10	4.15±1.66	1.63±0.82	2.13±0.48	0.49±0.07	0.38±0.20	0.30±0.10	0.37±0.07	1.25±1.43	10.72±2.54	17	0.52±0.10
SR11	3.29±1.44	1.69±0.69	2.06±0.47	0.55±0.07	0.13±0.13	0.19±0.06	0.28±0.05	1.46±1.21	9.64±2.27	13	0.79±0.05
SR12	1.98±0.90	1.17±0.47	1.42±0.33	0.47±0.06	0.19±0.03	0.14±0.10	0.16±0.03	2.38±1.37	7.92±1.87	11	1.03±0.05
SR13	1.04±0.51	0.60±0.34	0.86±0.22	0.27±0.03	0.07±0.03	0.11±0.02	0.07±0.01	0.37±0.42	3.39±0.94	5	1.09±0.05
SR14	0.52±0.26	0.28±0.13	0.44±0.12	0.16±0.02	0.03±0.02	0.10±0.02	0.03±0.00	0.23±0.26	1.78±0.47	6	1.14±0.05
SR15	0.28±0.16	0.11±0.06	0.24±0.08	0.10±0.02	0.03±0.01	0.04±0.01	0.03±0.01	0.14±0.16	0.96±0.29	0	1.12±0.06
SR16	0.13±0.09	0.01±0.02	0.09±0.03	0.04±0.00	0.00±0.00	0.03±0.00	0.01±0.00	0.03±0.06	0.33±0.14	1	0.91±0.06
SR17	0.00±0.00	0.02±0.02	0.01±0.00	0.00±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.25±0.32	0.29±0.32	0	0.26±0.03

Table 24: Prefit event yields in BDT regions for 2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	0.88±0.31	30.20±11.24	1.34±0.35	0.86±0.13	0.70±0.42	6.36±1.55	0.01±0.00	1.72±0.66	42.07±11.64	51	0.49±0.03
SR1	0.50±0.38	0.18±0.10	0.08±0.06	0.01±0.01	0.24±0.20	0.11±0.17	0.29±0.05	0.36±0.36	1.76±0.89	1	0.00±0.00
SR2	1.68±0.87	0.44±0.28	0.38±0.20	0.03±0.02	0.97±0.49	0.50±0.19	0.68±0.13	1.97±1.04	6.65±1.59	8	0.00±0.01
SR3	3.58±1.52	0.91±0.49	0.87±0.31	0.14±0.04	1.01±0.30	0.77±0.24	1.08±0.20	4.74±2.28	13.09±3.14	13	0.02±0.01
SR4	5.17±2.33	1.24±0.71	1.39±0.42	0.21±0.04	0.40±0.49	0.72±0.30	1.17±0.22	3.95±1.41	14.25±3.64	12	0.04±0.01
SR5	6.35±2.41	1.54±1.04	1.69±0.50	0.30±0.06	0.41±0.29	0.62±0.35	1.19±0.22	3.47±1.57	15.56±3.92	18	0.06±0.01
SR6	6.47±2.64	1.85±0.89	1.98±0.54	0.35±0.05	0.02±0.65	0.59±0.21	1.06±0.20	4.12±1.99	16.44±4.36	25	0.10±0.03
SR7	6.14±2.33	2.26±1.26	2.13±0.58	0.42±0.08	0.93±0.20	0.52±0.13	0.87±0.16	4.73±2.16	18.00±3.90	19	0.15±0.03
SR8	7.76±2.87	2.88±1.72	2.86±0.77	0.59±0.07	0.18±0.24	0.59±0.26	0.89±0.17	3.86±1.57	20.31±4.19	21	0.30±0.04
SR9	8.29±3.19	3.30±2.43	3.86±0.98	0.86±0.11	0.95±0.24	0.87±0.27	0.99±0.19	6.14±2.60	25.28±5.48	19	0.46±0.07
SR10	7.39±2.78	2.96±1.84	3.52±0.95	0.90±0.10	0.80±0.34	0.56±0.28	0.70±0.13	5.98±2.66	22.81±4.82	32	0.83±0.04
SR11	5.86±2.36	2.99±1.60	3.50±0.93	0.99±0.12	0.37±0.07	0.52±0.15	0.52±0.10	2.59±1.35	17.34±3.78	35	1.38±0.06
SR12	3.77±1.37	2.03±1.05	2.49±0.68	0.72±0.09	0.34±0.21	0.52±0.15	0.31±0.06	3.04±1.69	13.20±2.82	17	1.65±0.08
SR13	2.14±0.88	1.08±0.68	1.54±0.42	0.48±0.07	0.18±0.04	0.27±0.09	0.13±0.03	1.22±0.51	7.05±1.55	5	2.00±0.09
SR14	0.98±0.45	0.66±0.34	0.90±0.26	0.27±0.04	0.08±0.02	0.13±0.03	0.10±0.02	1.71±1.10	4.84±1.39	6	1.83±0.10
SR15	0.79±0.37	0.42±0.20	0.49±0.14	0.19±0.02	0.17±0.12	0.10±0.02	0.07±0.01	0.19±0.10	2.41±0.60	3	1.78±0.14
SR16	0.18±0.10	0.10±0.09	0.14±0.06	0.08±0.01	0.01±0.00	0.05±0.01	0.01±0.00	0.70±0.47	1.27±0.51	4	1.53±0.11
SR17	0.00±0.02	0.05±0.05	0.02±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.03	0.10±0.08	3	0.52±0.07

Table 25: Prefit event yields in BDT regions for 2016+2017+2018.


	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	1.92± 0.76	60.77±19.35	2.80± 0.61	1.94± 0.25	0.84± 0.35	12.94± 2.43	0.03± 0.00	2.94± 0.97	84.18±19.84	104	1.05± 0.05
SR1	1.02± 0.66	0.28± 0.15	0.19± 0.13	0.02± 0.01	0.40± 0.27	0.18± 0.24	0.66± 0.07	0.86± 0.55	3.61± 1.31	4	0.00± 0.00
SR2	3.45± 1.75	0.83± 0.45	0.80± 0.35	0.07± 0.03	2.19± 0.62	0.85± 0.27	1.52± 0.17	2.71± 1.28	12.43± 2.62	19	0.01± 0.01
SR3	6.83± 3.29	1.82± 0.80	1.72± 0.57	0.25± 0.05	1.65± 0.44	1.14± 0.38	2.26± 0.26	7.90± 2.79	23.58± 5.19	19	0.04± 0.02
SR4	10.53± 4.72	2.47± 1.13	2.83± 0.80	0.39± 0.08	1.66± 0.65	1.18± 0.45	2.51± 0.29	7.86± 2.55	29.44± 6.65	33	0.09± 0.03
SR5	12.38± 5.22	3.04± 1.56	3.45± 0.96	0.58± 0.10	1.49± 0.58	1.23± 0.54	2.59± 0.29	6.64± 2.23	31.41± 7.00	36	0.15± 0.03
SR6	12.60± 5.48	3.71± 1.54	4.12± 1.03	0.74± 0.09	0.97± 0.87	1.35± 0.37	2.17± 0.25	7.76± 2.77	33.41± 7.42	44	0.22± 0.06
SR7	12.35± 5.14	4.50± 2.09	4.43± 1.11	0.81± 0.14	2.21± 0.36	1.02± 0.26	1.86± 0.21	7.27± 2.42	34.45± 6.85	41	0.31± 0.07
SR8	15.32± 6.19	5.66± 2.87	6.10± 1.45	1.21± 0.15	1.61± 0.37	1.21± 0.43	1.95± 0.22	7.16± 2.61	40.22± 8.27	46	0.68± 0.05
SR9	17.01± 7.14	6.82± 3.69	7.95± 1.84	1.87± 0.25	2.14± 0.35	1.56± 0.41	2.02± 0.23	11.11± 4.16	50.48±10.65	48	1.12± 0.08
SR10	14.93± 6.06	6.20± 3.08	7.53± 1.71	1.87± 0.24	1.46± 0.46	1.14± 0.36	1.40± 0.16	8.86± 3.31	43.40± 8.48	61	1.81± 0.07
SR11	11.98± 5.15	6.11± 2.67	7.27± 1.67	2.03± 0.26	0.95± 0.19	0.92± 0.22	1.07± 0.12	4.79± 2.15	35.12± 7.15	62	2.83± 0.14
SR12	7.43± 3.07	4.18± 1.74	5.24± 1.22	1.54± 0.19	0.70± 0.22	0.89± 0.26	0.61± 0.07	6.41± 2.60	27.00± 5.09	40	3.54± 0.12
SR13	4.20± 1.87	2.40± 1.18	3.15± 0.77	1.01± 0.14	0.26± 0.06	0.52± 0.14	0.28± 0.03	2.31± 0.88	14.13± 3.00	15	4.01± 0.14
SR14	2.06± 1.01	1.32± 0.51	1.78± 0.48	0.55± 0.07	0.11± 0.08	0.30± 0.05	0.18± 0.02	2.21± 1.13	8.50± 1.93	16	3.93± 0.15
SR15	1.38± 0.73	0.68± 0.29	0.96± 0.28	0.37± 0.05	0.21± 0.11	0.17± 0.03	0.12± 0.01	0.44± 0.24	4.33± 1.13	4	3.84± 0.20
SR16	0.43± 0.25	0.20± 0.12	0.32± 0.13	0.15± 0.02	0.07± 0.04	0.10± 0.02	0.03± 0.00	0.84± 0.48	2.13± 0.66	7	3.19± 0.17
SR17	0.03± 0.03	0.07± 0.05	0.03± 0.02	0.02± 0.00	0.00± 0.00	0.02± 0.00	0.00± 0.00	0.25± 0.30	0.44± 0.33	3	1.01± 0.10

Table 26: Postfit event yields in SR+CR regions for 2016+2017+2018.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	2.72± 0.56	75.97±10.55	3.00± 0.67	1.98± 0.24	0.85± 0.37	13.37± 2.24	0.03± 0.00	3.24± 1.03	101.17±10.12	104	0.83± 0.49
CRW	142.07±27.77	34.87± 4.83	37.34± 8.06	7.39± 0.83	14.31± 1.79	9.82± 1.74	18.78± 2.53	66.66±19.89	331.24±18.64	338	3.88± 2.28
SR1	10.22± 2.17	3.29± 0.47	5.16± 1.17	1.23± 0.15	0.64± 0.13	0.62± 0.12	0.49± 0.07	3.99± 1.61	25.64± 2.09	33	1.98± 1.18
SR2	3.61± 0.86	0.69± 0.23	1.63± 0.41	0.39± 0.06	0.15± 0.05	0.17± 0.04	0.10± 0.01	2.40± 1.04	9.15± 1.26	9	1.13± 0.65
SR3	0.72± 0.34	0.20± 0.10	0.65± 0.20	0.13± 0.02	0.02± 0.02	0.04± 0.01	0.04± 0.01	0.21± 0.17	2.01± 0.58	3	0.73± 0.42
SR4	4.02± 0.93	1.58± 0.34	1.94± 0.46	0.61± 0.07	0.44± 0.13	0.31± 0.06	0.32± 0.04	2.13± 0.84	11.34± 1.25	14	1.58± 0.90
SR5	2.20± 0.57	0.61± 0.13	1.01± 0.27	0.32± 0.04	0.10± 0.04	0.26± 0.05	0.10± 0.02	0.44± 0.27	5.03± 0.77	5	1.68± 0.95
SR6	0.80± 0.21	0.18± 0.09	0.43± 0.13	0.10± 0.02	0.13± 0.05	0.05± 0.01	0.02± 0.00	0.56± 0.29	2.29± 0.40	8	1.20± 0.67
SR7	0.31± 0.11	0.12± 0.04	0.20± 0.06	0.04± 0.01	0.01± 0.01	0.03± 0.01	0.01± 0.00	0.00± 0.08	0.71± 0.20	0	0.88± 0.48
SR8	0.70± 0.28	0.28± 0.12	0.42± 0.15	0.22± 0.03	0.05± 0.02	0.11± 0.02	0.08± 0.01	1.44± 0.81	3.31± 0.95	5	2.20± 1.27
SR9	1.46± 0.41	2.24± 0.34	1.58± 0.36	0.31± 0.05	0.14± 0.02	0.16± 0.09	0.00± 0.00	0.94± 0.46	6.84± 0.80	6	0.71± 0.39
SR10	0.33± 0.11	0.63± 0.14	0.56± 0.14	0.11± 0.02	0.01± 0.01	0.05± 0.01	0.00± 0.00	0.42± 0.26	2.10± 0.31	3	0.35± 0.22
SR11	0.18± 0.07	0.32± 0.08	0.20± 0.05	0.04± 0.01	0.01± 0.01	0.02± 0.01	0.00± 0.00	0.60± 0.72	1.38± 0.75	1	0.23± 0.14
SR12	0.22± 0.08	0.61± 0.12	0.42± 0.10	0.09± 0.01	0.04± 0.01	0.06± 0.01	0.00± 0.00	0.59± 0.39	2.03± 0.48	2	0.59± 0.34
SR13	0.29± 0.11	0.36± 0.12	0.31± 0.09	0.07± 0.01	0.02± 0.01	0.04± 0.01	0.00± 0.00	0.00± 0.11	1.09± 0.28	2	0.69± 0.39
SR14	0.16± 0.05	0.23± 0.07	0.18± 0.06	0.04± 0.01	0.00± 0.00	0.03± 0.01	0.00± 0.00	0.23± 0.27	0.87± 0.30	1	0.80± 0.45

Table 27: Postfit event yields in BDT regions for 2016+2017+2018.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	2.50± 0.54	77.26±11.71	3.09± 0.64	1.99± 0.21	0.85± 0.35	13.33± 2.56	0.03± 0.00	3.23± 1.18	102.28±11.58	104	1.11± 0.43
SR1	1.25± 0.49	0.31± 0.10	0.19± 0.10	0.02± 0.01	0.32± 0.25	0.17± 0.19	0.66± 0.07	1.04± 0.64	3.95± 0.96	4	0.00± 0.00
SR2	4.35± 1.28	0.97± 0.32	0.83± 0.29	0.08± 0.02	2.20± 0.54	0.79± 0.23	1.53± 0.17	3.46± 1.44	14.19± 1.76	19	0.01± 0.01
SR3	8.53± 2.40	2.17± 0.50	1.75± 0.53	0.23± 0.05	1.46± 0.47	1.12± 0.38	2.26± 0.26	8.01± 2.77	25.53± 3.53	19	0.04± 0.03
SR4	13.52± 3.37	3.04± 0.68	3.00± 0.76	0.38± 0.08	1.57± 0.51	1.19± 0.40	2.52± 0.29	8.74± 3.07	33.96± 4.01	33	0.08± 0.05
SR5	15.80± 3.69	3.73± 0.93	3.68± 0.92	0.57± 0.08	1.61± 0.49	1.27± 0.46	2.59± 0.29	7.41± 2.61	36.67± 3.96	36	0.15± 0.07
SR6	16.34± 3.87	4.50± 0.99	4.41± 1.03	0.75± 0.08	1.26± 0.69	1.40± 0.33	2.18± 0.25	8.99± 3.45	39.81± 4.16	44	0.23± 0.12
SR7	15.96± 3.61	5.53± 1.27	4.75± 1.10	0.82± 0.12	2.25± 0.31	1.03± 0.26	1.86± 0.21	8.12± 3.05	40.32± 3.73	41	0.31± 0.16
SR8	19.81± 4.39	6.88± 1.74	6.61± 1.47	1.21± 0.14	1.65± 0.32	1.21± 0.34	1.95± 0.22	7.97± 2.80	47.28± 4.33	46	0.72± 0.28
SR9	22.22± 4.94	8.57± 2.22	8.63± 1.89	1.88± 0.21	2.15± 0.30	1.53± 0.42	2.02± 0.24	11.50± 3.92	58.51± 5.22	48	1.18± 0.46
SR10	19.68± 4.28	7.79± 1.84	8.31± 1.78	1.91± 0.21	1.39± 0.37	1.18± 0.29	1.40± 0.16	10.48± 3.93	52.15± 4.28	61	1.91± 0.74
SR11	15.96± 3.65	7.82± 1.52	8.03± 1.73	2.06± 0.23	0.98± 0.17	0.97± 0.22	1.07± 0.12	6.10± 2.62	43.00± 3.52	62	2.98± 1.19
SR12	9.66± 2.22	5.30± 1.00	5.81± 1.26	1.56± 0.18	0.81± 0.21	0.89± 0.24	0.61± 0.07	7.46± 3.03	32.10± 3.04	40	3.74± 1.41
SR13	5.57± 1.34	3.01± 0.66	3.50± 0.77	1.04± 0.12	0.27± 0.05	0.51± 0.12	0.28± 0.03	2.52± 0.93	16.71± 1.62	15	4.25± 1.63
SR14	2.74± 0.74	1.67± 0.28	1.97± 0.48	0.56± 0.07	0.11± 0.07	0.31± 0.06	0.18± 0.02	2.61± 1.23	10.14± 1.24	16	4.17± 1.59
SR15	1.81± 0.53	0.84± 0.23	1.06± 0.26	0.38± 0.04	0.20± 0.10	0.17± 0.03	0.12± 0.01	0.45± 0.22	5.03± 0.77	4	4.09± 1.55
SR16	0.55± 0.21	0.26± 0.10	0.36± 0.12	0.15± 0.02	0.07± 0.04	0.10± 0.02	0.03± 0.00	0.97± 0.54	2.49± 0.61	7	3.37± 1.25
SR17	0.05± 0.03	0.09± 0.05	0.04± 0.02	0.02± 0.00	0.00± 0.00	0.02± 0.00	0.00± 0.00	0.34± 0.35	0.57± 0.36	3	1.08± 0.42



(a)

Figure 58: Observed/expected likelihood scans for Run 2 with the BDT analysis

12 Results: interpretations

12.1 Type-II 2HDM

12.1.1 Introduction

Final states with 3 or more top quarks occur in many scenarios of new physics.

In in two Higgs doublet models (2HDM) [23, 24] of type-II, the couplings of the CP-even scalar h become exactly SM-like in the so-called "alignment limit", $\sin(\beta - \alpha) \rightarrow 1$. In such models, the couplings of the heavy scalar and pseudoscalar to the SM vector bosons are suppressed, vanishing as $\cos(\beta - \alpha) \rightarrow 0$. In such a limit, production is predominantly via gluon-fusion and then associated production with either $b\bar{b}$ or $t\bar{t}$. The sensitivity of the direct search for resonant $t\bar{t}$ production is significantly reduced due to interference with SM QCD production [25, 26]. As a result, at low $\tan\beta$, three and four top final states as seen in the diagrams in Figure 59 provide a promising window to probe this scenario [27].




Figure 59: Diagrams for scalar (pseudoscalar) production in association with top quarks. In some scenarios, these heavy bosons will preferentially decay to a pair of top quarks, generating final states with three or four top quarks.

12.1.2 Simulation

We interpret the results of the SM $t\bar{t}\bar{t}$ search as limits on the top-associated production of a scalar or pseudoscalar with subsequent decay to a pair of top quarks.

We generate a one-dimensional grid of points for scalar masses between 350 and 550 GeV in 20 GeV steps for the three processes shown in Figure 59 at LO with MadGraph in the 5-flavor scheme using the NN23LO pdf set. The $t\bar{t}H$ and tWH processes were generated with up to one additional parton, while the tHq process was generated with no additional partons due to the presence of a light flavor quark in the lowest order ME.

For the purpose of interpretation, we use LO cross sections for the production of a heavy scalar or pseudoscalar boson using the 2HDMtII.NLO MadGraph model with the NNPDF30_lo_as_0130 PDF. Processes involving H and A mediators are generated separately and charged higgses (H^\pm) are decoupled by setting their mass to 10TeV. We fix $\sin(\beta - \alpha) = 1$ for the alignment condition, and use $\tan\beta$ as an additional parameter. For consistency with previous results and after verifying consistency with the results of [27], $\tan\beta = 1$ is used for one-dimensional exclusions as a function of mediator mass. To construct a two-dimensional exclusion plane, we calculate cross-sections for the same mediator mass grid with $\tan\beta$ values ranging from 0.5 to 3. Note that with the type-II 2HDM, the branching ratio to up-type quarks (e.g., top quark) is proportional to $\frac{1}{\tan\beta}$.

For the pseudoscalar case, we use the same events as the scalar scenario but with cross sections determined for the case of a CP-odd scalar with the same mass.

988 **12.1.3 Kinematic comparison of scalar and pseudoscalar diagrams**

989 Figure 60 shows analysis-level quantities for $t\bar{t}H$ and $t\bar{t}A$, with the mediator mass set to 450 GeV
990 at $\tan\beta = 1$. The events were simulated with the `2HDMtII_NLO` MG5 model and reconstructed
991 with the 2017 `MINIAODSIM` workflow. In general, acceptance and kinematics are similar be-
992 tween scalar and pseudoscalar processes, especially the final distribution of yields with the
993 BDT-based regions, where a KS statistical test yields a p-value of 0.8. For this reason, scalar
994 samples can be used for both processes as the production cross-section is the primary differ-
995 ence.

DRAFT




Figure 60: Distributions comparing H_T , E_T^{miss} , N_{jets} , $N_{\text{b jets}}$, p_T of the leading jet, and the BDT signal region yields for scalar and pseudoscalar processes. The cross-section has been normalized away, so both shapes and acceptance are relevant here.

996 **12.1.4 Cross-sections vs $\tan\beta$**

997 Cross sections for the scalar and pseudoscalar processes are plotted in Figure 61 and summa-
 998 rized in Table 28, separated by the value of $\tan\beta$.




Figure 61: Cross-sections in units of fb in the plane of $\tan\beta$ vs mediator mass for a heavy scalar boson (top left) and heavy pseudoscalar boson (top right) The bottom plot contains 1-D projections for both mediators for a few values of $\tan\beta$.

Table 28: Cross sections for the case of scalar (H) and pseudoproduction (A), assuming a heavy higgs boson with SM-like top quark couplings. All cross-sections are reported in units of fb, and masses in units of GeV.

$\tan \beta$	mass	σ_{HH}	σ_{tWH}	σ_{tqH}	σ_{tHA}	σ_{tWA}	σ_{tqA}	$\tan \beta$	mass	σ_{HH}	σ_{tWH}	σ_{tqH}	σ_{tHA}	σ_{tWA}	σ_{tqA}
0.5	350	81.4	38.0	77.9	117	36.3	45.7	0.8	350	31.8	14.9	30.4	45.9	14.2	17.8
	370	71.4	34.3	66.7	103	33.0	40.4		370	27.9	13.4	26.1	40.4	12.9	15.8
	390	62.5	31.0	57.5	91.1	29.9	35.7		390	24.4	12.1	22.5	35.6	11.7	13.9
	410	55.2	28.1	49.9	80.1	27.1	31.7		410	21.6	11.0	19.5	31.3	10.6	12.4
	430	48.8	25.4	43.4	70.8	24.7	28.2		430	19.1	9.94	17.0	27.6	9.63	11.0
	450	43.6	23.1	37.9	62.4	22.3	25.1		450	17.0	9.02	14.8	24.4	8.73	9.83
	470	39.1	21.0	33.4	55.4	20.3	22.3		470	15.3	8.22	13.0	21.6	7.94	8.74
	490	35.0	19.1	29.3	49.2	18.5	20.0		490	13.7	7.47	11.4	19.2	7.23	7.82
	510	31.5	17.4	25.7	43.9	16.9	18.0		510	12.3	6.78	10.0	17.2	6.62	7.02
	530	28.3	15.8	22.7	39.1	15.4	16.1		530	11.1	6.17	8.89	15.3	6.03	6.29
	550	25.6	14.4	20.2	35.0	14.1	14.5		550	10.0	5.63	7.90	13.7	5.53	5.65
	570	23.2	13.1	18.0	31.3	12.9	13.0		570	9.07	5.13	7.03	12.2	5.05	5.07
	590	21.0	12.0	16.0	28.1	11.8	11.7		590	8.21	4.70	6.25	11.0	4.61	4.57
	610	19.1	11.0	14.2	25.2	10.8	10.6		610	7.46	4.29	5.56	9.83	4.21	4.13
630	17.3	10.1	12.8	22.6	9.89	9.54	630	6.77	3.95	4.99	8.83	3.86	3.72		
650	15.8	9.24	11.4	20.4	9.09	8.64	650	6.16	3.61	4.46	7.96	3.55	3.37		
1.0	350	18.5	8.66	17.7	29.0	8.98	11.3	1.2	350	11.6	5.40	11.1	20.0	6.20	7.80
	370	17.0	8.19	15.9	25.6	8.18	10.0		370	11.1	5.34	10.4	17.7	5.67	6.92
	390	15.1	7.49	13.9	22.6	7.43	8.85		390	10.0	4.98	9.25	15.7	5.15	6.14
	410	13.4	6.85	12.2	19.9	6.76	7.87		410	9.01	4.60	8.18	13.8	4.68	5.45
	430	11.9	6.24	10.6	17.6	6.13	7.00		430	8.07	4.21	7.18	12.2	4.25	4.86
	450	10.7	5.68	9.32	15.5	5.56	6.26		450	7.25	3.86	6.33	10.8	3.86	4.34
	470	9.61	5.18	8.22	13.8	5.06	5.57		470	6.54	3.53	5.60	9.55	3.51	3.86
	490	8.63	4.70	7.19	12.2	4.61	4.98		490	5.88	3.21	4.92	8.49	3.20	3.46
	510	7.77	4.28	6.35	10.9	4.20	4.47		510	5.31	2.94	4.34	7.58	2.92	3.10
	530	7.00	3.91	5.62	9.74	3.85	4.01		530	4.79	2.67	3.85	6.75	2.66	2.78
	550	6.34	3.56	5.00	8.71	3.51	3.60		550	4.35	2.44	3.43	6.04	2.45	2.50
	570	5.75	3.25	4.44	7.80	3.21	3.23		570	3.94	2.23	3.07	5.41	2.23	2.24
	590	5.20	2.98	3.96	6.99	2.94	2.92		590	3.57	2.05	2.73	4.85	2.05	2.02
	610	4.73	2.72	3.54	6.27	2.69	2.63		610	3.25	1.87	2.43	4.35	1.86	1.83
630	4.29	2.50	3.16	5.63	2.47	2.38	630	2.95	1.72	2.18	3.91	1.71	1.65		
650	3.91	2.28	2.83	5.08	2.25	2.15	650	2.69	1.58	1.95	3.52	1.57	1.49		
1.4	350	6.89	3.23	6.62	14.6	4.52	5.69	1.6	350	5.28	2.48	5.08	11.2	3.47	4.36
	370	7.29	3.52	6.84	13.0	4.15	5.07		370	5.58	2.70	5.24	9.92	3.18	3.89
	390	6.85	3.42	6.32	11.5	3.77	4.49		390	5.24	2.62	4.85	8.77	2.89	3.44
	410	6.27	3.20	5.70	10.1	3.43	4.00		410	4.80	2.45	4.37	7.72	2.63	3.07
	430	5.67	2.97	5.05	8.92	3.12	3.56		430	4.34	2.27	3.88	6.83	2.39	2.73
	450	5.13	2.73	4.48	7.88	2.83	3.17		450	3.93	2.09	3.44	6.03	2.17	2.44
	470	4.66	2.52	3.99	6.99	2.57	2.83		470	3.57	1.93	3.06	5.35	1.97	2.17
	490	4.21	2.30	3.52	6.22	2.34	2.53		490	3.22	1.77	2.71	4.76	1.80	1.94
	510	3.81	2.10	3.12	5.55	2.15	2.28		510	2.91	1.62	2.39	4.25	1.64	1.75
	530	3.44	1.92	2.78	4.95	1.96	2.04		530	2.64	1.48	2.13	3.79	1.50	1.56
	550	3.13	1.76	2.48	4.43	1.79	1.83		550	2.40	1.35	1.90	3.39	1.37	1.41
	570	2.85	1.61	2.21	3.97	1.64	1.65		570	2.18	1.24	1.70	3.04	1.25	1.26
	590	2.58	1.48	1.97	3.55	1.50	1.49		590	1.98	1.14	1.51	2.72	1.15	1.14
	610	2.35	1.36	1.76	3.19	1.37	1.34		610	1.80	1.04	1.35	2.44	1.05	1.03
630	2.14	1.25	1.58	2.86	1.26	1.21	630	1.64	0.956	1.21	2.19	0.964	0.929		
650	1.95	1.15	1.42	2.58	1.15	1.10	650	1.49	0.877	1.09	1.98	0.884	0.840		
1.8	350	3.12	1.47	3.01	8.73	2.72	3.42	2.0	350	1.79	0.845	1.73	7.00	2.19	2.75
	370	3.84	1.86	3.62	7.79	2.51	3.06		370	2.68	1.31	2.53	6.26	2.03	2.47
	390	3.79	1.90	3.51	6.89	2.28	2.72		390	2.80	1.41	2.60	5.55	1.84	2.19
	410	3.55	1.83	3.24	6.07	2.07	2.42		410	2.69	1.39	2.46	4.89	1.68	1.96
	430	3.26	1.72	2.92	5.37	1.88	2.15		430	2.50	1.32	2.25	4.33	1.53	1.74
	450	2.97	1.59	2.61	4.75	1.71	1.92		450	2.31	1.24	2.03	3.83	1.39	1.56
	470	2.72	1.48	2.34	4.21	1.56	1.71		470	2.12	1.16	1.83	3.40	1.26	1.39
	490	2.46	1.35	2.08	3.75	1.42	1.54		490	1.93	1.07	1.64	3.02	1.15	1.24
	510	2.24	1.25	1.84	3.35	1.30	1.38		510	1.76	0.987	1.46	2.70	1.05	1.12
	530	2.03	1.14	1.64	2.98	1.19	1.23		530	1.60	0.904	1.31	2.41	0.963	1.00
	550	1.85	1.05	1.47	2.67	1.08	1.11		550	1.47	0.833	1.17	2.16	0.878	0.900
	570	1.69	0.961	1.31	2.39	0.992	0.998		570	1.34	0.766	1.05	1.93	0.807	0.807
	590	1.53	0.884	1.18	2.14	0.910	0.900		590	1.22	0.706	0.940	1.73	0.736	0.730
	610	1.40	0.811	1.05	1.92	0.830	0.812		610	1.11	0.648	0.841	1.55	0.673	0.658
630	1.27	0.746	0.945	1.73	0.762	0.734	630	1.02	0.597	0.756	1.40	0.618	0.594		
650	1.16	0.687	0.849	1.56	0.696	0.664	650	0.928	0.551	0.681	1.26	0.567	0.538		
2.5	350	0.498	0.240	0.488	4.38	1.40	1.75	3.0	350	0.112	0.056	0.112	2.97	0.978	1.22
	370	1.28	0.633	1.22	3.95	1.30	1.58		370	0.682	0.349	0.668	2.69	0.915	1.11
	390	1.50	0.768	1.41	3.51	1.19	1.41		390	0.899	0.476	0.866	2.40	0.836	0.992
	410	1.52	0.800	1.41	3.10	1.08	1.25		410	0.948	0.514	0.902	2.12	0.764	0.886
	430	1.46	0.784	1.32	2.74	0.985	1.12		430	0.933	0.514	0.868	1.88	0.695	0.791
	450	1.37	0.748	1.22	2.43	0.895	1.00		450	0.889	0.502	0.810	1.66	0.634	0.706
	470	1.27	0.706	1.11	2.16	0.816	0.893		470	0.835	0.477	0.746	1.48	0.578	0.630
	490	1.17	0.657	1.00	1.92	0.743	0.800		490	0.774	0.449	0.678	1.32	0.526	0.565
	510	1.07	0.610	0.898	1.71	0.681	0.718		510	0.714	0.419	0.612	1.18	0.480	0.507
	530	0.982	0.564	0.808	1.53	0.622	0.646		530	0.656	0.389	0.553	1.05	0.440	0.456
	550	0.901	0.520	0.727	1.37	0.568	0.580		550	0.604	0.360	0.499	0.940	0.402	0.409
	570	0.825	0.480	0.654	1.23	0.521	0.521		570	0.554	0.332	0.451	0.842	0.368	0.368
	590	0.753	0.445	0.589	1.10	0.474	0.471		590	0.507	0.308	0.406	0.754	0.337	0.332
	610	0.690	0.409	0.529	0.986	0.435	0.425		610	0.466	0.284	0.365	0.677	0.307	0.300
630	0.631	0.377	0.476	0.887	0.398	0.384	630	0.426	0.262	0.329	0.609	0.281	0.272		
650	0.578	0.347	0.429	0.800	0.365	0.348	650	0.391	0.243	0.298	0.549	0.258	0.245		

999 **12.1.5 Exclusions**

1000 With the Run2 BDT analysis and the previous model assumptions, heavy scalar (pseudoscalar)
 1001 bosons are excluded as shown in Figure 62, as a function of mediator mass for $\tan\beta = 1$, and
 1002 in Figure 63, as a function of mediator mass and $\tan\beta$.




Figure 62: Observed and expected limits on heavy scalar (left) and pseudoscalar (right) processes as a function of the (pseudo)scalar mass using FullSim samples. The parameter $\tan\beta$ is assumed to be 1 here.




Figure 63: Two-dimensional observed and expected exclusions of heavy scalar only (top left), pseudoscalar only (top right), and both scalar+pseudoscalar (both) as a function of the mediator mass and $\tan\beta$. In each case, other 2HDM particles (except SM higgs) are decoupled.

12.2 Top Yukawa coupling

12.2.1 Introduction

In the SM there are contributions to $pp \rightarrow t\bar{t}\bar{t}$ from diagrams with virtual Higgs bosons, see for example Figure 64. The amplitude corresponding to these diagrams is proportional to the square of the top Yukawa coupling.




Figure 64: One of the Feynman diagrams for $t\bar{t}\bar{t}$ including a virtual Higgs.

Using the notation of Reference [28] the $t\bar{t}\bar{t}$ cross-section can be written as

$$\sigma(t\bar{t}\bar{t}) = \sigma^{SM}(t\bar{t}\bar{t})_{g+Z/\gamma} + k_t^4 \sigma^{SM}(t\bar{t}\bar{t})_H + k_t^2 \sigma_{\text{int}}^{SM} \quad (6)$$

where $k_t \equiv y_t/y_t^{SM}$, y_t is the top Yukawa coupling, and y_t^{SM} is its value in the SM. In equation 6 the first term on the right hand side corresponds to the SM contribution to the cross section from diagrams with virtual gluons or Z/γ , the second term is the contribution from diagrams with virtual H bosons, and the third term is the interference between the two. Therefore, given a theoretical calculation and a measurement of $\sigma(t\bar{t}\bar{t})$, one can put constraints on $|y_t/y_t^{SM}|$.

The authors of Reference [28] have calculated the cross-section terms at LO, and have provided us privately with the uncertainties under variations of the factorization and renormalizations scales. These are given in Table 29.

	$[\mu/2, \mu, 2\mu]$
$\sigma^{SM}(t\bar{t}\bar{t})_{g+Z/\gamma}$	[14.104, 9.997, 6.378] fb
$\sigma^{SM}(t\bar{t}\bar{t})_H$	[1.625, 1.167, 0.7655] fb
σ_{int}^{SM}	[-2.152, -1.547, -0.999] fb

Table 29: LO calculation of the terms in equation 6 from Reference [28]. The scale variations are private communications from the authors.

We have investigated the possibility of a full NLO treatment of the interpretation, and discussed with the authors of the NLO calculation [6], but Madgraph could not provide what we need yet. So, we have decided to continue using the LO calculation for this interpretation. The authors of Reference [28] have also argued that it is appropriate to apply the overall NLO/LO k-factor of 1.27 calculated at 14 TeV for the **total** $\sigma(t\bar{t}\bar{t})$ cross-section [29] to the individual components in equation 6. This would then result in an NLO cross-section of $12.2_{-4.4}^{+5.0}$ fb. This is

1023 in agreement with the NLO calculation of $11.97_{-2.51}^{+2.15}$ fb [6].¹ The authors of Reference [28] then
 1024 go on to extract a limit on $|y_t/y_t^{SM}|$ based on their calculation and the then-available experi-
 1025 mental limit on $\sigma(t\bar{t}\bar{t})$. Following this procedure in the 2016 result, in Fig. 65 (left) we show
 1026 the measurement of the cross section and its upper limit, as well as its SM prediction as a
 1027 function of the absolute value of the ratio of the top quark Yukawa to its SM value ($|y_t/y_t^{SM}|$).
 1028 The central (upper,lower) value of the theoretical cross section band resulted in a 95% CL limit
 1029 $|y_t/y_t^{SM}| < 2.27$ (2.03,2.56) (in the 2016 result).

1030 Noting that $t\bar{t}H$ is a non-negligible background, showing a flat line for the observed cross sec-
 1031 tion of $t\bar{t}\bar{t}$ (and observed upper limit) as a function of κ_t is not completely correct, as it neglects
 1032 the κ_t^2 scaling of $t\bar{t}H$, since $\sigma(t\bar{t}H) \propto (y_t/y_t^{SM})^2/(\Gamma_H/\Gamma_H^{SM})$, where Γ_H is the total width of the
 1033 Higgs. The limit plot therefore is also shown, on the right, for the case of the $t\bar{t}H$ background is
 1034 scaled by $(y_t/y_t^{SM})^2$ for each value of $|y_t/y_t^{SM}|$. As the $t\bar{t}H$ background becomes larger, fewer
 1035 $t\bar{t}\bar{t}$ events are allowed by the fit, and the observed $t\bar{t}\bar{t}$ upper limit and the measured $t\bar{t}\bar{t}$ cross
 1036 sections becomes smaller. With this prescription, the central (upper,lower) value of the theoret-
 1037 ical cross section band results in a 95% CL limit $|y_t/y_t^{SM}| < 2.10_{-0.27}^{+0.22}$ (in the 2016 result).

1038 For the full Run2 BDT analysis, this procedure resulted in a 95% CL limit of the central, upper,
 1039 and lower values of the theoretical cross section provide respective 95% CL limits for $|y_t/y_t^{SM}|$
 1040 < 1.7 , < 1.4 , and < 2.0 , as shown in Figure 66.




Figure 65: The expected $t\bar{t}\bar{t}$ cross-section, $\sigma(t\bar{t}\bar{t})$, as a function of $|y_T/y_T^{SM}|$, the absolute value of the ratio of the top quark Yukawa coupling and its SM value (diagonal band), compared with the measured $\sigma(t\bar{t}\bar{t})$ (horizontal band), and its 95% CL upper limit (horizontal line). The right plot includes the scaling of the $t\bar{t}H$ background as a function of y_T as discussed in the text.

1041 12.2.2 Alternative statistical treatment

1042 Note that this section is from 2016, but is left here as a reference

1043 An alternative analysis, still based on the LO cross-section (and its uncertainty) from Refer-
 1044 ence [28], scaled up by the k-factor of 1.27, consists of interpreting the experimental likelihood
 1045 as the posterior pdf for the cross-section.² We then construct (correlated) pdfs for the three

¹Note that the uncertainties in the full NLO calculation are smaller, but this is to be expected.

²This would correspond to using a Bayesian method with a flat prior in the cross-section. It is not entirely kosher




Figure 66: The expected $\bar{t}t\bar{t}$ cross-section, $\sigma(\bar{t}t\bar{t})$, as a function of $|y_T/y_T^{SM}|$, the absolute value of the ratio of the top quark Yukawa coupling and its SM value (diagonal band), compared with the measured $\sigma(\bar{t}t\bar{t})$ (horizontal band), and its 95% CL upper limit (horizontal line)

1046 terms on the right side of equation 6, and extract a pdf for $|y_T/y_T^{SM}|$ by propagating uncer-
 1047 tainties using equation 6 and a toy MC method. We do this in two ways, by either taking the
 1048 correlated pdfs as bifurcated gaussians (truncated to not allow a change of sign), or as “flat”
 1049 within the limits of Table 29. The 95% CL limit would then be $|y_T/y_T^{SM}| < 2.45$ (bifurcated gaus-
 1050 sians pdfs) or $|y_T/y_T^{SM}| < 2.25$ (flat pdfs). These results are to be compared with the “constant
 1051 ttH” interpretation of Figure 65 (left), which resulted in a 95% CL limits of $|y_T/y_T^{SM}| < 2.27$
 1052 (2.03,2.56).

1053 We prefer to present the results according to Figure 65. This is in the same spirit as what is
 1054 customary in the SUSY group, where the experimental limits are clearly separated from the
 1055 theoretical uncertainties in the interpretation (but maybe the TOP group does it differently).
 1056 In addition, we are reluctant to give too much weight to an error analysis that includes the
 1057 systematic uncertainty from Reference [28], which is somewhat of a hack (a LO calculation with
 1058 the pieces of the LO calculation scaled by the same k-factor as obtained by another group for
 1059 the full calculation at a slightly different energy...). Incidentally, the limit on $|y_T/y_T^{SM}|$ quoted in
 1060 Reference [28] ignores the uncertainty on the theoretical calculation. In any case, we are open
 1061 to suggestions.

1062 12.2.3 Kinematic dependence on top yukawa coupling

1063 Since MC is taken with a nominal top yukawa coupling and only the cross-section is scaled in
 1064 the scan above, it is worth verifying that there is no significant kinematic dependence on the
 1065 top yukawa coupling value. This is checked with LO MG5 (MG5_aMC.v2.6.3.2) with default
 1066 parameters, including a dynamical scale choice) using the following proc card.

```
1067 set default_unset_couplings 99
1068 set group_subprocesses Auto
1069 set ignore_six_quark_processes False
```

since the likelihood has been profiled with respect to the nuisances, while Bayesian approaches usually require marginalization.

```

1070 set loop_optimized_output True
1071 set loop_color_flows False
1072 set gauge unitary
1073 set complex_mass_scheme False
1074 set max_npoint_for_channel 0
1075 set nb_core 4
1076
1077
1078 import model sm
1079 define p = g u c d s u~ c~ d~ s~
1080 define p = p b b~
1081 generate p p > t t~ t t~ QED=99
1082
1083
1084 output ftlo_ytscan
1085 launch
1086
1087 # >>> import numpy as np
1088 # >>> x = np.arange(0.4,2.2,0.1)
1089 # >>> print ", ".join(map(lambda y: str(round(y,1)),x))
1090 # paste output in scan brackets below
1091
1092 set param_card yukawa 6 scan:[69.2,...,363.3]

```

1093 The LHE for $|y_T/y_T^{SM}|$ between 0.4 and 2.2 in steps of 0.1 was carried through the 2016 MINIAOD-
 1094 SIM workflow with a slightly modified Pythia fragment which imposes a dilepton (or more)
 1095 filter for better statistical uncertainties. The relevant part of the fragment is

```

1096 'JetMatching:setMad = off',
1097 'JetMatching:scheme = 1',
1098 'JetMatching:merge = on',
1099 'JetMatching:jetAlgorithm = 2',
1100 'JetMatching:etaJetMax = 5.',
1101 'JetMatching:coneRadius = 1.',
1102 'JetMatching:slowJetPower = 1',
1103 'JetMatching:qCut = 59.',
1104 'JetMatching:nQmatch = 5', #4 corresponds to 4-flavour scheme (no matching of b-
1105 'JetMatching:nJetMax = 0', #number of partons in born matrix element for highest
1106 'JetMatching:doShowerKt = off', #off for MLM matching, turn on for shower-kT mat
1107 '6:m0 = 172.5',
1108 '24:mMin = 0.1',
1109 '23:mMin = 0.1',
1110 'ResonanceDecayFilter:filter = on',
1111 'ResonanceDecayFilter:exclusive = off', #off: require at least the specified num
1112 'ResonanceDecayFilter:eMuAsEquivalent = off', #on: treat electrons and muons as
1113 'ResonanceDecayFilter:eMuTauAsEquivalent = on', #on: treat electrons, muons , an
1114 'ResonanceDecayFilter:allNuAsEquivalent = on', #on: treat all three neutrino fla
1115 'ResonanceDecayFilter:daughters = 11,11',
1116 'Check:abortIfVeto = on',

```

1117 To validate the MadGraph setup, cross-sections for points along the scan are compared with
 1118 Equation 6 (using values from Table 29) and they are found to agree within 2%, as shown in
 1119 Figure 67




Figure 67: Calculated cross-section from Equation 6 and from MG5 for varying $|y_T/y_T^{SM}|$ values

1120 Finally, plots of various kinematic quantities are shown in Figure 68 for the analysis baseline
 1121 selection, also allowing opposite-sign events in addition to same-sign events for augmented
 1122 statistics. There is no visible significant difference in the kinematic plots for different values of
 1123 $|y_T/y_T^{SM}|$.




Figure 68: Kinematic quantities (left to right, top to bottom: H_T , N_{jets} , N_{btags} , MET , $p_T(\text{lep1})$, $p_T(\text{lep2})$) for the baseline selection with same-sign and opposite-sign dileptons for $|y_T/y_T^{SM}|$ of 0.5, 1.0, 1.5, 2.0. All histograms are normalized to the same cross-section.

1124 12.3 Off-shell mediators decaying to top quark pairs

1125 12.3.1 Introduction

The production of $t\bar{t}\bar{t}$ may also be influenced by a neutral scalar mediator (ϕ) or neutral vector mediator (Z') which couple to top quarks and have masses less than twice the mass of the top quark, distinguishing them from similar processes within the 2HDM framework, for example. The off-shell contributions to the SM $t\bar{t}\bar{t}$ production can be large, as shown in Ref. [30]. For a large range of masses, the authors have shown that kinematics are identical when considering these additional processes, so that the total $t\bar{t}\bar{t}$ cross-section is subject to a simple rescaling. We consider coupling terms in the lagrangian of the form

$$\mathcal{L}_{Z'} = -g_{tZ'}\bar{t}_R Z' t_R \quad \mathcal{L}_\phi = -g_{t\phi}\bar{t}_L \phi t_R$$

1126 and calculate leading order cross-sections for the process

1127 `generate p p > t~ t t~ t QED=2`

1128 with MadGraph UFO models provided by the authors of Ref. [30], which have been copied to
 1129 <https://github.com/aminnj/FTInterpretations/tree/master/models>. We also
 1130 considered single top processes, similar to those for the 2HDM exclusions, but found their
 1131 cross-section to be small compared to $t\bar{t}$ -associated production, so they are not included.

1132 Due to the approximate independence of kinematics on the coupling strength and mediator
 1133 mass, we are able to use the upper limit result from the nominal analysis to place constraints
 1134 on couplings $g_{tZ'}$ and $g_{t\phi}$ as a function of masses $m_{Z'}$ and m_ϕ , respectively, without the use
 1135 of dedicated signal samples. The nominal analysis uses a NLO $t\bar{t}\bar{t}$ sample, so to justify this
 1136 procedure, we first show that the nominal NLO $t\bar{t}\bar{t}$ sample indeed has good shape agreement
 1137 with a LO $t\bar{t}\bar{t}$ generated with these models (setting coupling strengths to 0) in Figure 69. Next,
 1138 we show various couplings and mass points near the exclusion boundary compared to the LO
 1139 SM $t\bar{t}\bar{t}$ sample in Figure 70 for the Z' mediator, and Figure 71 for the ϕ mediator. Based on
 1140 the level of agreement for both mediator types, we include a 10% additional normalization
 1141 uncertainty on the SM $t\bar{t}\bar{t}$ signal to conservatively cover minor acceptance differences.




Figure 69: Distributions comparing H_T , E_T^{miss} , N_{jets} , $N_{b jets}$, and the BDT signal region yields for the nominal NLO and LO $t\bar{t}$ samples. Only shapes are relevant here.




Figure 70: Distributions comparing H_T , E_T^{miss} , N_{jets} , $N_{b jets}$, and the BDT signal region yields for the LO $t\bar{t}$ samples and various vector mediator mass points. Both shape and acceptance are relevant here.




Figure 71: Distributions comparing H_T , E_T^{miss} , N_{jets} , $N_{\text{b jets}}$, and the BDT signal region yields for the LO $tt\bar{t}\bar{t}$ samples and various scalar mediator mass points. Both shape and acceptance are relevant here.

1142 The BDT observed upper limit on $t\bar{t}$ production, including the extra 10% uncertainty previ-
 1143 ously motivated, is 23 fb^{-1} . Taking the ratio with the SM cross-section gives approximately 1.9.
 1144 This is represented as the horizontal dashed line in Figure 72, which includes curves for cross-
 1145 sections (normalized to SM) for various mediator masses as a function of coupling strengths.
 1146 Intersections between the cross-section curves and the horizontal dashed line are marked with
 1147 vertical dashed lines. To obtain smoother values, a quadratic interpolation between generated
 1148 points is used. These intersections form the exclusion boundary as shown in Figure 73.




Figure 72: Cross-section (normalized to SM) as a function of coupling values with varying mediator masses for vector (left) and scalar (right) mediators. The observed analysis upper limit is shown as a horizontal dashed line.




Figure 73: Observed 95% CL exclusions in the plane of coupling-mass for off-shell vector and scalar mediators.

12.4 Oblique Higgs parameter

12.4.1 Cross-section calculation

In a universal effective field theory framework, the Higgs oblique parameter \hat{H} , defined as the Wilson coefficient of the dimension-6 operator modifying the Higgs boson propagator, can result in deviations of the SM $t\bar{t}t\bar{t}$ cross-section, as shown in Ref. [31]. These (off-shell) deviations can be constrained to a level which is competitive with constraints from on-shell processes.

The two main characteristic effects of this oblique parameter are an additional contact term in the SM Higgs boson propagator

$$P_h(p^2) \approx \frac{i}{p^2 - m_h^2} - \frac{i\hat{H}}{m_h^2},$$

and a rescaling of the fermionic higgs couplings

$$\kappa_f = 1 - \hat{H}.$$

Using the latest combined fits of ATLAS for the (on-shell) fermionic couplings, with 80 fb^{-1} of 13TeV data, the authors of Ref. [31] find a constraint on the oblique parameter of $\hat{H} < 0.16$ at 95% CL.

The authors also calculate that the cross-section of (off-shell) $t\bar{t}t\bar{t}$ is subject to a fractional modification (with respect to the SM cross-section) at 14 TeV, given by,

$$\frac{\sigma_{\hat{H}+\text{SM}}}{\sigma_{\text{SM}}} = 1 + 0.03 \left(\frac{\hat{H}}{0.04} \right) + 0.15 \left(\frac{\hat{H}}{0.04} \right)^2.$$

For an oblique parameter value of 0.1, the formula predicts a doubling of the SM cross-section of $t\bar{t}t\bar{t}$ with MG 2.6.1 and 2.6.5

The SM model within MadGraph was modified to take into account the **extra term in the propagator**, as well as the **rescaling of the top-yukawa coupling**, by changing only the numerator of the scalar propagator affecting the Higgs boson:

$$\frac{i}{p^2 - m^2 + im\Gamma} \rightarrow \frac{(1 - \hat{H})^2 \left(1 - \frac{\hat{H}}{m^2} (p^2 - m^2) \right)}{p^2 - m^2 + im\Gamma}.$$

Note that the complex i in the numerator of the default propagator is removed due to an internal inconsistency found in the latest versions of MadGraph, as verified by the authors. Explicitly, we took the SM model UFO file and modified the numerator of S in file `propagators.py` to be

```
"(1-hhat) * (1-hhat) * (1 - (hhat / (Mass(id) * Mass(id)))" +
"* (P('mu', id) * P('mu', id) - Mass(id) * Mass(id))"
```

instead of "i", and added the line

```
propagator = Prop.S,
```

1168 to the SM Higgs particle definition in file `particles.py`. Finally, for convenience of scanning
 1169 the parameter \hat{H} , we added

```
1170 hhat = Parameter(
1171     name = 'hhat',
1172     nature = 'external',
1173     type = 'real',
1174     value = 0.,
1175     texname = '\\text{hhat}',
1176     lhablock = 'PROP',
1177     lhacode = [ 1 ],
1178 )
```

1179 to file `parameters.py`. The final model can be found at [https://github.com/aminnj/](https://github.com/aminnj/FTInterpretations/tree/master/models/Oblique_UFO)
 1180 `FTInterpretations/tree/master/models/Oblique_UFO`.

1181 12.4.2 Private generation details

1182 We privately generated five different values of \hat{H} with the above modifications to the SM model
 1183 in MadGraph 2.6.5 with the `nn231o1` PDF and default (dynamic) scale choices at leading order.
 1184 Each parameter point consists of three datasets of 50k events each with LHE→MINIAODSIM
 1185 configurations matching the `RunIISummer16MiniAODv3`, `RunIIFall17MiniAODv2`, and
 1186 `RunIIAutumn18MiniAOD` campaigns. To increase the baseline selection efficiency, we used
 1187 the dilepton Pythia filter described in Section 12.2, which requires events to have at least 2 gener-
 1188 ator leptons (of any flavor) with a filter efficiency of 0.3944. The cross-sections (times filter
 efficiencies) for these parameter points are tabulated below.

\hat{H}	$\sigma \times \epsilon_{\text{filter}}$ (fb)
0.0	4.721
0.04	5.411
0.08	6.752
0.12	8.456
0.16	10.235

1189

1190 12.4.3 Comparison of kinematics/acceptance

1191 Figure 74 shows analysis-level quantities for five different values of \hat{H} , one of which is analo-
 1192 gous to SM $t\bar{t}t\bar{t}$ ($\hat{H} = 0$). In the last BDT bin, the acceptance for the two values of non-zero \hat{H} are
 1193 within approximately 10% of the SM. In general, acceptance for those non-zero points is 10-20%
 1194 higher than the SM in the higher BDT bins, with visible trends in jet/lepton momentum, and
 1195 H_T . The nominal analysis upper limit cannot be directly used to place a constraint on \hat{H} as ac-
 1196 ceptance increases for higher values of \hat{H} . However, the next section outlines the method of ex-
 1197 clusion while directly using the nominal analysis result (assuming SM acceptance/kinematics).
 1198 The subsequent section then takes into account acceptance/kinematic differences by showing
 1199 the result with dedicated samples for each \hat{H} point.




Figure 74: Distributions comparing H_T , E_T^{miss} , N_{jets} , N_{bjets} , p_T of the leading lepton, p_T of the subleading lepton, p_T of the leading jet, and the BDT signal region yields for three values of the oblique parameter \hat{H} . Both shapes and acceptance are relevant here, as cross-section has been normalized to 12 fb^{-1} .

12.4.4 Exclusion, assuming SM kinematics

Through correspondence with the authors, we recalculated the 14TeV formula and find cross-sections to match closely with the predicted formula above, for values $\hat{H} < 0.04$, as the reference expands only to second order due to their tighter region of interest (HL-LHC projections). Figure 75 shows a recalculation of the 14TeV formula instead at 13TeV, as well as a cubic fit to the values in order to extract an upper limit on the oblique parameter using the nominal analysis BDT upper limit. We find $\hat{H} < 0.13$ at 95% CL.

As is the case for the top Yukawa interpretation from Section 12.2, the $t\bar{t}H$ background is also subject to modifications due to non-zero \hat{H} . The $t\bar{t}H$ background is affected by a cross-section scaling due to the reduction of the top yukawa constant ($y_t \rightarrow y_t - \hat{H}$). Thus, we can utilize the observed upper limits calculated for the $y_t < 1$ section of the curve in Figure 66. An increase in \hat{H} corresponds to slight increase of the observed upper limit, which translates into a weakening of the \hat{H} constraint by 0.005 (4% relative).




Figure 75: Cross-section (normalized to SM) as a function of oblique parameter \hat{H} . The red curve represents equation 5.4 from Ref. [31] at 14TeV. The blue and green curves are private calculations from MadGraph at 14TeV and 13TeV, respectively. The solid black curve is a cubic fit to the 13TeV private calculation, which is used to find the intersection with the (nearly) horizontal black dotted line (nominal analysis upper limit, corrected to account for deviations from $t\bar{t}H$, as described in the text). For the sake of comparison, the (completely) horizontal gray dotted line does not take this effect into account. The intersection of the black dotted lines provides the upper limit on \hat{H} .

12.4.5 Exclusion using dedicated samples

Repeating the previous subsection with privately-generated dedicated samples yields Figure 76 and excludes $\hat{H} > 0.12$. Explicitly, we substitute the privately-generated \hat{H} -modified $t\bar{t}t\bar{t}$ sam-

1216 ple for the $t\bar{t}\bar{t}$ signal, scale down the $t\bar{t}H$ background normalization by $y_t^2 = (1 - \hat{H})^2$ (≈ 0.7 at
 1217 $\hat{H} = 0.16$), and run the nominal limit-setting procedure.

1218 Note that the upper limit becomes more stringent for higher values of \hat{H} , consistent with the
 1219 increase in acceptance shown in Figure 74.

1220 We have produced official gridpacks and asked the TOP MC contacts to generate official CMS
 1221 samples using the configurations for 2016, 2017 and 2018 respectively. These samples are ex-
 1222 pected to be identical to the ones we generated privately, since the same process will be used.




Figure 76: See caption for Figure 75. An additional red curve was added connecting the five upper limits calculated with dedicated samples. The intersection of a linear interpolation between these points (in red) with the black curve defines the 95% CL upper limit on \hat{H} .

12.5 Dark matter

12.5.1 Introduction

Upper limits on additional $t\bar{t}\bar{\chi}$ production can be translated into exclusions/upper limits on simplified dark matter models which have a scalar or pseudoscalar mediator decaying into a pair of dark matter or standard model particles. The production of the mediator (decaying into invisible dark matter) in association with a pair of top quarks was performed by CMS with the 2016 dataset in Ref. [32]. Reference [33] found that production of the mediator in association with a single top quark contributes meaningfully to the total cross-section. Thus, a search that also included these single top associated production modes was carried out with the same dataset in Ref. [34], which we aim to complement in this analysis with an orthogonal final state. A brief overview is included below, and further theoretical details and motivation can be found in the Analysis Note for EXO-18-010 ([35]).

In the framework of a simplified dark matter model, where the scalar (ϕ) or pseudoscalar (a) mediator couples dark matter and SM particles, the relevant lagrangian terms are of the form

$$\mathcal{L}_\phi = g_\chi \phi \bar{\chi} \chi + \frac{g_q \phi}{\sqrt{2}} \sum_f y_f \bar{f} f \quad \mathcal{L}_a = i g_\chi a \bar{\chi} \gamma^5 \chi + \frac{i g_q a}{\sqrt{2}} \sum_f y_f \bar{f} \gamma^5 f$$

where y_f are the fermionic yukawa couplings. The coupling constants g_χ and g_q give the relative strengths of the mediator coupling to dark matter and SM particles, and are used interchangeably with g_{DM} and g_{SM} , respectively. The model has four free parameters (g_χ , g_q , m_χ , and m_a) which is reduced to two with the assumption of $g_\chi = g_q = 1$.

The relevant production diagrams are shown in Figure 77. When the mediator mass is above $2m_{\text{top}} \approx 350\text{GeV}$, on-shell decay to $t\bar{t}$ becomes kinematically accessible, resulting in 3 or 4 top quark final states, so we instead consider a version of the diagrams with a decay of the mediator into $t\bar{t}$ rather than a pair of (invisible) dark matter particles. Consequently, the production diagrams and kinematics are identical to that of the 2HDM interpretation which will allow us to use existing simulation samples for this interpretation, and we only need to calculate branching ratios into $t\bar{t}$.

In this way, our final state is complementary to the $t\bar{t}$ +MET final state used in Ref. [34] when the DM mass becomes large.

Using MadGraph 2.6.5, and the models from the official gridpacks used for the EXO-18-010 result we first verified consistency with the corresponding AN cross-sections before modifying the $\phi/a \rightarrow \chi\bar{\chi}$ decay to be $\phi/a \rightarrow t\bar{t}$. We use MadGraph to calculate cross-section (times branching ratio) for the $t\bar{t}$ associated process and two single-top processes for mediator masses matching the available 2HDM simulation samples at varying dark-matter masses. We use the NNPDF30_lo_as_0130 PDF.

While we use MadGraph-calculated values in the end to account for all effects, we independently verified that relative branching ratios for scalar and pseudoscalar, respectively, are given by

$$\Gamma_{\text{ratio}} \equiv \frac{\Gamma(\phi \rightarrow t\bar{t})}{\Gamma(\phi \rightarrow \chi\bar{\chi})} = \frac{3g_{\text{SM}}^2 y_t^2}{2g_{\text{DM}}^2} \left(\frac{M^2 - 4m_t^2}{M^2 - 4m_\chi^2} \right)^{3/2} \quad (7)$$

$$\Gamma_{\text{ratio}} \equiv \frac{\Gamma(\phi \rightarrow t\bar{t})}{\Gamma(\phi \rightarrow \chi\bar{\chi})} = \frac{3g_{\text{SM}}^2 y_t^2}{2g_{\text{DM}}^2} \left(\frac{M^2 - 4m_t^2}{M^2 - 4m_\chi^2} \right)^{1/2} \quad (8)$$




Figure 77: Diagrams for scalar (pseudoscalar) mediator production in association with a $t\bar{t}$ pair (top left), associated t-channel single top (top right), associated tW (bottom row). The mediator decays into a pair of invisible particles. As the s-channel single top production cross-section is relatively negligible (by at least an order of magnitude), it is not included in this result.

1254 where y_t is the top-quark yukawa coupling, M is the mediator mass, and ϕ represents both
 1255 the scalar and pseudoscalar. Note that the only difference is in the exponent of the mass term.
 1256 In practice, these formulae would be used after calculating the production cross-section with
 1257 $\text{BR}(\phi \rightarrow t\bar{t})=1$ with MadGraph for each mediator mass value (e.g., $M = 450\text{GeV}$ with $m_\chi >$
 1258 450GeV so that the DM decay is kinematically suppressed.) Then that production cross-section
 1259 is multiplied by $\text{BR}(t\bar{t}) = \frac{\Gamma(t\bar{t})}{\Gamma(t\bar{t}) + \Gamma(\text{DM})} = \left(1 - \Gamma_{\text{ratio}}^{-1}\right)^{-1}$ in order to get the $\sigma \cdot \text{BR}$ at an arbitrary
 1260 m_χ . The formulae assume that there are only two accessible decay modes of the mediator.

1261 **12.5.2 Exclusions**

1262 As noted previously, the processes are identical to those of the 2HDM interpretation with a
1263 different labeling of the mediator, so we can calculate a single cross-section upper limit for a
1264 given mediator mass using pre-existing samples. The single value is used to exclude along the
1265 vertical axis (m_{DM}) since kinematics remain identical and only the branching ratio to $t\bar{t}$ changes.
1266 This can be seen in Figure 78, which overlays exclusion contours on cross-sections calculated
1267 at discrete points. The diagonal line corresponds to the kinematic boundary between on- and
1268 off-shell mediator decays into dark matter, $m_{\text{mediator}} = 2m_{\text{DM}}$, above which $t\bar{t}$ is the dominant
1269 decay. For couplings set to unity, the region above the diagonal is excluded for both mediator
1270 types between approximately 350 and 500 GeV. When decreasing the relative branching ratio
1271 to DM ($g_{\text{DM}} = 0.5$), DM masses down to 1 GeV are excluded up to a pseudoscalar mediator
1272 mass of 470 GeV. Using Eqs. 7-8, Figure 79 shows that the gain is larger in the pseudoscalar case
1273 when compared to the scalar mediator due to a steeper decrease in $t\bar{t}$ branching ratio for the
1274 scalar mediator.

DRAFT




Figure 78: Expected and observed 95% CL exclusions in the plane of m_{DM} - $m_{mediator}$ for scalar (left) and pseudoscalar (right) mediators. The top row shows two exclusions: the nominal assumption ($g_{DM} = g_{SM} = 1$), and an alternate assumption ($g_{DM} = 0.5, g_{SM} = 1$). Calculated cross-section times branching ratio values for the first assumption are shown as markers with accompanying values in units of fb. The middle row shows the fraction of the total production cross-section attributed to single top processes. The the bottom row is identical to the top row but without cross-section value/markers for presentational purposes.




Figure 79: Cross-section times branching ratio to $t\bar{t}$ for scalar and pseudoscalar mediators with mass 450 GeV, shown at DM coupling values of $g_{\text{DM}} = 1$ and $g_{\text{DM}} = 0.5$, calculated with Eqs. 7-8.

References

- 1275
- 1276 [1] RA5 group, “Search for SUSY in same-sign dilepton events at 13 TeV”, *CMS Physics*
1277 *Analysis Note AN-2016/228* (2016).
- 1278 [2] CMS Collaboration, “Search for physics beyond the standard model in events with two
1279 leptons of same sign, missing transverse momentum, and jets in proton-proton collisions
1280 at $\sqrt{s} = 13$ TeV”, arXiv:1704.07323.
- 1281 [3] J. Alwall et al., “The automated computation of tree-level and next-to-leading order
1282 differential cross sections, and their matching to parton shower simulations”, *JHEP* **07**
1283 (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- 1284 [4] tttt same-sign and multilepton group, “Search for SM tttt in the same-sign dilepton and
1285 multi-lepton final states at sqrt(s) = 13 TeV”, *CMS Physics Analysis Note AN-2017/115*
1286 (2016).
- 1287 [5] CMS Collaboration, “Search for standard model production of four top quarks with
1288 same-sign and multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Eur.*
1289 *Phys. J.* **C78** (2018), no. 2, 140, doi:10.1140/epjc/s10052-018-5607-5,
1290 arXiv:1710.10614.
- 1291 [6] R. Frederix, D. Pagani, and M. Zaro, “Large NLO corrections in $t\bar{t}W^\pm$ and $t\bar{t}\bar{t}$
1292 hadroproduction from supposedly subleading EW contributions”, *JHEP* **02** (2018) 031,
1293 doi:10.1007/JHEP02(2018)031, arXiv:1711.02116.
- 1294 [7] <https://twiki.cern.ch/twiki/bin/view/CMS/MissingETOOptionalFilters>.
- 1295 [8] M. Franco Sevilla and A. Ovcharova, “Isr reweighting recommendations for moriond
1296 2017”, December, 2016.
1297 [https://indico.cern.ch/event/592621/contributions/2398559/
1298 attachments/1383909/2105089/16-12-05_ana_manuelf_isr.pdf](https://indico.cern.ch/event/592621/contributions/2398559/attachments/1383909/2105089/16-12-05_ana_manuelf_isr.pdf).
- 1299 [9] CMS SUSY fake-leptons working group, “Studies of methods to estimate the non-prompt
1300 lepton background to searches for new physics”, *CMS Physics Analysis Note*
1301 *AN-2014/261* (2015).
- 1302 [10] P. Pigard, “Multivariate electron id in 8x”, June, 2016.
1303 [https://indico.cern.ch/event/482674/contributions/2206032/
1304 attachments/1292177/1931287/20160621_EGM_cms_week_v5.pdf](https://indico.cern.ch/event/482674/contributions/2206032/attachments/1292177/1931287/20160621_EGM_cms_week_v5.pdf).
- 1305 [11] G. Zevi Della Porta, “Lepton id for the full 2016 dataset”, November, 2016.
1306 [https://indico.cern.ch/event/590228/contributions/2380031/
1307 attachments/1375541/2088587/EGMSUS_newIDs_17Nov16.pdf](https://indico.cern.ch/event/590228/contributions/2380031/attachments/1375541/2088587/EGMSUS_newIDs_17Nov16.pdf).
- 1308 [12] V. H. et al., “Electron mva id for susy”, April, 2017.
1309 [https://indico.cern.ch/event/719317/contributions/2963816/
1310 attachments/1630110/2598062/MVAidSUSY_10Apr18_SUSYMeeting.pdf](https://indico.cern.ch/event/719317/contributions/2963816/attachments/1630110/2598062/MVAidSUSY_10Apr18_SUSYMeeting.pdf).
- 1311 [13] S. L. S. F. team, “Lepton scale factors”, July, 2016.
1312 <https://twiki.cern.ch/twiki/bin/view/CMS/SUSLeptonSF>.
- 1313 [14] Baffioni, S. and others, “Electron Charge Identification using 8 TeV data”, *CMS Physics*
1314 *Analysis Note AN-2014/164* (2015).

- 1315 [15] G. Abbiendi et al., “Baseline muon selections”, October, 2014. https://twiki.cern.ch/twiki/bin/view/CMSPublic/SWGuideMuonId#Tight_Muon.
1316
- 1317 [16] G. Petrucciani and C. Botta, “Two step prompt muon identification”, January, 2015.
1318 <https://indico.cern.ch/event/368007/contribution/2/material/slides/0.pdf>.
1319
- 1320 [17] RA5 group, “Search for SUSY in same-sign dilepton events at 13 TeV”, *CMS Physics*
1321 *Analysis Note AN-2015/031* (2016).
- 1322 [18] D. Ferencek et al., “b-tagging offline guide”, November, 2014.
1323 <https://twiki.cern.ch/twiki/bin/view/CMSPublic/SWGuideBTagging>.
- 1324 [19] CMS Collaboration, “Performance of the missing transverse energy reconstruction by the
1325 CMS experiment in $\sqrt{s} = 8$ TeV pp data”, (2014). arXiv:1411.0511. Submitted to
1326 *JINST*.
- 1327 [20] [https://twiki.cern.ch/twiki/bin/view/CMS/](https://twiki.cern.ch/twiki/bin/view/CMS/MissingETUncertaintyPrescription#Instructions_for_9_4_X_X_9_for_2)
1328 [MissingETUncertaintyPrescription#Instructions_for_9_4_X_X_9_for_2](https://twiki.cern.ch/twiki/bin/view/CMS/MissingETUncertaintyPrescription#Instructions_for_9_4_X_X_9_for_2).
- 1329 [21] R. Barlow and C. Beeston, “Fitting using finite monte carlo samples”, *Computer Physics*
1330 *Communications* **77** (1993), no. 2, 219 – 228,
1331 doi:[https://doi.org/10.1016/0010-4655\(93\)90005-W](https://doi.org/10.1016/0010-4655(93)90005-W).
- 1332 [22] [https://twiki.cern.ch/twiki/bin/view/CMS/](https://twiki.cern.ch/twiki/bin/view/CMS/SWGuideHiggsAnalysisCombinedLimit)
1333 [SWGuideHiggsAnalysisCombinedLimit](https://twiki.cern.ch/twiki/bin/view/CMS/SWGuideHiggsAnalysisCombinedLimit).
- 1334 [23] K. J. F. Gaemers and F. Hoogeveen, “Higgs Production and Decay Into Heavy Flavors
1335 With the Gluon Fusion Mechanism”, *Phys. Lett.* **B146** (1984) 347–349,
1336 doi:10.1016/0370-2693(84)91711-8.
- 1337 [24] G. C. Branco et al., “Theory and phenomenology of two-Higgs-doublet models”, *Phys.*
1338 *Rept.* **516** (2012) 1–102, doi:10.1016/j.physrep.2012.02.002,
1339 arXiv:1106.0034.
- 1340 [25] D. Dicus, A. Stange, and S. Willenbrock, “Higgs decay to top quarks at hadron colliders”,
1341 *Phys. Lett.* **B333** (1994) 126–131, doi:10.1016/0370-2693(94)91017-0,
1342 arXiv:hep-ph/9404359.
- 1343 [26] N. Craig et al., “The Hunt for the Rest of the Higgs Bosons”, *JHEP* **06** (2015) 137,
1344 doi:10.1007/JHEP06(2015)137, arXiv:1504.04630.
- 1345 [27] N. Craig et al., “Heavy Higgs Bosons at Low $\tan\beta$: from the LHC to 100 TeV”,
1346 arXiv:1605.08744.
- 1347 [28] Q.-H. Cao, S.-L. Chen, and Y. Liu, “Probing higgs width and top quark yukawa coupling
1348 from $t\bar{t}h$ and $t\bar{t}t\bar{t}$ productions”, *Phys. Rev. D* **95** (Mar, 2017) 053004,
1349 doi:10.1103/PhysRevD.95.053004.
- 1350 [29] G. Bevilacqua and M. Worek, “Constraining BSM Physics at the LHC: Four top final
1351 states with NLO accuracy in perturbative QCD”, *JHEP* **07** (2012) 111,
1352 doi:10.1007/JHEP07(2012)111, arXiv:1206.3064.
- 1353 [30] E. Alvarez et al., “Four tops for lhc”, *Nuclear Physics B* **915** (2017) 19 – 43,
1354 doi:<https://doi.org/10.1016/j.nuclphysb.2016.11.024>.

- 1355 [31] C. Englert, G. F. Giudice, A. Greljo, and M. McCullough, “The \hat{H} -Parameter: An Oblique
1356 Higgs View”, arXiv:1903.07725.
- 1357 [32] CMS Collaboration, “Search for Dark Matter Particles Produced in Association with a
1358 Top Quark Pair at $\sqrt{s}=13\text{TeV}$ ”, *Phys. Rev. Lett.* **122** (Jan, 2019) 011803,
1359 doi:10.1103/PhysRevLett.122.011803.
- 1360 [33] D. Pinna, A. Zucchetta, M. R. Buckley, and F. Canelli, “Single top quarks and dark
1361 matter”, *Phys. Rev. D* **96** (Aug, 2017) 035031, doi:10.1103/PhysRevD.96.035031.
- 1362 [34] CMS Collaboration, “Search for dark matter produced in association with a single top
1363 quark or a top quark pair in proton-proton collisions at $\sqrt{s}=13\text{TeV}$ ”, *Journal of High
1364 Energy Physics* **2019** (Mar, 2019) 141, doi:10.1007/JHEP03(2019)141.
- 1365 [35] e. a. T. Bose, “Search for dark matter in single top + MET final states”, *CMS Physics
1366 Analysis Note* **AN-2017/327** (2017).
- 1367 [36] J. Steggeman, “Taudid for 13 tev run: recommendation from the tau pog (version 43)”,
1368 March, 2018. [https://twiki.cern.ch/twiki/bin/view/CMS/
1369 TauIDRecommendation13TeV?rev=43#New_Tau_Isolation_Discriminators.](https://twiki.cern.ch/twiki/bin/view/CMS/TauIDRecommendation13TeV?rev=43#New_Tau_Isolation_Discriminators)

A Statistical checks

A.1 Impacts

The leading 30 nuisance impacts for two sets of impacts, expected and observed for cut-based and BDT analyses, are shown in Figure 117 (expected cut-based analysis), Figure 118 (observed cut-based analysis), Figure 119 (expected BDT analysis), and Figure 120 (observed BDT analysis). The leading expected nuisance in both cases corresponds to the $\sigma(\text{ttbb})/\sigma(\text{ttjj})$ scaling. Note that the “prop binSS” nuisances for MC statistics include (and are dominated by) tight-loose sideband statistics.

The observed pulls show the most constrained/pulled nuisances correspond to normalization parameters for ttW and ttZ, as we would expect from the control regions. “TTWSF” is moved by approximately 1σ (0.8σ) with respect to the input nuisance sizes for the cut-based (BDT) analysis. “TTZSF” is moved up by approximately 0.6σ (0.7σ) for the cut-based (BDT) analysis.




Figure 80: Expected nuisance impacts for the cut-based analysis.




Figure 81: Observed nuisance impacts for the cut-based analysis.




Figure 82: Expected nuisance impacts for the BDT-based analysis.




Figure 83: Observed nuisance impacts for the BDT-based analysis.

1382 **A.2 Nuisance forms**

1383 Nuisance functional forms and widths are tabulated in Table 30.

Table 30: Rows preceded by (s) apply to signal as well. For “shape” uncertainties, width is not applicable as up and down variation envelopes are taken as the systematic uncertainties.

name	function	width
lumi	lnN	1.021-1.05
(s) jes	shape	-
(s) jer	shape	-
isr	shape	-
bb	shape	-
(s) lep	shape	-
(s) trig	shape	-
(s) btaghf	shape	-
(s) btaglf	shape	-
(s) isrvar	shape	-
(s) fsrvar	shape	-
(s) scale	shape	-
(s) pdf	shape	-
(s) alphas	shape	-
(s) pu	shape	-
TTWSF	lnN	1.4
TTZSF	lnN	1.4
TTH	lnN	1.25
TTVV	lnN	1.11
XG	lnN	1.11
rares	lnN	1.2
fakes	lnN	1.3
fakes_EWK	shape	-
flips	lnN	1.2

A.3 Nuisances

1384
1385 Two sets of nuisance pull values, expected and observed for cut-based and BDT analyses, are
1386 tabulated in Table 62 (expected cut-based analysis), Table 63 (observed cut-based analysis),
1387 Table 64 (expected BDT analysis), and Table 65 (observed BDT analysis).

1388 The most constrained nuisances correspond to normalization parameters for ttW and ttZ (
1389 “TTWSF” and “TTZSF”) due to high statistics in control regions and in the bulk (BDT). Their
1390 input normalization uncertainty is 40%.

DRAFT

Table 31: Expected nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s</i> + <i>b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.38, 0.99	+0.00, 0.97	-0.12
TTVV	+0.05, 1.00	+0.00, 0.99	-0.02
TTWSF	+0.09, 0.61	+0.00, 0.61	-0.06
TTZSF	-0.02, 0.45	+0.00, 0.45	+0.02
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	-0.00, 0.99	+0.00
bb	+0.74, 0.94	+0.00, 0.98	-0.24
fakes	+0.24, 0.92	+0.00, 0.92	-0.08
pdf	+0.07, 0.99	-0.00, 0.99	-0.02
rares	-0.01, 0.99	+0.00, 0.99	+0.00
scale	-0.19, 0.96	-0.00, 0.99	+0.00
y2016_btaghf	+0.04, 0.99	+0.00, 0.99	-0.03
y2016_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_fakes_EWK	+0.04, 1.00	+0.00, 0.99	-0.01
y2016_flips	-0.02, 0.99	+0.00, 0.99	+0.00
y2016_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.02
y2016_isr	-0.10, 0.99	-0.00, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	+0.00, 0.99	-0.02
y2016_jer	-0.13, 1.12	-0.00, 0.99	+0.03
y2016_jes	+0.07, 0.97	+0.00, 0.99	-0.05
y2016_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2016_lumi	+0.02, 0.99	+0.00, 0.99	-0.02
y2016_prefire	-0.00, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.01, 0.99	-0.00, 0.99	+0.00
y2016_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_btaghf	+0.04, 0.99	+0.00, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	+0.00, 0.99	-0.00, 0.98	-0.01
y2017_flips	-0.02, 0.99	+0.00, 0.99	+0.00
y2017_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2017_isr	+0.07, 0.99	+0.00, 0.99	-0.03
y2017_isrvar	+0.00, 0.99	-0.00, 0.99	-0.01
y2017_jer	-0.12, 1.20	-0.00, 0.99	+0.03
y2017_jes	+0.07, 0.99	+0.00, 0.99	-0.05
y2017_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2017_lumi	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_prefire	-0.01, 0.99	-0.00, 0.99	+0.01
y2017_pu	-0.01, 1.00	+0.00, 0.99	+0.00
y2017_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2018_btaghf	+0.09, 0.99	+0.00, 0.99	-0.05
y2018_btaglf	+0.12, 0.99	-0.00, 0.99	-0.04
y2018_fakes_EWK	+0.08, 1.03	+0.00, 0.99	-0.02
y2018_flips	-0.03, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_isr	+0.08, 0.99	+0.00, 0.99	-0.03
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_jer	-0.33, 1.08	-0.00, 0.99	+0.08
y2018_jes	+0.18, 0.96	-0.00, 0.98	-0.11
y2018_lep	+0.02, 0.99	+0.00, 0.99	-0.02
y2018_lumi	+0.03, 0.99	+0.00, 0.99	-0.04
y2018_pu	+0.00, 1.00	-0.00, 0.99	+0.00
y2018_trig	+0.02, 0.99	+0.00, 0.99	-0.03

Table 32: Observed nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	
TTH	+0.48, 1.02	+0.24, 1.00	-0.12
TTVV	+0.06, 1.00	+0.03, 1.00	-0.02
TTWSF	+1.03, 0.57	+0.99, 0.56	-0.06
TTZSF	+0.57, 0.41	+0.60, 0.42	+0.05
XG	+0.04, 1.00	+0.04, 1.00	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.62, 0.92	+0.10, 0.97	-0.27
fakes	+0.54, 1.02	+0.33, 1.00	-0.09
pdf	+0.15, 0.98	+0.06, 1.00	-0.04
rares	+0.06, 1.00	+0.07, 1.00	+0.01
scale	-0.08, 0.91	-0.02, 0.96	-0.00
y2016_btaghf	+0.07, 0.99	+0.04, 0.99	-0.03
y2016_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_fakes_EWK	+0.08, 1.00	+0.05, 0.99	-0.01
y2016_flips	+0.00, 0.99	+0.01, 1.00	+0.01
y2016_fsrvar	+0.00, 0.99	-0.01, 1.04	+0.01
y2016_isr	-0.12, 0.99	-0.04, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	-0.04, 1.06	-0.03
y2016_jer	-0.07, 1.04	+0.01, 0.99	+0.04
y2016_jes	+0.38, 1.08	+0.25, 1.12	-0.08
y2016_lep	+0.04, 0.99	+0.03, 0.99	-0.01
y2016_lumi	+0.05, 1.00	+0.04, 1.00	-0.02
y2016_prefire	-0.01, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.03, 0.99	+0.01, 0.99	-0.01
y2016_trig	+0.04, 0.99	+0.03, 0.99	-0.01
y2017_btaghf	+0.10, 1.00	+0.06, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	+0.23, 0.98	+0.22, 0.97	-0.01
y2017_flips	+0.00, 0.99	+0.01, 1.00	+0.01
y2017_fsrvar	+0.00, 0.99	-0.01, 0.99	+0.00
y2017_isr	+0.08, 0.99	+0.02, 0.99	-0.04
y2017_isrvar	+0.00, 0.99	-0.01, 0.99	-0.01
y2017_jer	-0.07, 1.24	+0.01, 0.96	+0.03
y2017_jes	+0.15, 0.93	+0.09, 0.93	-0.05
y2017_lep	+0.04, 0.99	+0.03, 0.99	-0.01
y2017_lumi	+0.05, 1.00	+0.04, 1.00	-0.01
y2017_prefire	-0.02, 0.99	-0.02, 0.99	+0.01
y2017_pu	+0.03, 1.00	+0.01, 1.00	-0.00
y2017_trig	+0.04, 0.99	+0.04, 0.99	-0.01
y2018_btaghf	+0.15, 0.99	+0.07, 0.99	-0.05
y2018_btaglf	+0.12, 0.99	+0.02, 0.99	-0.05
y2018_fakes_EWK	+0.15, 1.01	+0.09, 0.99	-0.02
y2018_flips	+0.01, 0.99	+0.02, 1.00	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_isr	+0.08, 0.99	+0.02, 0.99	-0.04
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_jer	-0.19, 1.17	+0.06, 1.05	+0.09
y2018_jes	+0.48, 1.02	+0.31, 1.08	-0.13
y2018_lep	+0.06, 0.99	+0.05, 0.99	-0.02
y2018_lumi	+0.10, 0.99	+0.08, 0.99	-0.03
y2018_pu	+0.00, 1.00	-0.00, 0.99	+0.00
y2018_trig	+0.08, 0.99	+0.07, 0.99	-0.02

Table 33: Expected nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s</i> + <i>b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.32, 0.98	+0.00, 0.96	-0.11
TTVV	+0.06, 1.00	+0.00, 0.99	-0.02
TTWSF	+0.08, 0.63	+0.00, 0.62	-0.08
TTZSF	-0.04, 0.51	+0.00, 0.48	+0.03
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	-0.00, 0.99	-0.00, 0.99	+0.00
bb	+0.91, 0.93	+0.00, 0.99	-0.27
fakes	+0.28, 0.91	+0.00, 0.90	-0.08
pdf	+0.12, 0.97	-0.00, 0.99	-0.03
rares	-0.02, 0.99	+0.00, 0.99	+0.00
scale	-0.25, 0.97	+0.00, 0.99	+0.01
y2016_btaghf	+0.04, 0.99	+0.00, 0.99	-0.03
y2016_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_fakes_EWK	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_flips	-0.02, 0.99	+0.00, 0.99	+0.00
y2016_fsrvar	-0.00, 0.99	+0.00, 0.99	+0.01
y2016_isr	-0.13, 0.99	+0.00, 0.99	+0.04
y2016_isrvar	-0.00, 0.99	-0.00, 0.99	-0.01
y2016_jer	-0.05, 1.03	+0.00, 0.99	+0.01
y2016_jes	+0.07, 0.90	+0.00, 0.99	-0.04
y2016_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2016_lumi	+0.02, 0.99	+0.00, 0.99	-0.02
y2016_prefire	-0.00, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.04, 0.99	-0.00, 0.99	-0.01
y2016_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_btaghf	+0.03, 0.99	+0.00, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	+0.03, 0.98	+0.00, 0.97	-0.02
y2017_flips	-0.02, 0.99	+0.00, 0.99	+0.01
y2017_fsrvar	-0.00, 0.99	-0.00, 0.99	+0.00
y2017_isr	+0.07, 0.99	-0.00, 0.99	-0.02
y2017_isrvar	-0.00, 0.99	+0.00, 0.99	-0.01
y2017_jer	-0.01, 1.19	+0.00, 0.99	+0.00
y2017_jes	+0.07, 0.90	+0.00, 0.99	-0.04
y2017_lep	+0.00, 0.99	+0.00, 0.99	-0.02
y2017_lumi	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_prefire	-0.00, 0.99	-0.00, 0.99	+0.01
y2017_pu	+0.03, 1.00	-0.00, 0.99	-0.01
y2017_trig	+0.00, 0.99	+0.00, 0.99	-0.02
y2018_btaghf	+0.08, 0.99	+0.00, 0.99	-0.05
y2018_btaglf	+0.12, 0.99	-0.00, 0.99	-0.05
y2018_fakes_EWK	+0.06, 0.98	+0.00, 0.98	-0.01
y2018_flips	-0.03, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	-0.00, 0.99	+0.00, 0.99	+0.00
y2018_isr	+0.04, 0.99	-0.00, 0.99	-0.02
y2018_isrvar	-0.00, 0.99	+0.00, 0.99	+0.00
y2018_jer	-0.01, 1.56	+0.00, 0.95	+0.00
y2018_jes	+0.15, 0.80	-0.00, 0.95	-0.10
y2018_lep	+0.01, 0.99	+0.00, 0.99	-0.02
y2018_lumi	+0.03, 0.99	+0.00, 0.99	-0.04
y2018_pu	+0.02, 1.01	-0.00, 0.99	-0.01
y2018_trig	+0.02, 0.99	+0.00, 0.99	-0.03

Table 34: Observed nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.57, 1.02	+0.33, 1.00	-0.09
TTVV	+0.08, 1.00	+0.04, 1.00	-0.02
TTWSF	+0.64, 0.66	+0.75, 0.60	-0.01
TTZSF	+0.61, 0.42	+0.65, 0.43	+0.04
XG	+0.02, 0.99	+0.04, 1.00	+0.00
alphas	+0.00, 0.99	+0.00, 1.00	+0.00
bb	+0.87, 0.95	+0.07, 0.98	-0.27
fakes	+0.93, 0.99	+0.32, 0.99	-0.14
pdf	+0.05, 0.99	-0.03, 1.01	-0.02
rares	+0.03, 0.99	+0.04, 0.99	+0.00
scale	-0.24, 0.94	-0.04, 0.94	+0.00
y2016_btaghf	+0.08, 0.99	+0.05, 0.99	-0.02
y2016_btaglf	+0.04, 1.00	+0.01, 0.99	-0.01
y2016_fakes_EWK	+0.16, 0.99	+0.08, 0.99	-0.02
y2016_flips	-0.01, 0.99	+0.00, 1.00	+0.00
y2016_fsrvar	+0.00, 0.99	+0.02, 1.07	+0.02
y2016_isr	-0.18, 0.99	-0.06, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	+0.01, 1.00	-0.01
y2016_jer	-0.06, 0.89	-0.03, 0.87	+0.01
y2016_jes	+0.03, 1.00	-0.04, 1.11	-0.03
y2016_lep	+0.04, 0.99	+0.03, 0.99	-0.01
y2016_lumi	+0.05, 1.00	+0.04, 1.00	-0.02
y2016_prefire	-0.01, 0.99	-0.00, 0.99	+0.00
y2016_pu	-0.01, 0.99	-0.05, 1.00	-0.01
y2016_trig	+0.05, 0.99	+0.04, 0.99	-0.02
y2017_btaghf	+0.09, 0.99	+0.07, 0.99	-0.03
y2017_btaglf	+0.05, 0.99	+0.02, 0.99	-0.01
y2017_fakes_EWK	+0.43, 1.04	+0.25, 0.97	-0.04
y2017_flips	-0.01, 0.99	+0.01, 1.00	+0.00
y2017_fsrvar	+0.00, 0.99	+0.02, 1.00	+0.01
y2017_isr	+0.08, 0.99	+0.02, 0.99	-0.02
y2017_isrvar	+0.00, 0.99	+0.00, 1.00	-0.01
y2017_jer	+0.09, 0.87	+0.07, 0.83	-0.01
y2017_jes	+0.14, 0.89	+0.09, 0.99	-0.04
y2017_lep	+0.05, 0.99	+0.04, 0.99	-0.01
y2017_lumi	+0.05, 1.00	+0.05, 1.00	-0.02
y2017_prefire	-0.01, 0.99	-0.01, 0.99	+0.01
y2017_pu	+0.05, 1.00	+0.03, 0.99	-0.01
y2017_trig	+0.04, 0.99	+0.04, 0.99	-0.01
y2018_btaghf	+0.13, 0.99	+0.09, 0.99	-0.05
y2018_btaglf	+0.05, 0.99	-0.03, 0.99	-0.04
y2018_fakes_EWK	+0.10, 0.98	-0.01, 0.99	-0.03
y2018_flips	-0.02, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_isr	+0.04, 0.99	+0.01, 0.99	-0.02
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_jer	+0.74, 0.77	+0.50, 0.97	-0.14
y2018_jes	+0.21, 0.67	+0.14, 0.81	-0.07
y2018_lep	+0.05, 0.99	+0.05, 0.99	-0.02
y2018_lumi	+0.08, 0.99	+0.07, 0.99	-0.03
y2018_pu	+0.10, 1.01	+0.06, 0.99	-0.02
y2018_trig	+0.05, 0.99	+0.05, 0.99	-0.03

1391 **A.4 Nuisance correlation matrix**

1392 Using the combine tool (via `combine -M FitDiagnostics card.txt --robustFit=1`
 1393 `--saveShapes --saveWithUncertainties --saveOverallShapes --numToysForShapes`
 1394 `200`), we can show the correlations between different nuisance parameters in the fit, in Figs. 84-
 1395 85.

Correlation matrix of fit parameters (cut-based)




Figure 84: Correlation plot of nuisance-vs-nuisance for the s+b fit for the cut-based regions

Correlation matrix of fit parameters (BDT)




Figure 85: Correlation plot of nuisance-vs-nuisance for the s+b fit for the BDT regions

1396 **A.5 Goodness of fits**

1397 The goodness of fit distributions (using the saturated, Kolmogorov-Smirnov, and Anderson-Darling test statistics) with the signal+background fit to data for the cut-based and BDT analyses are shown in Figure 121. We note that the observation is generally within the bulk of the expected distributions for both analyses and all three metrics.

1398

1399




Figure 86: GOF test for the cut-based analysis (left) and BDT-based analysis (right) using the saturated (top), KS (middle), AD (bottom) test statistics.

B Shape variations

B.1 BDT discriminator

Shape variations for the BDT discriminator for the largest backgrounds and signal, for b-tagging, JES, JER, PU, prefiring, ISR/FSR parton shower, and ISR jet reweighting systematics are shown in Figures 87, 88, 89, and 90. Note that complete MC is used with events normalized to 136.3 fb^{-1} .




Figure 87: Systematic variations for ttW




Figure 88: Systematic variations for ttZ




Figure 89: Systematic variations for ttH




Figure 90: Systematic variations for tttt signal

C Studies which did not result in analysis changes

C.1 Jet and b-jet thresholds

We studied whether the $t\bar{t}\bar{t}$ signal significance would improve by lowering the N_{jets} and $N_{\text{b jets}}$ counting thresholds from the 2016 ones which were 40 and 25 GeV, respectively. To do this, we used 2016 MC to estimate the expected significance of the analysis with an expected luminosity of 75 fb^{-1} , using the 2016+2017 cut based signal region binning. The result, in Figure 91, shows that the current configuration is close to optimal, with only a 3% increase in significance which could be obtained from reducing the $N_{\text{b jets}}$ threshold from 25 to 20 GeV. Given the minor improvement, and the potential of larger b-tagging scale factors at low p_T in the 2017 data, we decided to keep the 2016 thresholds.




Figure 91: Signal significance for different transverse momentum requirements used in counting N_{jets} and $N_{\text{b jets}}$.

C.2 Taus

We studied whether the $t\bar{t}\bar{t}$ signal significance would improve by including taus in addition to electrons and muons. For this study, we used an expected luminosity of 75 fb^{-1} , and we plotted the expected yields for events with reconstructed tau leptons, separating the truth-matched and non-truth-matched ones.

We focused on events with only one tau, and the other 1 or 2 leptons being e/μ , to avoid large yields of fake taus. Taus were selected with the following identification requirements: `decayModeFinding`, `againstMuonTight3`, `againstElectronTightMVA6`, `byTightIsolationMVArun2v1DBdR03oldDMwLT`. The last requirement was developed by the $t\bar{t}H$ analysis to be used in environments with high jet multiplicity [36]. The performance of tau identification is shown in Figure 92 for the $t\bar{t}\bar{t}$ signal and the $t\bar{t}W/Z/H$ background samples, where it is clear that there are large off-diagonal terms due to inefficiency and impurity of

1429 the tau selection. Figure 93 shows the expected yields for each sample as a function of signal
 1430 region for 1-tau events, split between $t\bar{t}\bar{\tau}$ events where the tau is truth-matched, and the sum of
 1431 $t\bar{t}W/Z/H$ events and $t\bar{t}\bar{\tau}$ events with fake taus. The plot shows some potential in the 3-lepton
 1432 regions ($e\bar{e}\tau/e\mu\tau/\mu\mu\tau$), particularly 14-15-16, which integrate to about 0.5 $t\bar{t}\bar{\tau}$ events and 1
 1433 background event, but this doesn't account for fake taus from non- $t\bar{t}W/Z/H$ backgrounds.
 1434 Given these small yields and large backgrounds, we decided not to include signal regions with
 1435 taus for the 2016+2017 analysis.




Figure 92: Number of events as a function of the number of true and reconstructed tau leptons, for different samples.




Figure 93: Top: number of events with 1 tau lepton and 1 or 2 e/μ leptons, as a function of the signal regions defined in Table 10, with SR11 and higher including events with 3 leptons ($ee\tau/e\mu\tau/\mu\mu\tau$), showing truth-matched $t\bar{t}\bar{t}$ signal (orange) and backgrounds from $t\bar{t}W/Z/H$ and non-truth-matched $t\bar{t}\bar{t}$ (blue). Middle and bottom: S/B and significance per bin, considering only the $t\bar{t}W/Z/H$ and non-truth-matched $t\bar{t}\bar{t}$ events as background.

C.3 Top-tagging

We explored resolved and merged top tagging as additional discrimination handles.

C.3.1 Resolved

Pairs of 3 jets, where one jet is a b-jet candidate, are fed into a pre-trained BDT from the single lepton and opposite-sign analysis, AN2017-146-v17. The BDT uses 6 inputs (b-tag discriminant, top candidate mass, W candidate mass, the ratio of the top p_T to the trijet p_T , and the $\Delta\phi$ between (top, W) and (top,b)). We can find the highest and second highest trijet discriminant in the event. For this analysis, the leading backgrounds can all have a hadronic top, so we focus on the subleading trijet discriminant.

Distributions of this quantity for two jet threshold schemes are shown in Figure 94. The nominal analysis uses jet thresholds of 40 GeV and b-tagged jet thresholds of 25 GeV. To explore the possibility that the high jet threshold is reducing the inputs to the BDT, we compare with looser jet thresholds of 20 GeV, 20 GeV, respectively.

In both cases, the ratio of signal to background shows only a slight trend. To test this more quantitatively, the subleading discriminator for the 20,20 scheme was put into a 19+1 variable BDT and was only ranked 9th. Compared to without the variable, the maximum significance only increased by 1%, so we do not pursue such tagging in the analysis. This is not unexpected since the same-sign and multilepton final state has one or two less hadronic tops to tag.




Figure 94: Distributions of subleading trijet discriminator in the signal regions for (jet,b-jet) p_T thresholds of (40,25) [left] and (20,20) [right].

C.3.2 Merged

Top decay products may become merged through accidental overlaps or boosts. The latter is not a large fraction in this final state due to lack of high MET, however, we explored the DeepAK8 tagger developed by the SUSY Heavy Object Tagging group, which provides numerical probabilities for objects such as top, W , Z , H , etc. for AK8 jets through a deep neural network acting on constituent particle flow candidates.

Following the same strategy as for the resolved top tagger, we look at the leading and subleading top discriminants for events passing the baseline selection for a few leading backgrounds in Figure 95.

1463 Again, while there is a small trend toward higher values for the signal sample, the quantity
1464 of these events is miniscule (on the order of a few percent), so this tagging method was not
1465 pursued further.




Figure 95: Maximum AK8 top discriminant in the event (left). Subleading AK8 top discriminant in the event (right). Histograms are normalized to the signal (red) cross-section.

D Impact of the L1 ECAL prefiring(2016/2017) and HEM15/16 loss (2018) on the results

D.1 L1 prefiring(2016/2017)

The L1 prefiring issue impacts 2016 and 2017 data collection periods. Due to mistiming and trigger rules, events with high η energy deposits can be preferentially lost. The impact of this is checked on 80X fastsim T1tttt signal samples. Inefficiency maps to be applied to simulation are taken from https://ncsmith.web.cern.ch/ncsmith/PrefireEfficiencyMaps/Preliminary/Jet_L1FinOReff_bxml_looseJet_SingleMuon_Run2016B-H.pdf (Jet map 2016B-H) and <https://lathomas.web.cern.ch/lathomas/TSGStuff/L1Prefiring/PrefiringMaps/> (Jet map 2017B-F) and shown in Figure 96. For each event, consider all jets and electrons that pick up a non-zero scale factor from the chosen inefficiency map, parameterized by jet p_T and η and obtain a multiplicative scale factor (< 1) to apply to MC.

The procedure matches the tool given in <https://twiki.cern.ch/twiki/bin/view/CMS/L1ECALPrefiringWeightRecipe>.

Figures 97 and 98 show the average SFs as a function of (selected) electron and jet multiplicities per event, separately for 2016 and 2017. The average scale factors are about 2.5% (4%) below unity for 2016 (2017). Compared to the central value of the scale factors, the variation for increasing jet or electron multiplicity is small. Note that as a function of raw (not analysis/selected) jets, the trend may be larger, as the analysis jet selection criteria requires $|\eta| < 2.5$.

Scale factors for the prefiring inefficiency will be applied to the 2016 and 2017 MC samples.




Figure 96: Prefiring inefficiency maps for 2016 (left) and 2017 (right)




Figure 97: Average preferring inefficiency as a function of electron multiplicity for 2016 (left) and 2017 (right)




Figure 98: Average preferring inefficiency as a function of jet multiplicity for 2016 (left) and 2017 (right)

D.2 HEM15/16 loss in 2018

Beginning in Run 319077 for the 2018 data collection period, HEM 15 and 16 sectors are switched off due to a power cut issue. This results in a gap of HCAL info for $\eta \in [-3, 1.3]$ and $\phi \in [-1.57, -0.87]$. This has the potential to increase electron fakes due to lower H/E and isolation values. The impact was checked in MC and in data.

The relative increase of fake electrons passing tight selection requirements was found to be approximately 12% in MC, while muons showed no increase. Since 23% of the Run2 dataset is impacted by the HEM issue, and considering fakes inclusively ($e+\mu$), this increase corresponds to a 1% overall increase in fakes across the full dataset. Data checks showed similar increases to MC.

D.2.1 HEM impact in MC

Dedicated samples were produced to perform an apples-to-apples comparison of the HEM15/16 issue. In particular, the dataset

```
/TTToSemiLeptonic_TuneCP5_13TeV-powheg-pythia8
/RunIISpring18MiniAOD-HEMPremix_100X_upgrade2018_realistic_v10-v3
/MINIAODSIM
```

was produced with HEM sectors disabled. A sample without “HEM” in the dataset name corresponds to the nominal HEM “in” case. For both samples, truth-matched fake single lepton efficiencies were calculated with nominal analysis tight ID+Iso as the numerator and reco-leptons from miniAOD as the denominator. Leptons were required to have $p_T > 25$ GeV. Labeling HEM “in” and “out” samples as “good” and “bad”, both good and bad samples were checked inclusively and also restricting to the η - ϕ region impacted by the HEM loss. The efficiencies are listed in Table 35. Muons are not impacted, while electrons show a $(12 \pm 5)\%$ relative increase in fakes inclusively, or a factor of 2 increase in the HEM region specifically.

Table 35: Single lepton efficiencies for good and bad HEM samples.

region	scenario	flavor	sig. eff. (%)	bkg. eff. (%)
all	good	e	65.8 ± 0.1	0.15 ± 0.01
		μ	84.6 ± 0.1	0.31 ± 0.01
	bad	e	65.9 ± 0.1	0.17 ± 0.0
		μ	84.7 ± 0.1	0.29 ± 0.01
hemregion	good	e	53.9 ± 0.8	0.21 ± 0.04
		μ	80.7 ± 1.0	0.21 ± 0.06
	bad	e	59.7 ± 0.7	0.47 ± 0.03
		μ	82.3 ± 0.9	0.33 ± 0.07

D.2.2 HEM impact in data

To assess the impact in data, the 2018 dataset was split into two subsections: “before” (runs before 319077, corresponding to 20.25 fb^{-1}) and “after” (runs including and after 319077, corresponding to 30.73 fb^{-1}). After normalizing both sections to the same luminosity, plots of lepton flavor ($\mu\mu, e\mu, ee$) for 3 kinematic regions ($Z \rightarrow \ell\ell$ -dominated, $t\bar{t} \rightarrow \ell\ell$ -dominated, and the (fake-dominated) tight-loose control region) are shown in Figure 99. While the first two

1516 prompt-enriched regions show a relative increase afterwards, this could be attributed to a variety of data-taking condition differences between the two periods. The tight-loose region shows
 1517
 1518 similar counts for dimuon events, but after/before ratios of 1.13 ± 0.08 for $e\mu$ and 1.37 ± 0.20 for
 1519 dielectron events. Such an increase is consistent with the 12% increase found from simulation.

1520 Figure 100 shows the lepton ϕ distribution for Z-dominated and tight-loose regions. Muons
 1521 show no relative changes. When focusing on $\phi \in [-1.57, -0.87]$, no increase is observed for
 1522 prompt electrons, while a $(59 \pm 23)\%$ increase is seen for the tight-loose electron counts.




Figure 99: Z-dominated (left), $t\bar{t}$ -dominated (center), tight-loose (right).




Figure 100: Z-dominated region (left), tight-loose region (right), for electrons (top) and muons (bottom)

E Analysis changes

E.1 Changes from ANv7

This section details some changes and updates with respect to V7 of this note. Different items are accompanied by the relative change in expected significance from the ANV7 number (cut-based gave 3.069σ , and BDT gave 3.328σ).

However, it is worth noting that JECs for 2018 data and MC are not yet available, so 2017 JECs are being used instead. Several of the changes enumerated below and their conclusions are sensitive to JECs.

The compounded effect of these changes brings the cut-based significance to 2.371σ and the BDT-based significance to 2.591σ .

- V32 JEC

- With respect to the previous version of JECs (V6), V32 reparameterizes the L1 correction for better stability. While the product of L1,L2L3 remained the same from V6 to V32, the relative ratios did not. Our lepton isolation quantities, p_T^{ratio} and p_T^{rel} depend on these two corrections separately, not only on their product. Upgrading to V32 caused a per lepton efficiency loss of a few percent, so we re-tuned our isolation working points for 2017 and 2018 to approximately match the signal and background efficiencies from V6.

- The updated working point values can be found in Table 7.

- Relevant discussion in <https://hypernews.cern.ch/HyperNews/CMS/get/JetMET/1891.html>

- σ decreases by 4%.

- Prefiring SF

- The latest recommended prefiring inefficiency maps (from https://twiki.cern.ch/twiki/bin/view/CMS/SUSRecommendations18#Prefire_Issue) were incorporated for 2016 and 2017 MC. Application of the maps scales down four top signal by 2.5% (4.7%) in 2016 (2017) and the $t\bar{t}W$ background by 1.5% (3.3%) in 2016 (2017).

- After application of the inefficiency maps, disagreements in high- $|\eta|$ tails of the opposite-sign control region are largely cured.

- For reference, the affected datasets (2016+2017) are only 58% of the Run2 dataset by integrated luminosity.

- σ decreases by 1%.

- Merging 4b signal region bins

- Signal regions 8, 9, and 10 (now just 8), corresponding to at least 4 b-tagged jets and at least 5 jets, were merged (removing the jet multiplicity boundaries) to be more conservative when dealing with the tight-loose control region yields for the fake background prediction

- The cut-based signal region definition now has 14 signal regions (+2 $t\bar{t}W/t\bar{t}Z$ dominated regions, as before).

- σ decreases by 4%.

- Single card

- Card inputs to the HiggsCombine tool are no longer treated as 3 sepa-

- rate channels in a simultaneous fit. Instead, yields (including control region yields) are summed into single histograms (2016+2017+2018). Uncorrelated nuisances, for example, prefiring uncertainties affecting only 2016/2017, are considered as shape variations only on the 2016/2017 component of the summed nominal histogram. This has the benefit of not being as susceptible to MC statistics issues for each of the 3 years. It also is more conservative and takes advantage of larger control region statistics for the fake prediction.
- σ decreases by 5%.
 - 2017/2018 MC
 - The Autumn18 MC campaign is nearly complete, so we have switched to taking relevant 2018 predictions from the correct MC rather than using 2017 as a stand-in.
 - Explicitly, we have all the desired samples in the 2018 campaign except the signal sample is still lacking 25% MC statistics. This does not matter as we don't suffer MC statistics issues for $t\bar{t}\bar{t}$.
 - In 2017, we are still lacking a very small $Z+\gamma$ background sample, which we take from the 2018 campaign instead.
 - σ decreases by 3%.
 - 2017/2018 ISR reweighting
 - For the 2016 dataset, we reweight the $t\bar{t}W$ and $t\bar{t}Z$ background predictions in order to match the data $N_J^{\text{ISR/FSR}}$ distribution, as detailed in Section 2. This had the effect of scaling down high jet multiplicity events, decreasing the background prediction. In 2017 and 2018, however, the Pythia tune was changed to CP5, so we requested dedicated samples with configurations matching $t\bar{t}W$ and $t\bar{t}Z$ in order to derive new weights. This results in the opposite trend to 2016, i.e., high jet multiplicity events can get scaled up by almost 30%, increasing the background prediction.
 - In 2017/2018, the samples for $t\bar{t}Z$ and $t\bar{t}W$ with 0 and 1 extra parton, respectively, are
 - /TT_DiLept_TuneCP5_13TeV-amcatnlo-pythia8/
 - /TTPlus1Jet_DiLept_TuneCP5_13TeV-amcatnloFXFX-pythia8/
 which are completed by the appropriate processing string for each campaign. In the case of 2017, full event statistics are split between a nominal sample and an extension ("ext1") sample.
 - σ decreases by 2%.
 - 2018 b-tag WPs
 - The 2018 b-tag medium WP for DeepCSV preserves the b tagging and light mis-tagging efficiencies compared to 2017. However, it appears the charm mis-tagging efficiency relatively increased by about 15%. In the 3 and 4 b-tag bins, $t\bar{t}W$ can enter if there is additional heavy flavor contribution (e.g., $t\bar{t}W +bb$ via gluon splitting), or through mistags of W decays containing a charm quark.
 - Relevant discussion in <https://hypernews.cern.ch/HyperNews/CMS/get/btag/1637.html>
 - σ decreases by 3%.

- 1613 • ttW+bb, ttZ+bb, ttH+bb scaling
- 1614 • As a response to a 7 Dec 2018 question from Otto about Figs 12, 13 (<https://twiki.cern.ch/twiki/bin/viewauth/CMS/TOP18003>), we showed
- 1615 that scaling the gluon splitting (events with extra $b\bar{b}$) components up by
- 1616 the systematic uncertainty of 1.35 brings better agreement in the b-tag
- 1617 multiplicity distribution.
- 1618
- 1619 • We now revert to the procedure of the previous iteration of this analysis
- 1620 for 2016 by scaling $(t\bar{t}W/t\bar{t}Z/t\bar{t}H)+b\bar{b}$ up by 1.7 ± 0.6 with a systematic
- 1621 uncertainty corresponding to the specified error, from the result of TOP-
- 1622 16-010, which found a Data/MC discrepancy in the ttbb/ttj ratio.
- 1623 • σ decreases by 6%.

DRAFT

F Unblinding of 2016 dataset

Following the pre-approval talk for the analysis on Feb. 15, 2019, the analysis was unblinded using the 2016 dataset. The yields and results are shown in Section F.1 with consistency checks of cut-based and BDT results in Section F.2. We include the results of various statistical checks, including nuisance pulls (Section F.3), and nuisance impacts (Section F.4).

F.1 Yields and results

Plots for the unblinded prefit and postfit event yields for the 2016 data, with a total luminosity of 35.9 fb^{-1} , are shown for both the cut-based and BDT based analysis in Figure 101. Numerical yields are also tabulated in Table 36 (prefit cut-based analysis), Table 37 (postfit cut-based analysis), Table 38 (prefit BDT analysis), Table 39 (postfit BDT analysis).

A binned likelihood fit is performed over signal regions using the Higgs Combine tool [22]; exclusion limits at 95% CL are calculated with the Asymptotic CLs method.

With the cut-based analysis, an observed (expected) upper limit on the production cross section of 33.67 fb ($20.48_{-6.83}^{+11.18} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 1.012 (1.374) standard deviations, corresponding to a measured signal strength parameter of $0.812_{-0.800}^{+1.054}$.

With the BDT analysis, an observed (expected) upper limit on the production cross section of 42.59 fb ($19.36_{-6.46}^{+10.11} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 1.812 (1.424) standard deviations, corresponding to a measured signal strength parameter of $1.538_{-0.935}^{+1.137}$.

F.2 Cut-based and BDT consistency checks

One noteworthy excess for the cut-based is SR4, which has 8 observed events with about 2.2 ± 0.5 predicted background. This is a 2-lepton, 5 jet, 3 b-tagged jet bin. In the 2016 result, this corresponded to part of oldSR4 (2-lepton, 5 or 6 jets, 3 b-tagged jets), and had 8 observed events. With the new binning which splits oldSR4 into SR4 and SR5, the data events remain in the 5 jet bin, SR4, leaving the slightly higher s/b bin SR5 with 0 data.

It is also worth pointing out the mildness of the disagreement between cut-based and BDT significances (both have differing directions for observed significance with respect to their expected significance). We have checked with toy pseudo-datasets that the probability of this happening is close to 50%.

The asimov-like toys were constructed in a way that would preserve the correlation between the cut-based and BDT significance results. The procedure for creating one toy is to consider all background events for the analysis, storing the cut-based and BDT SR bin number and event weight. For each background/signal process, calculate the total yield y in the cut-based/BDT signal regions and sample a number N from a Poisson distribution with parameter y . Next, draw N random events from the set of events for a particular process with relative probability given by the absolute value of the event weight (so there is a slight bias due to negatively weighted events). These N sampled events then are used to fill a histogram for the cut-based and BDT shapes. In this way, we create $\mu \cdot s + b$ toys with $\mu = 1$, which is then an input to HiggsCombine to produce a pair of significances (cut-based σ and BDT σ).

Several thousand pairs of values from the toy pseudo-datasets are binned into 2D histogram with the 1,2, and 3- σ contours overlaid. The threshold for the contours is calculated with a 2-sample χ^2 metric. Overlaying the observed significances from above, we find it lies within

1667 the 0.7- σ contour with a p-value of 0.5, as shown in Figure 108. As described in the caption, the
 1668 observed point is “corrected” (only within the scope of this toy study) before plotting it.




Figure 101: 2016: data yields compared to prefit (top) and postfit (bottom) SM predictions in CRZ, CRW, and the cut-based signal regions (left), and CRZ with the BDT signal regions (right). Note that we plan to retrain the BDT with latest JECs/corrections for 2018 (with identical settings)




Figure 102: Comparison of correlated expected significances between cut-based and BDT fits with observed point overlaid. Because significances from toy datasets are used, we get a correction factor for expected BDT/cut-based significances with the full fit and the mean of the significances from the toy datasets. This correction factor is then applied to the observed point such that the expected significance values from the full analysis would match the mean values from the toy pseudo-datasets.

Table 36: Prefit event yields in cut-based regions for 2016.

	t \bar{t} W	t \bar{t} Z	t \bar{t} H	t \bar{t} VV	X $+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t \bar{t} t \bar{t}
CRZ	0.50 \pm 0.18	14.01 \pm 4.84	0.69 \pm 0.17	0.50 \pm 0.06	0.06 \pm 0.01	2.87 \pm 0.58	0.02 \pm 0.00	0.42 \pm 0.18	19.06 \pm 4.95	31	0.25 \pm 0.03
CRW	22.56 \pm 7.66	6.31 \pm 2.14	8.44 \pm 2.07	1.76 \pm 0.21	4.41 \pm 0.63	2.33 \pm 0.48	4.96 \pm 0.95	10.05 \pm 3.79	60.82 \pm 9.06	82	1.23 \pm 0.11
SR1	1.87 \pm 0.68	0.68 \pm 0.24	1.12 \pm 0.30	0.27 \pm 0.04	0.18 \pm 0.08	0.21 \pm 0.05	0.12 \pm 0.02	0.70 \pm 0.32	5.16 \pm 0.93	8	0.58 \pm 0.06
SR2	0.61 \pm 0.24	0.20 \pm 0.09	0.35 \pm 0.09	0.08 \pm 0.02	0.05 \pm 0.02	0.03 \pm 0.01	0.02 \pm 0.00	0.48 \pm 0.23	1.83 \pm 0.37	2	0.39 \pm 0.03
SR3	0.18 \pm 0.10	-0.11 \pm 0.06	0.17 \pm 0.06	0.05 \pm 0.01	0.00 \pm 0.01	0.01 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.17	0.53 \pm 0.25	1	0.23 \pm 0.05
SR4	0.63 \pm 0.24	0.26 \pm 0.11	0.47 \pm 0.12	0.13 \pm 0.02	-0.17 \pm 0.07	0.07 \pm 0.01	0.09 \pm 0.02	0.36 \pm 0.21	2.19 \pm 0.46	8	0.46 \pm 0.06
SR5	0.40 \pm 0.18	0.09 \pm 0.03	0.22 \pm 0.07	0.06 \pm 0.02	0.03 \pm 0.01	0.03 \pm 0.01	0.02 \pm 0.00	0.31 \pm 0.30	1.17 \pm 0.39	0	0.51 \pm 0.05
SR6	0.09 \pm 0.06	0.03 \pm 0.04	0.10 \pm 0.04	0.01 \pm 0.01	0.07 \pm 0.05	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.07	0.32 \pm 0.16	2	0.34 \pm 0.04
SR7	0.06 \pm 0.04	0.04 \pm 0.02	0.05 \pm 0.02	0.01 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.01	0.16 \pm 0.07	0	0.28 \pm 0.08
SR8	0.15 \pm 0.08	0.04 \pm 0.02	0.08 \pm 0.03	0.05 \pm 0.01	0.00 \pm 0.00	0.02 \pm 0.00	0.03 \pm 0.01	0.17 \pm 0.11	0.55 \pm 0.17	0	0.63 \pm 0.06
SR9	0.25 \pm 0.09	0.49 \pm 0.17	0.36 \pm 0.09	0.08 \pm 0.01	0.02 \pm 0.00	0.02 \pm 0.01	0.00 \pm 0.00	0.22 \pm 0.11	1.44 \pm 0.27	1	0.24 \pm 0.07
SR10	0.08 \pm 0.03	0.16 \pm 0.07	0.14 \pm 0.04	0.02 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.01	0.00 \pm 0.00	0.16 \pm 0.19	0.57 \pm 0.22	0	0.11 \pm 0.02
SR11	0.04 \pm 0.02	0.09 \pm 0.05	0.05 \pm 0.02	0.01 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.01	0.00 \pm 0.00	0.05 \pm 0.04	0.26 \pm 0.09	0	0.07 \pm 0.02
SR12	0.04 \pm 0.02	0.13 \pm 0.05	0.09 \pm 0.03	0.02 \pm 0.00	0.00 \pm 0.00	0.02 \pm 0.00	0.00 \pm 0.00	0.17 \pm 0.18	0.46 \pm 0.20	1	0.17 \pm 0.02
SR13	0.04 \pm 0.02	0.05 \pm 0.03	0.08 \pm 0.02	0.01 \pm 0.00	0.02 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.02	0.20 \pm 0.07	0	0.21 \pm 0.02
SR14	0.01 \pm 0.02	0.05 \pm 0.03	0.04 \pm 0.02	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.01	0.11 \pm 0.05	1	0.25 \pm 0.04

Table 37: Postfit event yields in cut-based regions for 2016.

	t \bar{t} W	t \bar{t} Z	t \bar{t} H	t \bar{t} VV	X $+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t \bar{t} t \bar{t}
CRZ	0.72 \pm 0.18	21.58 \pm 4.13	0.75 \pm 0.17	0.51 \pm 0.06	0.06 \pm 0.01	3.00 \pm 0.59	0.02 \pm 0.00	0.47 \pm 0.21	27.09 \pm 4.12	31	0.20 \pm 0.22
CRW	32.12 \pm 7.40	9.64 \pm 1.87	9.08 \pm 2.03	1.80 \pm 0.22	4.57 \pm 0.66	2.45 \pm 0.50	5.03 \pm 0.84	11.13 \pm 4.44	75.82 \pm 7.26	82	0.99 \pm 1.03
SR1	2.72 \pm 0.68	1.04 \pm 0.21	1.21 \pm 0.27	0.28 \pm 0.03	0.19 \pm 0.06	0.22 \pm 0.05	0.12 \pm 0.02	0.77 \pm 0.35	6.56 \pm 0.71	8	0.47 \pm 0.52
SR2	0.87 \pm 0.23	0.31 \pm 0.07	0.38 \pm 0.09	0.09 \pm 0.01	0.05 \pm 0.01	0.04 \pm 0.01	0.02 \pm 0.00	0.51 \pm 0.23	2.26 \pm 0.33	2	0.32 \pm 0.30
SR3	0.27 \pm 0.10	0.18 \pm 0.05	0.18 \pm 0.05	0.05 \pm 0.01	0.00 \pm 0.00	0.01 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.18	0.71 \pm 0.26	1	0.19 \pm 0.21
SR4	0.94 \pm 0.25	0.42 \pm 0.09	0.52 \pm 0.12	0.14 \pm 0.02	0.20 \pm 0.05	0.07 \pm 0.01	0.10 \pm 0.02	0.46 \pm 0.24	2.83 \pm 0.40	8	0.37 \pm 0.35
SR5	0.59 \pm 0.16	0.13 \pm 0.03	0.24 \pm 0.07	0.07 \pm 0.01	0.03 \pm 0.00	0.03 \pm 0.01	0.02 \pm 0.00	0.28 \pm 0.26	1.39 \pm 0.33	0	0.41 \pm 0.38
SR6	0.14 \pm 0.06	0.06 \pm 0.03	0.11 \pm 0.03	0.01 \pm 0.01	0.08 \pm 0.05	0.01 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.08	0.44 \pm 0.16	2	0.29 \pm 0.26
SR7	0.09 \pm 0.05	0.06 \pm 0.02	0.05 \pm 0.02	0.01 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.01	0.22 \pm 0.07	0	0.23 \pm 0.19
SR8	0.23 \pm 0.08	0.06 \pm 0.02	0.09 \pm 0.03	0.05 \pm 0.01	0.00 \pm 0.00	0.02 \pm 0.00	0.03 \pm 0.01	0.18 \pm 0.11	0.67 \pm 0.16	0	0.52 \pm 0.50
SR9	0.35 \pm 0.09	0.75 \pm 0.16	0.39 \pm 0.09	0.08 \pm 0.01	0.03 \pm 0.00	0.02 \pm 0.01	0.00 \pm 0.00	0.23 \pm 0.13	1.85 \pm 0.26	1	0.20 \pm 0.16
SR10	0.11 \pm 0.03	0.24 \pm 0.06	0.15 \pm 0.04	0.02 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.01	0.00 \pm 0.00	0.15 \pm 0.17	0.69 \pm 0.20	0	0.09 \pm 0.11
SR11	0.06 \pm 0.02	0.15 \pm 0.04	0.05 \pm 0.02	0.01 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.00	0.05 \pm 0.05	0.34 \pm 0.09	0	0.06 \pm 0.07
SR12	0.06 \pm 0.02	0.20 \pm 0.05	0.10 \pm 0.02	0.02 \pm 0.00	0.00 \pm 0.00	0.02 \pm 0.00	0.00 \pm 0.00	0.19 \pm 0.19	0.58 \pm 0.21	1	0.14 \pm 0.14
SR13	0.05 \pm 0.03	0.08 \pm 0.04	0.08 \pm 0.02	0.01 \pm 0.00	0.02 \pm 0.00	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.02	0.26 \pm 0.08	0	0.17 \pm 0.16
SR14	0.02 \pm 0.02	0.07 \pm 0.03	0.04 \pm 0.01	0.01 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.01	0.15 \pm 0.05	1	0.21 \pm 0.21

Table 38: Prefit event yields in BDT regions for 2016.

	$t\bar{t}W$	$t\bar{t}Z$	$t\bar{t}H$	$t\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$t\bar{t}t\bar{t}$
CRZ	0.50±0.17	14.01±4.41	0.69±0.17	0.50±0.06	0.06±0.01	2.87±0.56	0.02±0.00	0.42±0.17	19.06±4.49	31	0.25±0.03
SR1	0.08±0.04	0.02±0.01	0.01±0.01	0.00±0.00	0.02±0.01	0.01±0.01	0.07±0.01	0.05±0.05	0.27±0.08	1	0.00±0.00
SR2	0.84±0.30	0.16±0.06	0.21±0.05	0.02±0.00	0.38±0.19	0.41±0.18	0.47±0.10	0.23±0.15	2.71±0.58	3	0.00±0.01
SR3	1.75±0.65	0.52±0.19	0.47±0.15	0.06±0.01	0.75±0.21	0.13±0.05	0.62±0.13	1.62±0.69	5.93±1.09	7	0.01±0.01
SR4	2.70±0.89	0.72±0.32	0.94±0.24	0.14±0.02	0.38±0.19	0.35±0.10	0.69±0.14	1.21±0.57	7.14±1.20	12	0.04±0.01
SR5	3.19±1.07	1.00±0.45	1.20±0.31	0.21±0.03	1.25±0.29	0.36±0.20	0.75±0.15	1.88±0.73	9.84±1.60	9	0.08±0.01
SR6	3.49±1.15	1.33±0.45	1.37±0.35	0.26±0.03	0.40±0.21	0.44±0.12	0.70±0.14	1.29±0.48	9.29±1.45	17	0.11±0.05
SR7	3.47±1.14	1.37±0.48	1.53±0.37	0.32±0.04	0.46±0.15	0.29±0.06	0.54±0.11	1.68±0.70	9.65±1.50	8	0.18±0.03
SR8	3.22±1.06	1.16±0.43	1.35±0.34	0.29±0.06	0.59±0.15	0.12±0.10	0.48±0.10	1.14±0.48	8.35±1.26	7	0.23±0.02
SR9	3.78±1.40	1.48±0.51	1.87±0.48	0.51±0.06	0.37±0.22	0.31±0.07	0.46±0.09	1.95±0.92	10.72±1.91	15	0.42±0.04
SR10	3.52±1.21	1.68±0.56	2.05±0.52	0.49±0.06	0.30±0.07	0.31±0.06	0.36±0.07	1.15±0.52	9.85±1.56	20	0.74±0.07
SR11	1.94±0.70	1.21±0.41	1.31±0.36	0.42±0.06	0.40±0.09	0.30±0.08	0.22±0.04	0.69±0.39	6.48±1.07	7	0.84±0.05
SR12	1.19±0.46	0.62±0.22	0.87±0.23	0.20±0.03	0.11±0.02	0.11±0.03	0.10±0.02	1.07±0.45	4.26±0.75	12	0.91±0.06
SR13	0.70±0.28	0.29±0.11	0.52±0.18	0.14±0.03	0.00±0.03	0.08±0.02	0.07±0.01	0.54±0.32	2.33±0.48	1	0.91±0.04
SR14	0.34±0.16	0.20±0.08	0.24±0.08	0.07±0.01	0.00±0.03	0.05±0.01	0.04±0.01	0.00±0.11	0.94±0.28	3	0.86±0.07
SR15	0.14±0.08	0.09±0.04	0.13±0.05	0.04±0.01	0.06±0.06	0.02±0.01	0.02±0.00	0.17±0.12	0.68±0.21	1	0.69±0.06
SR16	0.05±0.03	0.02±0.02	0.03±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.03	0.12±0.07	1	0.39±0.05
SR17	0.01±0.01	0.01±0.01	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.06	0.02±0.07	0	0.05±0.01

Table 39: Postfit event yields in BDT regions for 2016.

	$t\bar{t}W$	$t\bar{t}Z$	$t\bar{t}H$	$t\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$t\bar{t}t\bar{t}$
CRZ	0.66±0.19	21.93±4.77	0.74±0.17	0.51±0.06	0.06±0.01	2.98±0.55	0.02±0.00	0.47±0.18	27.37±4.71	31	0.38±0.24
SR1	0.11±0.04	0.03±0.01	0.01±0.01	0.00±0.00	0.02±0.01	0.01±0.01	0.07±0.01	0.06±0.06	0.33±0.09	1	0.00±0.00
SR2	1.10±0.33	0.24±0.08	0.22±0.05	0.02±0.00	0.39±0.18	0.41±0.18	0.48±0.09	0.27±0.17	3.14±0.54	3	0.00±0.01
SR3	2.30±0.71	0.80±0.21	0.51±0.14	0.06±0.01	0.77±0.17	0.13±0.05	0.63±0.11	1.82±0.80	7.02±1.04	7	0.02±0.02
SR4	3.54±0.97	1.11±0.40	1.01±0.22	0.15±0.02	0.43±0.18	0.37±0.09	0.70±0.13	1.44±0.71	8.75±1.15	12	0.06±0.04
SR5	4.18±1.18	1.56±0.57	1.28±0.28	0.21±0.03	1.24±0.26	0.36±0.16	0.76±0.14	2.06±0.77	11.66±1.41	9	0.12±0.08
SR6	4.64±1.26	2.06±0.48	1.47±0.32	0.27±0.04	0.46±0.19	0.48±0.11	0.71±0.13	1.50±0.62	11.58±1.24	17	0.17±0.16
SR7	4.55±1.22	2.13±0.52	1.63±0.35	0.32±0.04	0.46±0.15	0.29±0.06	0.54±0.10	1.83±0.76	11.75±1.23	8	0.27±0.17
SR8	4.23±1.12	1.79±0.49	1.43±0.31	0.30±0.05	0.57±0.13	0.13±0.08	0.48±0.09	1.26±0.57	10.20±1.11	7	0.36±0.21
SR9	5.00±1.48	2.30±0.56	2.00±0.45	0.52±0.06	0.37±0.17	0.32±0.07	0.46±0.08	2.24±1.07	13.22±1.68	15	0.66±0.39
SR10	4.69±1.32	2.64±0.62	2.18±0.49	0.49±0.06	0.31±0.06	0.32±0.06	0.37±0.07	1.36±0.64	12.37±1.37	20	1.14±0.67
SR11	2.59±0.74	1.88±0.46	1.40±0.34	0.42±0.06	0.40±0.09	0.31±0.07	0.22±0.04	0.76±0.43	7.98±0.96	7	1.30±0.76
SR12	1.62±0.52	1.00±0.27	0.93±0.22	0.21±0.03	0.11±0.02	0.12±0.03	0.10±0.02	1.30±0.65	5.39±0.84	12	1.41±0.80
SR13	0.94±0.29	0.45±0.13	0.56±0.17	0.14±0.02	0.00±0.02	0.08±0.01	0.07±0.01	0.54±0.33	2.78±0.48	1	1.41±0.83
SR14	0.45±0.18	0.33±0.10	0.26±0.08	0.08±0.01	0.00±0.02	0.05±0.01	0.04±0.01	0.00±0.11	1.22±0.30	3	1.35±0.76
SR15	0.19±0.09	0.15±0.06	0.14±0.05	0.05±0.01	0.06±0.05	0.02±0.00	0.02±0.00	0.18±0.12	0.82±0.22	1	1.09±0.60
SR16	0.07±0.04	0.04±0.03	0.03±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.03	0.16±0.08	1	0.62±0.36
SR17	0.01±0.01	0.01±0.02	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.08	0.02±0.08	0	0.07±0.06

1669 **F.3 Nuisances**

1670 Four sets of nuisance pull values, expected and observed for cut-based and BDT analyses, are
 1671 tabulated in Table 40 (expected cut-based analysis), Table 41 (observed cut-based analysis),
 1672 Table 42 (expected BDT analysis), Table 43 (observed BDT analysis).

Table 40: Expected nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	
TTH	+0.14, 0.99	+0.00, 0.98	-0.08
TTVV	+0.02, 1.00	+0.00, 0.99	-0.01
TTWSF	+0.16, 0.76	+0.00, 0.78	-0.14
TTZSF	+0.06, 0.69	+0.00, 0.69	-0.05
XG	+0.01, 0.99	+0.00, 0.99	-0.00
alphas	+0.00, 0.99	-0.00, 0.99	+0.00
bb	+0.25, 0.96	+0.00, 0.99	-0.13
btaghf	+0.07, 0.99	+0.00, 0.99	-0.07
btaglf	+0.05, 0.99	+0.00, 0.99	-0.03
fakes	+0.09, 0.97	+0.00, 0.97	-0.07
fakes.EWK	+0.07, 0.96	+0.00, 0.96	-0.05
flips	-0.01, 0.99	+0.00, 0.99	+0.00
fsrvar	+0.00, 0.99	-0.00, 0.99	+0.04
isr	-0.14, 0.98	-0.00, 0.99	+0.09
isrvar	+0.00, 0.99	+0.00, 0.99	-0.03
jer	-0.09, 0.97	-0.00, 0.99	+0.05
jes	+0.10, 0.91	+0.00, 0.99	-0.10
lep	+0.02, 0.99	+0.00, 0.99	-0.03
lumi	+0.04, 0.99	+0.00, 0.99	-0.02
pdf	+0.09, 0.96	-0.00, 0.99	-0.05
prefire	-0.01, 0.99	-0.00, 0.99	+0.01
pu	+0.00, 0.99	-0.00, 0.99	+0.02
rares	-0.00, 0.99	+0.00, 0.99	-0.00
scale	-0.11, 1.00	+0.00, 0.99	+0.04
trig	+0.03, 0.99	+0.00, 0.99	-0.04

Table 41: Observed nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.32, 1.02	+0.23, 1.01	-0.08
TTVV	+0.04, 1.00	+0.03, 1.00	-0.01
TTWSF	+1.09, 0.73	+0.97, 0.74	-0.17
TTZSF	+1.23, 0.61	+1.18, 0.62	-0.06
XG	+0.09, 1.00	+0.07, 1.00	-0.01
alphas	-0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.29, 0.97	+0.17, 0.98	-0.11
btaghf	+0.33, 0.99	+0.27, 0.99	-0.07
btaglf	+0.14, 0.99	+0.09, 0.99	-0.04
fakes	+0.25, 1.02	+0.21, 1.01	-0.04
fakes.EWK	+0.17, 0.98	+0.14, 0.95	-0.02
flips	+0.07, 1.00	+0.08, 1.00	+0.01
fsrvar	-0.00, 0.99	-0.03, 1.15	+0.00
isr	-0.24, 0.97	-0.15, 0.98	+0.09
isrvar	-0.00, 0.99	+0.06, 1.01	+0.02
jer	-0.05, 0.52	-0.01, 0.50	+0.04
jes	+0.13, 0.66	+0.07, 0.62	-0.09
lep	+0.16, 0.99	+0.14, 0.99	-0.03
lumi	+0.25, 0.99	+0.22, 0.99	-0.03
pdf	+0.23, 0.94	+0.10, 0.97	-0.12
prefire	-0.04, 0.99	-0.04, 0.99	+0.01
pu	-0.11, 0.98	-0.10, 0.99	+0.01
rares	+0.11, 1.00	+0.12, 1.00	+0.00
scale	+0.01, 0.96	+0.07, 0.95	+0.04
trig	+0.21, 0.99	+0.19, 0.99	-0.04

Table 42: Expected nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{in}, \sigma_{out}/\sigma_{in}$	$\Delta x/\sigma_{in}, \sigma_{out}/\sigma_{in}$	
TTH	+0.12, 0.99	-0.00, 0.98	-0.08
TTVV	+0.02, 1.00	+0.00, 0.99	-0.01
TTWSF	+0.20, 0.76	-0.00, 0.78	-0.16
TTZSF	+0.04, 0.69	+0.00, 0.69	-0.05
XG	-0.00, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	-0.00, 0.99	+0.00
bb	+0.33, 0.95	+0.00, 0.99	-0.16
btaghf	+0.07, 0.99	+0.00, 0.99	-0.07
btaglf	+0.05, 0.99	+0.00, 0.99	-0.03
fakes	+0.07, 0.97	-0.00, 0.96	-0.06
fakes.EWK	+0.06, 0.94	+0.00, 0.95	-0.06
flips	-0.00, 0.99	-0.00, 0.99	+0.00
fsrvar	+0.00, 0.99	-0.00, 0.99	+0.02
isr	-0.17, 0.97	-0.00, 0.99	+0.10
isrvar	+0.00, 0.99	-0.00, 0.99	-0.03
jer	-0.07, 1.07	+0.00, 0.99	+0.04
jes	+0.11, 0.98	+0.00, 0.97	-0.09
lep	+0.02, 0.99	+0.00, 0.99	-0.03
lumi	+0.04, 0.99	+0.00, 0.99	-0.02
pdf	+0.09, 0.97	-0.00, 0.99	-0.05
prefire	-0.01, 0.99	-0.00, 0.99	+0.01
pu	+0.04, 0.99	-0.00, 0.99	-0.01
rares	+0.00, 0.99	+0.00, 0.99	-0.00
scale	-0.14, 1.01	+0.00, 0.99	+0.04
trig	+0.03, 0.99	+0.00, 0.99	-0.05

Table 43: Observed nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	
TTH	+0.33, 1.02	+0.21, 1.01	-0.07
TTVV	+0.03, 1.00	+0.02, 1.00	-0.01
TTWSF	+1.05, 0.74	+0.74, 0.78	-0.20
TTZSF	+1.28, 0.62	+1.24, 0.62	-0.05
XG	-0.02, 0.99	-0.01, 0.99	+0.00
alphas	+0.00, 0.99	+0.00, 1.00	+0.00
bb	+0.63, 0.95	+0.18, 0.98	-0.16
btaghf	+0.25, 0.99	+0.16, 0.99	-0.08
btaglf	+0.12, 0.99	+0.04, 0.99	-0.04
fakes	+0.30, 1.01	+0.26, 1.00	-0.05
fakes.EWK	+0.24, 0.99	+0.21, 0.96	-0.04
flips	+0.05, 1.00	+0.06, 1.00	+0.01
fsrvar	+0.00, 0.99	-0.04, 1.01	+0.00
isr	-0.44, 0.97	-0.17, 0.98	+0.12
isrvar	+0.00, 0.99	+0.01, 1.01	-0.03
jer	-0.15, 0.93	-0.03, 0.83	+0.06
jes	+0.08, 0.78	-0.02, 0.75	-0.09
lep	+0.15, 0.99	+0.13, 0.99	-0.03
lumi	+0.23, 0.99	+0.18, 0.99	-0.02
pdf	+0.23, 0.95	+0.08, 0.97	-0.08
prefire	-0.05, 0.99	-0.04, 0.99	+0.02
pu	-0.18, 1.04	-0.23, 1.03	-0.00
rares	+0.10, 1.00	+0.11, 1.00	+0.00
scale	+0.00, 0.96	+0.13, 0.95	+0.02
trig	+0.19, 0.99	+0.16, 0.99	-0.05

F.4 Impacts

1673

1674 The leading 30 nuisance impacts for four sets of impacts, expected and observed for cut-based
 1675 and BDT analyses, are shown in Figure 103 (expected cut-based analysis), Figure 104 (observed
 1676 cut-based analysis), Figure 105 (expected BDT analysis), Figure 106 (observed BDT analysis).
 1677 The most constrained nuisances correspond to normalization parameters for ttW and ttZ; as we
 1678 would expect from the control regions, “TTWSF” and “TTZSF”, are moved by approximately
 1σ with respect to the input nuisance sizes.




Figure 103: Expected nuisance impacts for the cut-based analysis.

1679




Figure 104: Observed nuisance impacts for the cut-based analysis.




Figure 105: Expected nuisance impacts for the BDT-based analysis.




Figure 106: Observed nuisance impacts for the BDT-based analysis.

1680 G Unblinding of 2017 dataset

1681 Following the 2016 unblinding, the analysis was unblinded using the 2017 dataset. The yields
 1682 and results are shown in Section G.1 with consistency checks of cut-based and BDT results
 1683 in Section G.2. We include the results of various statistical checks, including nuisance pulls
 1684 (Section G.3), nuisance impacts (Section G.4), and goodness of fits (Section G.5). Additionally,
 1685 Section G.6 shows information about the combination of unblinded 2017 with unblinded 2016
 1686 data.

1687 G.1 Yields and results

1688 Plots for the unblinded prefit and postfit event yields for the 2017 data, with a total luminosity
 1689 of 41.5 fb^{-1} , are shown for both the cut-based and BDT based analysis in Figure 107. Numerical
 1690 yields are also tabulated in Table 44 (prefit cut-based analysis), Table 45 (postfit cut-based
 1691 analysis), Table 46 (prefit BDT analysis), Table 47 (postfit BDT analysis).

1692 A binned likelihood fit is performed over signal regions using the Higgs Combine tool [22];
 1693 exclusion limits at 95% CL are calculated with the Asymptotic CLs method.

1694 With the cut-based analysis, an observed (expected) upper limit on the production cross section
 1695 of 30.18 fb ($17.77^{+9.63}_{-5.93} \text{ fb}$), assuming the signal process does not exist. The observed (expected)
 1696 significance is 0.853 (1.543) standard deviations, corresponding to a measured signal strength
 1697 parameter of $0.703^{+0.985}_{-0.703}$.

1698 With the BDT analysis, an observed (expected) upper limit on the production cross section of
 1699 29.85 fb ($15.94^{+8.71}_{-5.36} \text{ fb}$), assuming the signal process does not exist. The observed (expected)
 1700 significance is 1.105 (1.756) standard deviations, corresponding to a measured signal strength
 1701 parameter of $0.806^{+0.924}_{-0.736}$.

1702 G.2 Cut-based and BDT consistency checks

1703 In 2016, cut-based SR4 had a high data yield with respect to the SM prediction. In 2017, this
 1704 region is in agreement. The situation is reversed for cut-based SR3, for example.

1705 For the 2016 unblinding, we quantified the mildness of the disagreement between cut-based
 1706 and BDT significances (both had differing directions for observed significance with respect to
 1707 their expected significance), using fits of toy pseudodatasets correlated between cut-based and
 1708 BDT regions.

1709 Although 2017 shows much better agreement between the two fits, we repeat the same procedure
 1710 for 2017. The observed values lie within a $0.3\text{-}\sigma$ contour, with a p-value of nearly 0.8, as
 1711 shown in Figure 108.




Figure 107: 2017: data yields compared to prefit (top) and postfit (bottom) SM predictions in CRZ, CRW, and the cut-based signal regions (left), and CRZ with the BDT signal regions (right). Note that we plan to retrain the BDT with latest JECs/corrections for 2018 (with identical settings)




Figure 108: Comparison of correlated expected significances between cut-based and BDT fits with observed point overlaid. Because significances from toy datasets are used, we get a correction factor for expected BDT/cut-based significances with the full fit and the mean of the significances from the toy datasets. This correction factor is then applied to the observed point such that the expected significance values from the full analysis would match the mean values from the toy pseudo-datasets.

Table 44: Prefit event yields in cut-based regions for 2017.

	t̄W	t̄Z	t̄H	t̄VV	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t̄t̄
CRZ	0.53±0.19	16.56±5.59	0.77±0.21	0.58±0.07	0.33±0.13	3.69±0.78	0.00±0.00	0.79±0.42	23.26±5.81	22	0.32±0.02
CRW	27.27±9.20	7.44±2.50	9.52±2.48	1.98±0.25	3.60±0.50	2.01±0.41	4.98±0.95	15.64±8.38	72.45±12.99	92	1.43±0.10
SR1	1.92±0.66	0.68±0.24	1.31±0.35	0.33±0.04	0.15±0.04	0.16±0.05	0.14±0.03	1.36±1.04	6.05±1.33	10	0.73±0.06
SR2	0.62±0.25	0.08±0.07	0.43±0.12	0.11±0.01	0.01±0.01	0.03±0.01	0.02±0.00	0.48±0.37	1.79±0.47	3	0.38±0.03
SR3	0.13±0.07	0.00±0.02	0.17±0.06	0.03±0.01	0.00±0.00	0.02±0.00	0.01±0.00	0.14±0.16	0.48±0.21	0	0.28±0.05
SR4	0.62±0.24	0.25±0.12	0.43±0.13	0.14±0.02	0.06±0.01	0.05±0.01	0.08±0.02	0.60±0.50	2.23±0.65	2	0.56±0.07
SR5	0.35±0.18	0.10±0.04	0.23±0.07	0.07±0.01	0.01±0.01	0.08±0.02	0.01±0.00	0.08±0.09	0.94±0.23	2	0.56±0.04
SR6	0.13±0.06	0.01±0.03	0.10±0.03	0.03±0.00	0.01±0.00	0.02±0.00	0.01±0.00	0.05±0.06	0.36±0.11	3	0.42±0.04
SR7	0.05±0.03	0.01±0.01	0.04±0.02	0.01±0.00	0.01±0.00	0.01±0.00	0.01±0.00	0.00±0.04	0.14±0.07	0	0.30±0.03
SR8	0.06±0.04	0.04±0.02	0.08±0.03	0.05±0.01	0.02±0.01	0.02±0.00	0.01±0.00	0.28±0.27	0.57±0.29	0	0.73±0.08
SR9	0.32±0.13	0.48±0.16	0.43±0.11	0.09±0.01	0.03±0.00	0.04±0.01	0.00±0.00	0.08±0.07	1.47±0.26	2	0.23±0.02
SR10	0.00±0.03	0.16±0.07	0.15±0.05	0.03±0.00	0.01±0.01	0.01±0.00	0.00±0.00	0.19±0.21	0.55±0.24	3	0.15±0.01
SR11	0.02±0.01	0.06±0.03	0.05±0.02	0.01±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.03	0.14±0.06	0	0.08±0.01
SR12	0.02±0.01	0.12±0.05	0.08±0.03	0.03±0.00	0.01±0.00	0.01±0.00	0.00±0.00	0.26±0.30	0.53±0.30	1	0.23±0.03
SR13	0.05±0.03	0.07±0.03	0.06±0.02	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.07	0.21±0.09	0	0.25±0.02
SR14	0.06±0.03	0.05±0.03	0.04±0.02	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.07	0.16±0.09	0	0.29±0.03

Table 45: Postfit event yields in cut-based regions for 2017.

	t̄W	t̄Z	t̄H	t̄VV	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t̄t̄
CRZ	0.65±0.20	16.04±3.53	0.81±0.20	0.59±0.08	0.34±0.13	3.70±0.77	0.00±0.00	0.98±0.72	23.11±3.77	22	0.22±0.23
CRW	32.98±9.73	7.23±1.61	9.98±2.40	2.00±0.26	3.63±0.51	2.01±0.41	5.00±0.84	25.44±12.71	88.27±10.26	92	1.01±1.04
SR1	2.36±0.74	0.67±0.15	1.39±0.33	0.34±0.04	0.16±0.04	0.17±0.04	0.14±0.02	2.35±1.56	7.56±1.37	10	0.51±0.53
SR2	0.76±0.29	0.08±0.06	0.46±0.11	0.11±0.01	0.01±0.01	0.03±0.01	0.02±0.00	0.85±0.59	2.33±0.57	3	0.27±0.29
SR3	0.15±0.07	0.00±0.01	0.18±0.05	0.03±0.01	0.00±0.00	0.02±0.00	0.01±0.00	0.13±0.27	0.51±0.30	0	0.20±0.19
SR4	0.75±0.25	0.24±0.07	0.45±0.12	0.14±0.02	0.06±0.01	0.05±0.01	0.08±0.01	0.91±0.63	2.67±0.64	2	0.39±0.39
SR5	0.43±0.18	0.10±0.03	0.24±0.07	0.07±0.01	0.01±0.00	0.08±0.02	0.01±0.00	0.14±0.12	1.09±0.22	2	0.39±0.41
SR6	0.17±0.07	0.01±0.03	0.11±0.03	0.03±0.00	0.01±0.00	0.02±0.00	0.01±0.00	0.10±0.11	0.46±0.13	3	0.29±0.28
SR7	0.07±0.03	0.01±0.01	0.04±0.02	0.01±0.00	0.01±0.00	0.01±0.00	0.01±0.00	0.00±0.08	0.15±0.10	0	0.21±0.22
SR8	0.07±0.05	0.04±0.01	0.08±0.03	0.05±0.01	0.02±0.01	0.02±0.00	0.01±0.00	0.23±0.45	0.52±0.46	0	0.52±0.52
SR9	0.40±0.15	0.46±0.11	0.45±0.11	0.09±0.01	0.03±0.01	0.04±0.01	0.00±0.00	0.15±0.12	1.62±0.22	2	0.16±0.16
SR10	0.00±0.03	0.16±0.05	0.16±0.04	0.03±0.00	0.01±0.00	0.01±0.00	0.00±0.00	0.44±0.35	0.81±0.34	3	0.11±0.11
SR11	0.03±0.01	0.06±0.02	0.05±0.01	0.01±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.01±0.06	0.16±0.07	0	0.05±0.06
SR12	0.02±0.01	0.12±0.03	0.08±0.03	0.03±0.00	0.01±0.00	0.01±0.00	0.00±0.00	0.37±0.41	0.63±0.41	1	0.16±0.16
SR13	0.07±0.03	0.07±0.02	0.06±0.02	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.11	0.22±0.12	0	0.17±0.17
SR14	0.07±0.03	0.05±0.02	0.04±0.02	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.10	0.18±0.11	0	0.21±0.21

Table 46: Prefit event yields in BDT regions for 2017.

	t̄W	t̄Z	t̄H	t̄VV	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t̄t̄
CRZ	0.53±0.21	16.56±5.54	0.77±0.16	0.58±0.08	0.33±0.15	3.69±0.76	0.00±0.00	0.79±0.44	23.26±5.72	22	0.32±0.02
SR1	0.09±0.05	0.02±0.01	0.02±0.01	0.00±0.00	0.01±0.01	0.02±0.03	0.05±0.01	0.00±0.09	0.21±0.13	0	0.00±0.00
SR2	1.07±0.46	0.26±0.15	0.20±0.06	0.03±0.00	0.26±0.12	0.10±0.06	0.40±0.08	0.42±0.53	2.75±0.80	4	0.00±0.00
SR3	2.35±1.01	0.65±0.31	0.62±0.14	0.08±0.02	0.70±0.24	0.44±0.17	0.63±0.12	1.71±1.01	7.18±1.52	7	0.01±0.01
SR4	3.52±1.38	1.05±0.53	0.91±0.21	0.14±0.03	0.36±0.20	0.30±0.14	0.73±0.14	2.70±1.31	9.71±2.10	16	0.04±0.01
SR5	4.02±1.69	1.34±0.60	1.28±0.29	0.22±0.03	0.61±0.19	0.33±0.17	0.70±0.13	2.93±1.88	11.44±2.67	5	0.08±0.01
SR6	4.33±1.73	1.45±0.64	1.58±0.34	0.31±0.04	0.85±0.21	0.30±0.11	0.67±0.13	2.05±1.52	11.54±2.46	14	0.14±0.02
SR7	3.97±1.63	1.44±0.65	1.63±0.35	0.32±0.05	0.54±0.08	0.42±0.15	0.60±0.11	1.63±1.41	10.56±2.17	14	0.17±0.03
SR8	3.42±1.37	1.41±0.57	1.63±0.36	0.41±0.05	0.28±0.09	0.21±0.05	0.50±0.10	2.97±1.60	10.85±2.13	17	0.26±0.02
SR9	4.33±1.74	1.65±0.73	2.19±0.46	0.55±0.06	0.40±0.09	0.26±0.06	0.46±0.09	1.88±1.83	11.73±2.57	14	0.49±0.06
SR10	3.92±1.59	1.78±0.67	2.22±0.49	0.53±0.07	0.23±0.10	0.31±0.07	0.40±0.08	2.30±1.48	11.70±2.39	15	0.83±0.05
SR11	2.18±1.00	1.01±0.45	1.45±0.33	0.43±0.05	0.30±0.08	0.16±0.04	0.21±0.04	1.87±1.45	7.62±1.82	16	1.00±0.04
SR12	1.21±0.55	0.68±0.25	0.91±0.23	0.30±0.04	0.01±0.04	0.08±0.02	0.13±0.02	0.89±0.60	4.21±0.95	6	1.08±0.05
SR13	0.56±0.30	0.30±0.14	0.49±0.13	0.15±0.02	0.02±0.01	0.10±0.02	0.05±0.01	0.16±0.28	1.84±0.51	2	1.03±0.05
SR14	0.36±0.21	0.06±0.04	0.27±0.08	0.10±0.02	0.03±0.01	0.07±0.01	0.02±0.00	0.19±0.17	1.11±0.32	3	1.05±0.05
SR15	0.12±0.07	0.07±0.04	0.12±0.04	0.05±0.01	0.00±0.00	0.02±0.01	0.02±0.00	0.04±0.06	0.45±0.13	1	0.76±0.06
SR16	0.02±0.02	0.02±0.01	0.03±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.10±0.07	0.16±0.07	0	0.49±0.03
SR17	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.25±0.32	0.26±0.32	0	0.05±0.01

Table 47: Postfit event yields in BDT regions for 2017.

	t̄W	t̄Z	t̄H	t̄VV	X+γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	t̄t̄
CRZ	0.61±0.20	15.95±3.76	0.80±0.18	0.59±0.07	0.32±0.12	3.68±0.77	0.00±0.00	1.00±0.74	22.96±4.00	22	0.26±0.22
SR1	0.11±0.04	0.02±0.01	0.02±0.01	0.00±0.00	0.01±0.01	0.02±0.03	0.05±0.01	0.00±0.15	0.21±0.17	0	0.00±0.00
SR2	1.22±0.43	0.25±0.10	0.21±0.07	0.03±0.00	0.25±0.11	0.10±0.07	0.40±0.07	0.91±0.78	3.37±0.81	4	0.00±0.00
SR3	2.66±0.95	0.61±0.21	0.64±0.17	0.08±0.02	0.70±0.24	0.42±0.15	0.63±0.11	2.38±1.37	8.13±1.43	7	0.01±0.02
SR4	4.00±1.31	1.00±0.36	0.95±0.26	0.14±0.03	0.38±0.18	0.30±0.16	0.73±0.13	4.35±2.37	11.85±2.25	16	0.03±0.03
SR5	4.56±1.57	1.28±0.38	1.32±0.35	0.22±0.03	0.58±0.15	0.32±0.20	0.70±0.12	3.78±1.98	12.76±2.24	5	0.07±0.07
SR6	4.92±1.61	1.38±0.39	1.63±0.40	0.31±0.04	0.87±0.18	0.30±0.13	0.67±0.11	3.44±2.08	13.52±2.00	14	0.11±0.11
SR7	4.54±1.52	1.37±0.39	1.70±0.41	0.32±0.05	0.54±0.09	0.43±0.17	0.60±0.10	3.07±2.10	12.56±1.85	14	0.14±0.15
SR8	3.90±1.28	1.36±0.34	1.70±0.43	0.41±0.05	0.29±0.10	0.21±0.05	0.51±0.09	4.86±2.79	13.24±2.49	17	0.21±0.19
SR9	4.95±1.59	1.59±0.45	2.29±0.53	0.56±0.06	0.40±0.10	0.26±0.08	0.46±0.08	3.58±2.60	14.09±2.31	14	0.40±0.32
SR10	4.51±1.47	1.72±0.42	2.32±0.57	0.54±0.07	0.23±0.10	0.31±0.08	0.40±0.07	3.71±2.04	13.74±1.97	15	0.67±0.63
SR11	2.54±0.93	0.98±0.29	1.52±0.38	0.43±0.05	0.31±0.07	0.17±0.04	0.21±0.04	3.89±2.56	10.06±2.31	16	0.81±0.72
SR12	1.41±0.48	0.65±0.16	0.96±0.25	0.30±0.03	0.01±0.03	0.08±0.02	0.13±0.02	1.35±0.84	4.89±0.92	6	0.87±0.76
SR13	0.64±0.26	0.29±0.09	0.52±0.14	0.15±0.02	0.02±0.01	0.10±0.02	0.05±0.01	0.41±0.45	2.19±0.51	2	0.83±0.73
SR14	0.42±0.16	0.07±0.04	0.28±0.09	0.11±0.02	0.03±0.01	0.07±0.01	0.02±0.00	0.30±0.30	1.29±0.32	3	0.85±0.74
SR15	0.14±0.06	0.07±0.03	0.13±0.05	0.05±0.01	0.00±0.00	0.02±0.01	0.02±0.00	0.09±0.11	0.52±0.14	1	0.61±0.53
SR16	0.03±0.02	0.02±0.01	0.03±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.10±0.10	0.10±0.10	0	0.40±0.35
SR17	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.14±0.52	0.15±0.52	0	0.04±0.03

1712 **G.3 Nuisances**

1713 Four sets of nuisance pull values, expected and observed for cut-based and BDT analyses, are
 1714 tabulated in Table 48 (expected cut-based analysis), Table 49 (observed cut-based analysis),
 1715 Table 50 (expected BDT analysis), Table 51 (observed BDT analysis).

Table 48: Expected nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{in}, \sigma_{out} / \sigma_{in}$	$\Delta x / \sigma_{in}, \sigma_{out} / \sigma_{in}$	
TTH	+0.15, 0.99	-0.00, 0.98	-0.08
TTVV	+0.02, 1.00	-0.00, 0.99	-0.01
TTWSF	+0.13, 0.80	-0.00, 0.81	-0.08
TTZSF	+0.05, 0.67	-0.00, 0.67	-0.03
XG	-0.00, 0.99	-0.00, 0.99	+0.00
alphas	+0.00, 0.99	-0.00, 0.99	+0.00
bb	+0.25, 0.96	-0.00, 0.99	-0.12
btaghf	+0.10, 0.98	-0.00, 0.98	-0.08
btaglf	+0.05, 0.99	-0.00, 0.99	-0.03
fakes	+0.14, 0.95	-0.00, 0.96	-0.07
fakes.EWK	+0.10, 0.85	-0.00, 0.85	-0.08
flips	-0.01, 0.99	-0.00, 0.99	+0.00
fsrvar	+0.00, 0.99	-0.00, 0.99	+0.01
isr	+0.10, 0.98	+0.00, 0.99	-0.05
isrvar	+0.00, 0.99	+0.00, 0.99	-0.02
jer	-0.08, 0.95	+0.00, 0.99	+0.05
jes	+0.10, 0.92	+0.00, 0.99	-0.10
lep	+0.03, 0.99	-0.00, 0.99	-0.04
lumi	+0.03, 0.99	-0.00, 0.99	-0.02
pdf	+0.01, 1.01	+0.00, 0.99	+0.01
prefire	-0.02, 0.99	+0.00, 0.99	+0.02
pu	-0.02, 0.99	+0.00, 0.99	+0.01
rares	+0.01, 0.99	-0.00, 0.99	-0.00
scale	-0.05, 0.99	+0.00, 0.99	-0.00
trig	+0.03, 0.99	-0.00, 0.99	-0.04

Table 49: Observed nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{in}, \sigma_{out} / \sigma_{in}$	$\Delta x / \sigma_{in}, \sigma_{out} / \sigma_{in}$	
TTH	+0.26, 1.02	+0.20, 1.01	-0.07
TTVV	+0.03, 1.00	+0.02, 1.00	-0.01
TTWSF	+0.59, 0.96	+0.56, 0.93	-0.03
TTZSF	-0.08, 0.71	-0.10, 0.69	-0.03
XG	+0.01, 0.99	+0.01, 1.00	+0.00
alphas	-0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.05, 0.98	-0.04, 0.99	-0.09
btaghf	+0.13, 0.98	+0.03, 0.98	-0.11
btaglf	+0.00, 0.99	-0.02, 0.99	-0.03
fakes	+0.70, 0.97	+0.58, 0.98	-0.13
fakes.EWK	+1.00, 0.95	+0.84, 1.01	-0.17
flips	+0.02, 0.99	+0.03, 1.00	+0.01
fsrvar	-0.00, 0.99	+0.02, 1.01	+0.03
isr	+0.21, 0.99	+0.13, 0.99	-0.09
isrvar	-0.00, 0.99	-0.01, 0.99	-0.02
jer	-0.15, 1.06	-0.05, 0.82	+0.08
jes	+0.17, 0.80	+0.13, 0.81	-0.07
lep	+0.05, 0.99	+0.04, 0.99	-0.02
lumi	+0.07, 0.99	+0.06, 0.99	-0.01
pdf	-0.02, 1.05	-0.02, 1.05	-0.00
prefire	-0.03, 0.99	-0.02, 0.99	+0.02
pu	+0.02, 1.01	-0.01, 1.01	-0.03
rares	-0.01, 0.99	-0.01, 0.99	-0.01
scale	-0.11, 1.05	-0.10, 1.07	-0.01
trig	+0.07, 0.99	+0.06, 0.99	-0.03

Table 50: Expected nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	
TTH	+0.17, 0.99	+0.00, 0.98	-0.08
TTVV	+0.03, 1.00	+0.00, 0.99	-0.01
TTWSF	+0.16, 0.80	+0.00, 0.82	-0.09
TTZSF	+0.04, 0.67	+0.00, 0.67	-0.03
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.36, 0.95	+0.00, 0.99	-0.14
btaghf	+0.11, 0.98	+0.00, 0.98	-0.09
btaglf	+0.05, 0.99	+0.00, 0.99	-0.03
fakes	+0.05, 0.94	+0.00, 0.94	-0.04
fakes.EWK	+0.12, 0.86	+0.00, 0.82	-0.10
flips	-0.01, 0.99	+0.00, 0.99	+0.00
fsrvar	+0.00, 0.99	+0.00, 0.99	+0.01
isr	+0.08, 0.99	+0.00, 0.99	-0.04
isrvar	+0.00, 0.99	+0.00, 0.99	-0.01
jer	-0.10, 1.32	+0.00, 0.99	+0.04
jes	+0.12, 0.97	+0.00, 0.97	-0.08
lep	+0.03, 0.99	+0.00, 0.99	-0.04
lumi	+0.04, 0.99	+0.00, 0.99	-0.02
pdf	+0.03, 1.01	+0.00, 0.99	-0.01
prefire	-0.02, 0.99	+0.00, 0.99	+0.02
pu	+0.05, 0.99	+0.00, 0.99	-0.02
rares	+0.02, 0.99	+0.00, 0.99	-0.00
scale	-0.02, 0.97	+0.00, 0.99	-0.01
trig	+0.04, 0.99	+0.00, 0.99	-0.05

Table 51: Observed nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{in}, \sigma_{out}/\sigma_{in}$	$\Delta x/\sigma_{in}, \sigma_{out}/\sigma_{in}$	
TTH	+0.25, 1.02	+0.17, 1.01	-0.07
TTVV	+0.04, 1.00	+0.02, 1.00	-0.01
TTWSF	+0.42, 0.98	+0.38, 0.95	-0.03
TTZSF	-0.11, 0.69	-0.13, 0.69	-0.02
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.23, 0.97	+0.04, 0.99	-0.14
btaghf	+0.10, 0.98	+0.03, 0.98	-0.09
btaglf	+0.06, 0.99	+0.02, 0.99	-0.04
fakes	+0.75, 0.97	+0.64, 0.97	-0.09
fakes.EWK	+1.01, 0.88	+0.83, 0.91	-0.17
flips	+0.00, 0.99	+0.01, 0.99	+0.01
fsrvar	+0.00, 0.99	-0.01, 0.99	-0.00
isr	+0.14, 0.99	+0.07, 0.99	-0.05
isrvar	+0.00, 0.99	-0.01, 0.99	-0.02
jer	-0.09, 1.15	-0.05, 1.06	+0.03
jes	+0.24, 1.01	+0.17, 1.10	-0.09
lep	+0.03, 0.99	+0.02, 0.99	-0.02
lumi	+0.05, 0.99	+0.04, 0.99	-0.01
pdf	+0.01, 1.03	-0.03, 1.03	-0.01
prefire	-0.01, 0.99	-0.01, 0.99	+0.02
pu	+0.08, 1.00	+0.04, 1.00	-0.03
rares	-0.02, 0.99	-0.03, 0.99	-0.01
scale	+0.03, 0.93	+0.04, 0.94	-0.01
trig	+0.05, 0.99	+0.04, 0.99	-0.03

1716 **G.4 Impacts**

1717 The leading 30 nuisance impacts for four sets of impacts, expected and observed for cut-based
 1718 and BDT analyses, are shown in Figure 109 (expected cut-based analysis), Figure 110 (observed
 1719 cut-based analysis), Figure 111 (expected BDT analysis), Figure 112 (observed BDT analysis).
 1720 Note that statistical nuisances tend to show up in the impact plots because the binning for the
 1721 cut-based and BDT analyses were optimized for the full luminosity (3.3 times what is shown
 1722 here for 2017). As in 2016, the most constrained nuisances correspond to normalization pa-
 1723 rameters for ttW and ttZ, as we would expect from the control regions. “TTWSF” is moved
 1724 by approximately 1σ with respect to the input nuisance sizes. Because the ttZ control region
 1725 has better data, simulation agreement in 2017 compared to 2016, the “TTZSF” nuisance is not
 pulled up here.




Figure 109: Expected nuisance impacts for the cut-based analysis.




Figure 110: Observed nuisance impacts for the cut-based analysis.




Figure 111: Expected nuisance impacts for the BDT-based analysis.




Figure 112: Observed nuisance impacts for the BDT-based analysis.

1727 **G.5 Goodness of fits**

1728 The goodness of fit distributions (using the saturated test statistic with the signal+background fit to data and asimov toys) for the cut-based and BDT analyses are shown in Figure 113.




Figure 113: GOF test for the cut-based analysis (left) and BDT-based analysis (right)

1729

1730 **G.6 Combination with 2016**

1731 This subsection presents the results of the combination of the unblinded 2017 dataset from
 1732 above with the 2016 dataset from Appendix F following the correlation model from Section 9.

1733 **G.6.1 Yields and results**

1734 Plots for the unblinded prefit and postfit event yields for the 2016+2017 data, with a total lu-
 1735 minosity of $35.9 \text{ fb}^{-1} + 41.5 \text{ fb}^{-1} = 77.4 \text{ fb}^{-1}$, are shown for both the cut-based and BDT based
 1736 analysis in Figure 114. Numerical yields are also tabulated in Table 52 (postfit cut-based analy-
 1737 sis), Table 53 (postfit BDT analysis).

1738 With the cut-based analysis, an observed (expected) upper limit on the production cross section
 1739 of 23.15 fb ($12.34^{+6.30}_{-4.00} \text{ fb}$), assuming the signal process does not exist. The observed (expected)
 1740 significance is 1.208 (2.074) standard deviations, corresponding to a measured signal strength
 1741 parameter of $0.696^{+0.690}_{-0.584}$.

1742 With the BDT analysis, an observed (expected) upper limit on the production cross section of
 1743 28.47 fb ($11.36^{+5.84}_{-3.71} \text{ fb}$), assuming the signal process does not exist. The observed (expected)
 1744 significance is 2.221 (2.272) standard deviations, corresponding to a measured signal strength
 1745 parameter of $1.167^{+0.697}_{-0.593}$.




Figure 114: 2016+2017: data yields compared to prefit (top) and postfit (bottom) SM predictions in CRZ, CRW, and the cut-based signal regions (left), and CRZ with the BDT signal regions (right). Note that we plan to retrain the BDT with latest JECs/corrections for 2018 (with identical settings)

Table 52: Postfit event yields in cut-based regions for 2016+2017.

	$\bar{t}\bar{W}$	$\bar{t}\bar{Z}$	$\bar{t}\bar{H}$	$\bar{t}\bar{V}V$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	1.45± 0.36	38.30± 6.11	1.57± 0.38	1.10± 0.13	0.40± 0.09	6.71± 1.25	0.02± 0.00	1.50± 0.74	51.05± 6.13	53	0.39± 0.35
CRW	70.12±16.58	17.21± 2.75	19.31± 4.64	3.79± 0.45	8.14± 0.98	4.44± 0.83	9.98± 1.16	35.72±18.98	168.72±14.28	174	1.85± 1.56
SR1	5.38± 1.34	1.71± 0.28	2.63± 0.66	0.62± 0.07	0.34± 0.07	0.39± 0.08	0.25± 0.03	2.96± 1.89	14.28± 1.65	18	0.91± 0.79
SR2	1.72± 0.45	0.35± 0.08	0.85± 0.21	0.20± 0.02	0.06± 0.02	0.07± 0.01	0.04± 0.01	1.32± 0.86	4.62± 0.79	5	0.54± 0.45
SR3	0.44± 0.15	0.14± 0.04	0.36± 0.10	0.08± 0.01	0.01± 0.00	0.03± 0.01	0.02± 0.00	0.14± 0.34	1.21± 0.40	1	0.36± 0.31
SR4	1.79± 0.44	0.64± 0.11	0.98± 0.23	0.28± 0.04	0.24± 0.05	0.12± 0.02	0.18± 0.02	1.47± 1.07	5.69± 0.92	10	0.71± 0.57
SR5	1.07± 0.31	0.23± 0.05	0.49± 0.13	0.14± 0.02	0.04± 0.01	0.11± 0.02	0.04± 0.00	0.46± 0.44	2.58± 0.53	2	0.75± 0.61
SR6	0.34± 0.11	0.06± 0.04	0.22± 0.06	0.05± 0.01	0.09± 0.03	0.03± 0.01	0.01± 0.00	0.12± 0.17	0.92± 0.24	5	0.54± 0.44
SR7	0.16± 0.06	0.06± 0.02	0.10± 0.03	0.02± 0.00	0.01± 0.00	0.01± 0.00	0.01± 0.00	0.00± 0.07	0.36± 0.12	0	0.40± 0.32
SR8	0.30± 0.09	0.10± 0.02	0.17± 0.04	0.10± 0.01	0.03± 0.01	0.05± 0.01	0.04± 0.01	0.44± 0.42	1.22± 0.45	0	0.96± 0.80
SR9	0.79± 0.20	1.21± 0.19	0.85± 0.20	0.17± 0.02	0.06± 0.01	0.06± 0.01	0.00± 0.00	0.39± 0.22	3.53± 0.34	3	0.33± 0.24
SR10	0.11± 0.04	0.41± 0.08	0.32± 0.08	0.05± 0.01	0.01± 0.00	0.02± 0.01	0.00± 0.00	0.55± 0.43	1.47± 0.44	3	0.18± 0.17
SR11	0.09± 0.03	0.19± 0.05	0.10± 0.03	0.02± 0.00	0.00± 0.00	0.01± 0.00	0.00± 0.00	0.06± 0.09	0.48± 0.12	0	0.10± 0.09
SR12	0.08± 0.03	0.31± 0.05	0.19± 0.04	0.04± 0.01	0.01± 0.00	0.03± 0.01	0.00± 0.00	0.57± 0.50	1.23± 0.49	2	0.28± 0.23
SR13	0.13± 0.05	0.15± 0.04	0.15± 0.04	0.03± 0.00	0.02± 0.00	0.02± 0.00	0.00± 0.00	0.00± 0.10	0.49± 0.15	0	0.32± 0.27
SR14	0.10± 0.03	0.12± 0.03	0.09± 0.02	0.02± 0.00	0.00± 0.00	0.01± 0.00	0.00± 0.00	0.01± 0.09	0.34± 0.12	1	0.38± 0.33

Table 53: Postfit event yields in BDT regions for 2016+2017.

	$\bar{t}\bar{W}$	$\bar{t}\bar{Z}$	$\bar{t}\bar{H}$	$\bar{t}\bar{V}V$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	1.46± 0.36	38.83± 6.01	1.58± 0.36	1.10± 0.12	0.40± 0.10	6.70± 1.18	0.02± 0.00	1.34± 0.52	51.42± 6.02	53	0.66± 0.35
SR1	0.25± 0.07	0.05± 0.02	0.04± 0.02	0.00± 0.00	0.03± 0.01	0.04± 0.03	0.12± 0.02	0.06± 0.13	0.58± 0.16	1	0.00± 0.00
SR2	2.69± 0.68	0.53± 0.14	0.44± 0.12	0.05± 0.01	0.64± 0.22	0.52± 0.18	0.88± 0.12	0.81± 0.57	6.56± 0.86	7	0.00± 0.01
SR3	5.76± 1.52	1.46± 0.31	1.17± 0.31	0.14± 0.02	1.46± 0.28	0.57± 0.15	1.26± 0.17	3.67± 1.37	15.50± 1.55	14	0.03± 0.02
SR4	8.76± 2.13	2.23± 0.54	1.99± 0.49	0.28± 0.03	0.80± 0.21	0.67± 0.18	1.43± 0.20	4.72± 1.82	20.89± 2.05	28	0.09± 0.05
SR5	10.13± 2.50	2.96± 0.68	2.66± 0.65	0.43± 0.04	1.82± 0.28	0.69± 0.30	1.46± 0.20	5.06± 2.04	25.21± 2.50	14	0.18± 0.11
SR6	11.04± 2.67	3.50± 0.60	3.17± 0.77	0.58± 0.06	1.31± 0.26	0.77± 0.18	1.37± 0.19	3.96± 1.86	25.70± 2.27	31	0.29± 0.19
SR7	10.50± 2.58	3.55± 0.66	3.40± 0.81	0.65± 0.07	1.00± 0.18	0.73± 0.18	1.14± 0.15	3.80± 1.91	24.77± 2.33	22	0.41± 0.23
SR8	9.40± 2.32	3.26± 0.60	3.21± 0.80	0.72± 0.09	0.88± 0.13	0.34± 0.08	0.99± 0.13	4.66± 1.92	23.45± 2.21	24	0.58± 0.30
SR9	11.50± 2.89	3.98± 0.71	4.40± 1.08	1.08± 0.11	0.77± 0.20	0.57± 0.12	0.92± 0.13	4.52± 2.34	27.73± 2.73	29	1.07± 0.54
SR10	10.62± 2.62	4.40± 0.70	4.63± 1.15	1.03± 0.12	0.53± 0.11	0.63± 0.12	0.77± 0.10	4.13± 1.95	26.75± 2.45	35	1.84± 1.00
SR11	5.91± 1.57	2.83± 0.51	3.00± 0.76	0.86± 0.10	0.71± 0.10	0.48± 0.10	0.44± 0.06	3.15± 1.88	17.37± 2.03	23	2.16± 1.14
SR12	3.46± 0.91	1.66± 0.29	1.94± 0.48	0.51± 0.06	0.12± 0.04	0.20± 0.05	0.23± 0.03	2.33± 0.96	10.45± 1.08	18	2.33± 1.21
SR13	1.79± 0.49	0.74± 0.13	1.11± 0.29	0.30± 0.04	0.02± 0.02	0.18± 0.03	0.12± 0.02	0.75± 0.50	4.99± 0.67	3	2.28± 1.22
SR14	1.00± 0.33	0.35± 0.08	0.55± 0.14	0.18± 0.02	0.03± 0.03	0.12± 0.02	0.06± 0.01	0.23± 0.23	2.53± 0.46	6	2.25± 1.18
SR15	0.38± 0.11	0.20± 0.04	0.27± 0.07	0.09± 0.01	0.07± 0.04	0.05± 0.01	0.03± 0.00	0.23± 0.13	1.33± 0.19	2	1.71± 0.88
SR16	0.11± 0.04	0.05± 0.03	0.06± 0.02	0.02± 0.00	0.00± 0.00	0.02± 0.00	0.01± 0.00	0.00± 0.06	0.27± 0.10	1	1.05± 0.56
SR17	0.01± 0.00	0.02± 0.01	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.19± 0.33	0.22± 0.33	0	0.11± 0.05

H Unblinding of 2018 dataset, BDT retraining

For reference, a table with “historical” numbers can be found here: <https://docs.google.com/spreadsheets/d/140yqrUwEsJtOJ8OmDdOun-J8iUrV49aAdUkvfNjpn8k/edit#gid=0> The last row of the table corresponds to the results presented below.

H.1 BDT retraining

With the availability of latest recipes for 2018, the BDT was retrained, keeping identical settings to what was used previously, in Section 6.3. In particular, the finalized retraining included two new key components: Autumn18V8 MC JECs and 2018 b-tag scale factors.

H.2 Yields and results

Plots for the prefit and postfit event yields for the 2018 data, with a total luminosity of 59.6 fb^{-1} , are shown for both the cut-based and BDT based analysis in Figure 115. Numerical yields are also tabulated in Table 54 (prefit cut-based analysis), Table 55 (postfit cut-based analysis), Table 56 (prefit BDT analysis), and Table 57 (postfit BDT analysis).

Similarly, plots of yields for combination of the full Run2 dataset, corresponding to 137.2 fb^{-1} , are shown for both the cut-based and BDT-based analyses in Figure 116. Full Run2 dataset numerical yields are also tabulated in Table 58 (prefit cut-based analysis), Table 59 (postfit cut-based analysis), Table 60 (prefit BDT analysis), and Table 61 (postfit BDT analysis).

A binned likelihood fit is performed over signal regions using the Higgs Combine tool [22] to extract signal strength and significance; exclusion limits at 95% CL are calculated with the Asymptotic CLs method.

Using the full run2 data, the *cut-based* analysis sets an observed (expected) upper limit on the production cross section of 20.31 fb ($9.40^{+4.38}_{-2.87} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 1.718 (2.461) standard deviations, corresponding to a measured observed (expected) signal strength parameter of $0.795^{+0.520}_{-0.475}$ ($1.000^{+0.485}_{-0.436}$)

The *BDT* analysis sets an observed (expected) upper limit on the production cross section of 22.86 fb ($8.56^{+3.96}_{-2.63} \text{ fb}$), assuming the signal process does not exist. The observed (expected) significance is 2.568 (2.680) standard deviations, corresponding to a measured observed (expected) signal strength parameter of $1.069^{+0.490}_{-0.443}$ ($1.000^{+0.447}_{-0.403}$)

To isolate the effect of the updates to 2018 data and MC, we recalculated the expected limit and significance with the latest corrections used above, but with the old BDT function, which gave an expected upper limit on the production cross section of $8.95^{+4.25}_{-2.80}$ (central value 4.6% worse than new training, relative), an expected significance of 2.636 (1.6% worse than new training, relative), and an expected signal strength of $1.000^{+0.470}_{-0.422}$ (measurement precision 5% worse than new training, relative).

Due to the retraining and updated SFs, updated numbers for the BDT analysis observed (expected) significances are as follows: 1.342 (1.496) for 2016, 1.182 (1.740) for 2017, and 2.950 (1.924) for 2018.




Figure 115: 2018: data yields compared to prefit (top) and postfit (bottom) SM predictions in CRZ, CRW, and the cut-based signal regions (left), and CRZ with the BDT signal regions (right).




Figure 116: Run2: data yields compared to prefit (top) and postfit (bottom) SM predictions in CRZ, CRW, and the cut-based signal regions (left), and CRZ with the BDT signal regions (right).

Table 54: Prefit event yields in cut-based regions for 2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	0.87±0.33	29.83±11.02	1.33±0.36	0.85±0.12	0.71±0.41	6.28±1.30	0.01±0.00	1.72±0.60	41.61±11.49	51	0.48±0.03
CRW	48.99±18.52	13.77±4.92	16.49±4.22	3.45±0.44	5.75±1.14	5.18±1.11	8.79±1.75	33.41±9.77	135.83±23.00	163	2.21±0.08
SR1	3.21±1.37	1.23±0.47	2.29±0.64	0.57±0.09	0.30±0.09	0.22±0.05	0.23±0.05	1.32±0.62	9.37±1.96	15	1.16±0.07
SR2	1.27±0.53	0.25±0.16	0.70±0.24	0.19±0.04	0.08±0.04	0.10±0.03	0.06±0.01	1.27±0.79	3.92±1.09	4	0.64±0.04
SR3	0.18±0.17	0.04±0.08	0.26±0.10	0.05±0.01	0.01±0.02	0.02±0.00	0.03±0.01	0.05±0.04	0.62±0.31	2	0.39±0.07
SR4	1.48±0.62	0.71±0.31	0.86±0.27	0.32±0.05	0.20±0.06	0.18±0.05	0.15±0.03	0.89±0.42	4.79±1.01	4	0.96±0.05
SR5	0.76±0.32	0.28±0.18	0.46±0.15	0.16±0.03	0.05±0.03	0.14±0.05	0.07±0.01	0.01±0.05	1.94±0.54	3	1.04±0.05
SR6	0.29±0.14	0.09±0.04	0.18±0.07	0.05±0.01	0.04±0.01	0.02±0.00	0.02±0.00	0.36±0.24	1.06±0.33	3	0.70±0.08
SR7	0.10±0.05	0.05±0.03	0.09±0.04	0.02±0.00	0.00±0.00	0.02±0.01	0.00±0.00	0.00±0.04	0.28±0.11	0	0.52±0.07
SR8	0.25±0.16	0.14±0.09	0.22±0.09	0.11±0.02	0.02±0.01	0.06±0.01	0.04±0.01	0.90±0.75	1.75±0.83	5	1.37±0.13
SR9	0.46±0.23	0.83±0.36	0.67±0.19	0.13±0.02	0.09±0.02	0.11±0.09	0.00±0.00	0.56±0.34	2.85±0.71	3	0.43±0.04
SR10	0.15±0.09	0.16±0.08	0.23±0.07	0.05±0.01	0.00±0.01	0.02±0.01	0.00±0.00	0.00±0.04	0.62±0.17	0	0.18±0.04
SR11	0.07±0.05	0.10±0.06	0.09±0.03	0.02±0.00	0.01±0.01	0.01±0.01	0.00±0.00	0.69±0.75	0.98±0.75	1	0.14±0.02
SR12	0.09±0.05	0.23±0.09	0.22±0.06	0.05±0.01	0.03±0.01	0.03±0.01	0.00±0.00	0.14±0.11	0.79±0.20	0	0.35±0.03
SR13	0.10±0.06	0.16±0.08	0.14±0.05	0.04±0.01	0.00±0.00	0.02±0.01	0.00±0.00	0.00±0.05	0.47±0.16	2	0.42±0.04
SR14	0.04±0.02	0.09±0.06	0.09±0.04	0.02±0.01	0.00±0.00	0.02±0.00	0.00±0.00	0.23±0.25	0.48±0.27	0	0.45±0.06

Table 55: Postfit event yields in cut-based regions for 2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	1.11±0.28	36.09±6.39	1.40±0.36	0.87±0.11	0.75±0.54	6.53±1.39	0.01±0.00	1.89±0.67	48.65±6.64	51	0.45±0.31
CRW	62.04±14.89	16.59±2.94	17.26±4.06	3.52±0.40	5.90±1.46	5.36±1.19	8.87±1.59	36.98±12.30	156.51±12.23	163	2.02±1.45
SR1	4.15±1.27	1.49±0.29	2.42±0.62	0.60±0.09	0.31±0.10	0.23±0.05	0.24±0.04	1.52±0.84	10.96±1.46	15	1.06±0.83
SR2	1.62±0.57	0.33±0.16	0.75±0.24	0.20±0.04	0.09±0.07	0.10±0.04	0.06±0.01	1.25±0.88	4.39±1.10	4	0.59±0.40
SR3	0.25±0.28	0.06±0.11	0.28±0.11	0.05±0.02	0.01±0.02	0.02±0.00	0.03±0.00	0.05±0.04	0.75±0.48	2	0.37±0.30
SR4	1.88±0.60	0.85±0.23	0.91±0.26	0.33±0.05	0.21±0.10	0.19±0.05	0.15±0.03	0.93±0.38	5.44±0.87	4	0.88±0.61
SR5	0.97±0.30	0.35±0.16	0.49±0.15	0.17±0.03	0.06±0.06	0.15±0.05	0.07±0.01	0.02±0.05	2.28±0.48	3	0.95±0.67
SR6	0.39±0.16	0.11±0.04	0.20±0.08	0.05±0.01	0.04±0.02	0.03±0.01	0.02±0.00	0.43±0.30	1.27±0.36	3	0.66±0.47
SR7	0.13±0.06	0.06±0.03	0.10±0.04	0.02±0.00	0.00±0.00	0.02±0.01	0.00±0.00	0.00±0.05	0.32±0.12	0	0.49±0.36
SR8	0.34±0.19	0.17±0.10	0.24±0.10	0.12±0.02	0.03±0.02	0.07±0.02	0.04±0.01	1.18±0.68	2.18±0.75	5	1.28±0.92
SR9	0.58±0.29	1.00±0.24	0.71±0.18	0.14±0.02	0.09±0.02	0.12±0.15	0.00±0.00	0.60±0.37	3.24±0.64	3	0.39±0.26
SR10	0.18±0.14	0.20±0.11	0.24±0.07	0.06±0.01	0.00±0.02	0.02±0.01	0.00±0.00	0.00±0.04	0.71±0.22	0	0.17±0.15
SR11	0.09±0.08	0.13±0.07	0.10±0.03	0.02±0.01	0.01±0.01	0.01±0.01	0.00±0.00	0.68±0.69	1.03±0.70	1	0.13±0.08
SR12	0.12±0.07	0.28±0.10	0.23±0.07	0.05±0.01	0.03±0.01	0.03±0.01	0.00±0.00	0.14±0.11	0.88±0.22	0	0.32±0.21
SR13	0.14±0.08	0.20±0.10	0.15±0.05	0.05±0.01	0.00±0.00	0.02±0.01	0.00±0.00	0.01±0.05	0.56±0.16	2	0.39±0.27
SR14	0.05±0.03	0.11±0.06	0.10±0.04	0.02±0.01	0.00±0.00	0.02±0.00	0.00±0.00	0.19±0.27	0.48±0.29	0	0.42±0.30

Table 56: Prefit event yields in BDT regions for 2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	0.87±0.31	29.83±11.11	1.33±0.35	0.85±0.13	0.71±0.43	6.28±1.56	0.01±0.00	1.72±0.66	41.61±11.56	51	0.48±0.03
SR1	0.49±0.38	0.18±0.10	0.08±0.06	0.01±0.01	0.24±0.20	0.11±0.17	0.29±0.05	0.36±0.36	1.75±0.87	1	0.00±0.00
SR2	1.66±0.86	0.43±0.28	0.38±0.20	0.03±0.02	0.95±0.49	0.49±0.19	0.68±0.13	1.97±1.04	6.60±1.58	8	0.00±0.01
SR3	3.54±1.50	0.89±0.48	0.86±0.31	0.13±0.04	0.97±0.29	0.76±0.24	1.08±0.20	4.74±2.28	12.98±3.15	13	0.02±0.01
SR4	5.09±2.29	1.23±0.70	1.38±0.42	0.21±0.04	0.40±0.47	0.71±0.30	1.17±0.22	3.95±1.41	14.14±3.61	12	0.04±0.01
SR5	6.26±2.38	1.53±1.03	1.68±0.50	0.29±0.06	0.38±0.32	0.61±0.35	1.19±0.22	3.47±1.57	15.40±3.92	17	0.06±0.01
SR6	6.38±2.61	1.83±0.89	1.96±0.54	0.35±0.05	0.03±0.62	0.59±0.21	1.06±0.20	4.10±1.97	16.30±4.32	25	0.10±0.02
SR7	6.05±2.30	2.23±1.25	2.11±0.58	0.42±0.08	0.91±0.20	0.51±0.13	0.87±0.16	4.74±2.16	17.83±3.90	19	0.15±0.03
SR8	7.65±2.83	2.85±1.70	2.84±0.77	0.58±0.08	0.86±0.24	0.59±0.26	0.89±0.17	3.86±1.57	20.12±4.17	21	0.29±0.04
SR9	8.18±3.15	3.26±2.41	3.83±0.98	0.85±0.12	0.93±0.24	0.86±0.27	0.99±0.19	6.14±2.59	25.05±5.46	19	0.45±0.06
SR10	7.28±2.75	2.93±1.82	3.49±0.95	0.89±0.11	0.79±0.33	0.55±0.28	0.69±0.13	5.99±2.66	22.60±4.84	32	0.82±0.04
SR11	5.77±2.33	2.94±1.59	3.47±0.93	0.98±0.12	0.37±0.07	0.51±0.15	0.52±0.10	2.60±1.35	17.14±3.76	35	1.36±0.06
SR12	3.71±1.35	2.00±1.04	2.47±0.68	0.71±0.09	0.33±0.21	0.51±0.15	0.31±0.06	3.04±1.69	13.08±2.83	17	1.63±0.07
SR13	2.10±0.87	1.07±0.68	1.53±0.42	0.48±0.07	0.18±0.04	0.27±0.09	0.13±0.03	1.22±0.51	6.98±1.54	5	1.97±0.09
SR14	0.97±0.44	0.66±0.33	0.89±0.26	0.27±0.04	0.08±0.02	0.13±0.03	0.10±0.02	1.71±1.10	4.81±1.39	6	1.81±0.10
SR15	0.78±0.37	0.41±0.20	0.48±0.14	0.19±0.02	0.16±0.11	0.10±0.02	0.07±0.01	0.19±0.10	2.37±0.59	3	1.75±0.14
SR16	0.17±0.10	0.10±0.09	0.13±0.05	0.08±0.01	0.01±0.00	0.05±0.01	0.01±0.00	0.70±0.47	1.26±0.51	4	1.51±0.11
SR17	0.00±0.02	0.05±0.05	0.02±0.01	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.00±0.03	0.10±0.08	3	0.51±0.07

Table 57: Postfit event yields in BDT regions for 2018.

	tW	tZ	tH	tVV	X+ γ	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	ttf
CRZ	1.01±0.25	36.74±6.82	1.43±0.31	0.89±0.12	0.71±0.41	6.60±1.34	0.01±0.00	1.86±0.67	49.25±6.88	51	0.70±0.29
SR1	0.50±0.23	0.18±0.06	0.07±0.04	0.01±0.00	0.14±0.15	0.09±0.09	0.29±0.05	0.34±0.33	1.62±0.59	1	0.00±0.00
SR2	1.83±0.58	0.45±0.17	0.35±0.14	0.03±0.01	0.85±0.32	0.40±0.15	0.68±0.12	2.30±1.17	6.89±1.34	8	0.00±0.01
SR3	3.82±1.08	0.95±0.29	0.79±0.26	0.11±0.03	0.82±0.28	0.73±0.22	1.08±0.19	5.04±1.99	13.34±2.15	13	0.02±0.02
SR4	5.79±1.59	1.42±0.37	1.36±0.35	0.19±0.04	0.30±0.37	0.71±0.23	1.17±0.21	4.08±1.53	15.02±2.15	12	0.05±0.03
SR5	7.02±1.78	1.76±0.52	1.69±0.42	0.29±0.05	0.53±0.24	0.65±0.24	1.19±0.21	3.74±1.42	16.86±2.17	17	0.08±0.04
SR6	7.41±1.92	2.04±0.52	1.99±0.47	0.35±0.05	0.37±0.46	0.63±0.17	1.06±0.19	4.97±2.09	18.83±2.36	25	0.14±0.08
SR7	6.97±1.75	2.56±0.66	2.15±0.51	0.43±0.07	0.91±0.14	0.51±0.11	0.87±0.15	5.03±1.92	19.43±2.11	19	0.19±0.10
SR8	8.77±2.15	3.18±0.92	2.96±0.69	0.57±0.08	0.90±0.21	0.56±0.18	0.89±0.16	4.09±1.71	21.94±2.44	21	0.42±0.19
SR9	9.62±2.42	3.95±1.19	3.99±0.89	0.86±0.12	0.94±0.18	0.84±0.23	0.99±0.17	5.54±2.32	26.73±3.16	19	0.65±0.30
SR10	8.65±2.14	3.56±0.91	3.76±0.86	0.92±0.12	0.69±0.21	0.60±0.20	0.69±0.12	7.02±2.90	25.88±2.88	32	1.19±0.52
SR11	7.01±1.78	3.70±0.83	3.73±0.85	1.01±0.13	0.38±0.07	0.56±0.14	0.52±0.09	3.48±1.58	20.39±2.05	35	1.94±0.85
SR12	4.24±1.06	2.45±0.53	2.67±0.62	0.72±0.10	0.46±0.16	0.50±0.13	0.31±0.05	3.28±1.49	14.63±1.65	17	2.36±0.99
SR13	2.52±0.68	1.27±0.33	1.65±0.39	0.51±0.07	0.20±0.03	0.26±0.08	0.13±0.02	1.23±0.45	7.79±0.92	5	2.86±1.24
SR14	1.15±0.35	0.79±0.17	0.95±0.25	0.28±0.04	0.08±0.02	0.13±0.03	0.10±0.02	1.62±1.02	5.10±1.12	6	2.66±1.13
SR15	0.89±0.30	0.48±0.13	0.51±0.14	0.19±0.03	0.16±0.09	0.10±0.02	0.07±0.01	2.59±0.51	3	2.61±1.13	
SR16	0.19±0.09	0.13±0.05	0.14±0.06	0.09±0.01	0.01±0.00	0.05±0.01	0.01±0.00	0.78±0.40	1.40±0.42	4	2.20±0.93
SR17	0.00±0.01	0.06±0.03	0.03±0.02	0.02±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.12±0.07	3	0.77±0.34	

Table 58: Prefit event yields in cut-based regions for Run2.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	1.91±0.70	60.40±22.00	2.79±0.73	1.93±0.22	0.85±0.41	12.86±2.77	0.03±0.00	2.94±1.10	83.71±22.75	104	1.05±0.06
CRW	98.82±35.54	27.53±10.10	34.45±8.64	7.20±0.79	13.76±1.80	9.33±2.08	18.72±2.14	59.11±21.59	268.90±45.98	337	4.87±0.20
SR1	7.00±2.66	2.59±0.96	4.72±1.30	1.18±0.15	0.63±0.14	0.60±0.14	0.48±0.06	3.37±1.41	20.57±3.76	33	2.47±0.14
SR2	2.49±0.96	0.53±0.28	1.49±0.47	0.38±0.06	0.14±0.05	0.16±0.04	0.10±0.01	2.23±1.24	7.53±1.87	9	1.41±0.07
SR3	0.49±0.33	0.15±0.11	0.59±0.22	0.12±0.02	0.01±0.02	0.04±0.01	0.04±0.01	0.19±0.18	1.64±0.61	3	0.91±0.12
SR4	2.74±1.00	1.22±0.53	1.76±0.49	0.59±0.07	0.43±0.14	0.29±0.07	0.32±0.04	1.86±0.87	9.21±1.74	14	1.97±0.10
SR5	1.51±0.63	0.47±0.21	0.91±0.29	0.30±0.04	0.10±0.04	0.26±0.07	0.10±0.01	0.40±0.28	4.05±0.97	5	2.10±0.10
SR6	0.52±0.20	0.13±0.10	0.39±0.14	0.10±0.02	0.12±0.05	0.05±0.01	0.02±0.00	0.41±0.27	1.75±0.48	8	1.46±0.09
SR7	0.22±0.09	0.10±0.04	0.18±0.07	0.04±0.01	0.01±0.01	0.03±0.01	0.01±0.00	0.00±0.08	0.59±0.19	0	1.09±0.10
SR8	0.47±0.24	0.21±0.11	0.38±0.14	0.21±0.03	0.05±0.02	0.11±0.02	0.08±0.01	1.36±0.81	2.86±0.93	5	2.73±0.17
SR9	1.03±0.44	1.79±0.66	1.46±0.39	0.30±0.04	0.15±0.02	0.16±0.09	0.00±0.00	0.86±0.45	5.75±1.18	6	0.90±0.08
SR10	0.23±0.10	0.48±0.17	0.52±0.16	0.11±0.02	0.01±0.01	0.04±0.01	0.00±0.00	0.35±0.24	1.73±0.39	3	0.44±0.06
SR11	0.13±0.06	0.25±0.11	0.19±0.06	0.04±0.01	0.01±0.01	0.02±0.01	0.00±0.00	0.74±0.74	1.38±0.76	1	0.29±0.03
SR12	0.15±0.08	0.48±0.17	0.39±0.10	0.09±0.01	0.04±0.01	0.06±0.02	0.00±0.00	0.57±0.37	1.77±0.46	2	0.75±0.04
SR13	0.20±0.10	0.28±0.11	0.28±0.09	0.07±0.01	0.02±0.01	0.04±0.01	0.00±0.00	0.00±0.09	0.89±0.24	2	0.87±0.05
SR14	0.11±0.05	0.18±0.09	0.17±0.07	0.04±0.01	0.00±0.00	0.03±0.01	0.00±0.00	0.23±0.26	0.76±0.32	1	1.00±0.09

Table 59: postfit event yields in cut-based regions for Run2.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	2.74±0.57	75.94±10.52	3.00±0.68	1.97±0.24	0.86±0.38	13.33±2.24	0.03±0.00	3.25±1.03	101.14±10.08	104	0.84±0.50
CRW	141.84±27.96	34.54±4.77	37.01±8.03	7.31±0.83	14.04±1.78	9.71±1.73	18.78±2.53	66.93±19.96	330.16±18.88	337	3.88±2.28
SR1	10.20±2.18	3.26±0.47	5.12±1.17	1.21±0.15	0.64±0.12	0.62±0.12	0.49±0.07	4.01±1.61	25.55±2.11	33	1.98±1.18
SR2	3.60±0.86	0.70±0.22	1.62±0.41	0.39±0.06	0.15±0.05	0.17±0.03	0.10±0.01	2.41±1.05	9.13±1.27	9	1.13±0.65
SR3	0.72±0.34	0.20±0.10	0.64±0.20	0.12±0.02	0.02±0.02	0.04±0.01	0.04±0.01	0.21±0.17	2.00±0.59	3	0.73±0.42
SR4	4.01±0.93	1.56±0.34	1.93±0.46	0.60±0.07	0.43±0.13	0.30±0.06	0.32±0.04	2.14±0.85	11.30±1.26	14	1.58±0.90
SR5	2.20±0.58	0.60±0.13	1.00±0.27	0.31±0.05	0.10±0.04	0.26±0.05	0.10±0.02	0.44±0.27	5.01±0.77	5	1.69±0.95
SR6	0.80±0.21	0.18±0.08	0.43±0.13	0.10±0.02	0.13±0.05	0.05±0.01	0.02±0.00	0.56±0.29	2.29±0.40	8	1.20±0.67
SR7	0.31±0.11	0.12±0.04	0.20±0.06	0.04±0.01	0.01±0.01	0.03±0.01	0.01±0.00	0.00±0.08	0.71±0.20	0	0.89±0.48
SR8	0.70±0.28	0.28±0.12	0.42±0.15	0.22±0.03	0.05±0.02	0.11±0.02	0.08±0.01	1.45±0.82	3.30±0.95	5	2.21±1.27
SR9	1.47±0.42	2.24±0.34	1.58±0.36	0.31±0.05	0.14±0.02	0.16±0.09	0.00±0.00	0.94±0.46	6.85±0.80	6	0.72±0.39
SR10	0.33±0.11	0.62±0.14	0.56±0.14	0.11±0.02	0.01±0.01	0.05±0.01	0.00±0.00	0.42±0.26	2.10±0.31	3	0.36±0.22
SR11	0.19±0.07	0.32±0.07	0.20±0.06	0.04±0.01	0.01±0.01	0.02±0.01	0.00±0.00	0.60±0.72	1.38±0.75	1	0.23±0.14
SR12	0.22±0.08	0.61±0.12	0.42±0.10	0.09±0.01	0.04±0.01	0.06±0.01	0.00±0.00	0.59±0.40	2.04±0.48	2	0.59±0.34
SR13	0.29±0.11	0.36±0.12	0.31±0.09	0.07±0.01	0.02±0.01	0.04±0.01	0.00±0.00	0.00±0.11	1.10±0.28	2	0.70±0.40
SR14	0.16±0.05	0.23±0.07	0.19±0.06	0.04±0.01	0.00±0.00	0.03±0.01	0.00±0.00	0.23±0.27	0.87±0.30	1	0.81±0.45

Table 60: Prefit event yields in BDT regions for Run2.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	1.91±0.76	60.40±19.18	2.79±0.60	1.93±0.25	0.85±0.36	12.86±2.43	0.03±0.00	2.94±0.97	83.71±19.68	104	1.05±0.06
SR1	1.01±0.65	0.27±0.15	0.19±0.13	0.02±0.01	0.39±0.27	0.18±0.24	0.66±0.07	0.86±0.55	3.58±1.29	4	0.00±0.00
SR2	3.40±1.73	0.81±0.44	0.79±0.34	0.07±0.03	0.21±0.61	0.84±0.27	1.52±0.17	2.72±1.28	12.31±2.61	19	0.01±0.01
SR3	6.73±3.26	1.79±0.78	1.70±0.57	0.25±0.05	1.61±0.43	1.12±0.38	2.26±0.26	7.91±2.79	23.35±5.16	19	0.04±0.02
SR4	10.36±4.68	2.44±1.11	2.80±0.80	0.38±0.08	1.64±0.63	1.17±0.45	2.51±0.28	7.87±2.55	29.17±6.61	33	0.08±0.03
SR5	12.19±5.18	3.01±1.54	3.41±0.95	0.57±0.10	1.44±0.61	1.21±0.53	2.59±0.29	6.65±2.23	31.08±7.00	35	0.15±0.05
SR6	12.42±5.43	3.66±1.52	4.07±1.02	0.74±0.09	0.94±0.85	1.34±0.37	2.17±0.25	7.74±2.76	33.08±7.38	44	0.22±0.06
SR7	12.18±5.10	4.43±2.07	4.38±1.10	0.80±0.14	2.17±0.35	1.01±0.26	1.86±0.21	7.27±2.42	34.09±6.84	41	0.31±0.07
SR8	15.09±6.14	5.59±2.83	6.03±1.44	1.19±0.15	1.60±0.37	1.19±0.42	1.95±0.22	7.17±2.61	39.80±8.25	46	0.67±0.05
SR9	16.78±7.08	6.74±3.65	7.87±1.83	1.85±0.25	2.11±0.34	1.54±0.42	2.02±0.23	11.11±4.16	50.01±10.62	48	1.11±0.08
SR10	14.71±6.01	6.14±3.04	7.45±1.70	1.85±0.24	1.43±0.44	1.12±0.36	1.40±0.16	8.86±3.31	42.96±8.47	61	1.78±0.08
SR11	11.80±5.11	6.03±2.64	7.20±1.66	2.00±0.26	0.95±0.19	0.91±0.22	1.07±0.12	4.80±2.15	34.75±7.12	62	2.80±0.15
SR12	7.32±3.04	4.14±1.72	5.18±1.21	1.51±0.18	0.69±0.22	0.88±0.26	0.61±0.07	6.41±2.60	26.76±5.08	40	3.50±0.14
SR13	4.14±1.85	2.38±1.16	3.13±0.77	1.00±0.13	0.25±0.06	0.51±0.13	0.28±0.03	2.31±0.88	14.00±2.99	15	3.97±0.16
SR14	2.04±1.00	1.30±0.50	1.76±0.47	0.54±0.07	0.11±0.08	0.30±0.05	0.18±0.02	2.21±1.13	8.43±1.93	16	3.88±0.17
SR15	1.37±0.73	0.67±0.29	0.95±0.27	0.37±0.05	0.20±0.10	0.17±0.03	0.12±0.01	0.44±0.24	4.27±1.12	4	3.79±0.22
SR16	0.42±0.24	0.20±0.12	0.31±0.13	0.14±0.02	0.07±0.04	0.10±0.02	0.03±0.00	0.84±0.48	2.12±0.66	7	3.15±0.19
SR17	0.03±0.03	0.07±0.05	0.03±0.02	0.02±0.00	0.00±0.00	0.02±0.00	0.00±0.00	0.25±0.30	0.44±0.33	3	1.00±0.10

Table 61: postfit event yields in BDT regions for Run2.

	$\bar{t}\bar{t}W$	$\bar{t}\bar{t}Z$	$\bar{t}\bar{t}H$	$\bar{t}\bar{t}VV$	$X+\gamma$	Rares	Charge misid.	Nonprompt lep.	SM expected	Data	$\bar{t}\bar{t}\bar{t}$
CRZ	2.51±0.54	77.27±11.74	3.09±0.65	1.98±0.21	0.87±0.36	13.29±2.58	0.03±0.00	3.25±1.19	102.30±11.59	104	1.12±0.43
SR1	1.24±0.49	0.31±0.11	0.19±0.10	0.01±0.01	0.32±0.25	0.17±0.19	0.66±0.07	1.05±0.65	3.95±0.95	4	0.00±0.00
SR2	4.33±1.28	0.96±0.32	0.82±0.29	0.08±0.02	0.16±0.53	0.78±0.23	1.53±0.17	3.49±1.46	14.15±1.77	19	0.01±0.01
SR3	8.48±2.41	2.14±0.50	1.74±0.53	0.23±0.05	1.43±0.47	1.10±0.38	2.26±0.26	8.08±2.79	25.45±3.53	19	0.04±0.03
SR4	13.42±3.36	3.02±0.68	2.98±0.77	0.37±0.08	1.56±0.49	1.18±0.40	2.51±0.29	8.81±3.10	33.86±4.00	33	0.08±0.05
SR5	15.71±3.69	3.71±0.93	3.66±0.93	0.57±0.08	1.57±0.51	1.25±0.45	2.59±0.29	7.42±2.61	36.48±4.00	35	0.15±0.07
SR6	16.26±3.87	4.47±0.99	4.38±1.03	0.74±0.08	1.22±0.67	1.39±0.33	2.17±0.25	9.05±3.47	39.68±4.14	44	0.23±0.12
SR7	15.88±3.61	5.48±1.28	4.72±1.10	0.81±0.12	2.21±0.31	1.01±0.27	1.86±0.21	8.18±3.08	40.17±3.72	41	0.32±0.16
SR8	19.71±4.39	6.83±1.75	6.58±1.47	1.19±0.14	1.64±0.32	1.19±0.34	1.95±0.22	8.03±2.83	47.11±4.34	46	0.72±0.28
SR9	22.13±4.94	8.52±2.23	8.59±1.90	1.87±0.21	2.13±0.30	1.52±0.43	2.02±0.24	11.59±3.95	58.35±5.27	48	1.18±0.46
SR10	19.58±4.27	7.76±1.85	8.27±1.79	1.89±0.21	1.36±0.36	1.17±0.29	1.40±0.16	10.58±3.98	52.01±4.30	61	1.92±0.74
SR11	15.88±3.65	7.77±1.53	7.99±1.74	2.04±0.23	0.98±0.17	0.96±0.22	1.07±0.12	6.16±2.65	42.86±3.51	62	3.00±1.19
SR12	9.61±2.22	5.28±1.01	5.78±1.26	1.54±0.18	0.81±0.21	0.88±0.24	0.61±0.07	7.52±3.06	32.03±3.07	40	3.76±1.41
SR13	5.55±1.35	3.00±0.66	3.49±0.77	1.03±0.12	0.27±0.05	0.50±0.12	0.28±0.03	2.54±0.94	16.67±1.62	15	4.28±1.63
SR14	2.73±0.74	1.65±0.28	1.96±0.48	0.55±0.07	0.11±0.07	0.31±0.06	0.18±0.02	2.62±1.24	10.12±1.25	16	4.19±1.59
SR15	1.81±0.53	0.84±0.24	1.06±0.26	0.37±0.04	0.19±0.10	0.17±0.03	0.12±0.01	0.46±0.22	5.01±0.77	4	4.11±1.55
SR16	0.56±0.21	0.26±0.10	0.35±0.12	0.15±0.02	0.07±0.04	0.10±0.02	0.03±0.00	0.97±0.54	2.50±0.61	7	3.39±1.26
SR17	0.05±0.03	0.09±0.05	0.04±0.02	0.02±0.00	0.00±0.00	0.02±0.00	0.00±0.00	0.34±0.35	0.57±0.36	3	1.08±0.42

1783 H.3 Nuisances

1784 Two sets of nuisance pull values, expected and observed for cut-based and BDT analyses, are
1785 tabulated in Table 62 (expected cut-based analysis), Table 63 (observed cut-based analysis),
1786 Table 64 (expected BDT analysis), and Table 65 (observed BDT analysis).

1787 The most constrained nuisances correspond to normalization parameters for ttW and ttZ (
1788 “TTWSF” and “TTZSF”) due to high statistics in control regions and in the bulk (BDT). Their
1789 input normalization uncertainty is 40%.

DRAFT

Table 62: Expected nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.37, 0.99	+0.00, 0.97	-0.12
TTVV	+0.05, 1.00	+0.00, 0.99	-0.02
TTWSF	+0.09, 0.62	+0.00, 0.62	-0.06
TTZSF	-0.02, 0.45	+0.00, 0.45	+0.02
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.72, 0.94	+0.00, 0.98	-0.23
fakes	+0.25, 0.91	+0.00, 0.91	-0.09
pdf	+0.07, 0.99	-0.00, 0.99	-0.02
rares	-0.01, 0.99	+0.00, 0.99	+0.00
scale	-0.20, 0.97	-0.00, 0.99	+0.00
y2016_btaghf	+0.04, 0.99	+0.00, 0.99	-0.03
y2016_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_fakes_EWK	+0.04, 1.00	+0.00, 0.99	-0.01
y2016_flips	-0.02, 0.99	+0.00, 0.99	+0.00
y2016_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.02
y2016_isr	-0.10, 0.99	-0.00, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	-0.00, 0.99	-0.02
y2016_jer	-0.13, 1.12	+0.00, 0.99	+0.03
y2016_jes	+0.07, 0.97	+0.00, 0.99	-0.05
y2016_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2016_lumi	+0.02, 0.99	+0.00, 0.99	-0.02
y2016_prefire	-0.00, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.01, 0.99	-0.00, 0.99	+0.00
y2016_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_btaghf	+0.04, 0.99	+0.00, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	-0.00, 0.99	+0.00, 0.98	-0.00
y2017_flips	-0.02, 0.99	+0.00, 0.99	+0.01
y2017_fsrvar	+0.00, 0.99	-0.00, 0.99	+0.00
y2017_isr	+0.07, 0.99	-0.00, 0.99	-0.03
y2017_isrvar	+0.00, 0.99	-0.00, 0.99	-0.01
y2017_jer	-0.12, 1.20	+0.00, 0.99	+0.03
y2017_jes	+0.07, 1.00	+0.00, 0.99	-0.05
y2017_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2017_lumi	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_prefire	-0.01, 0.99	-0.00, 0.99	+0.01
y2017_pu	-0.01, 0.99	-0.00, 0.99	+0.00
y2017_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2018_btaghf	+0.09, 0.99	+0.00, 0.99	-0.05
y2018_btaglf	+0.12, 0.99	+0.00, 0.99	-0.04
y2018_fakes_EWK	+0.08, 1.03	+0.00, 0.99	-0.02
y2018_flips	-0.03, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_isr	+0.08, 0.99	-0.00, 0.99	-0.03
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_jer	-0.32, 1.08	+0.00, 0.99	+0.08
y2018_jes	+0.15, 0.98	+0.00, 0.99	-0.09
y2018_lep	+0.02, 0.99	+0.00, 0.99	-0.02
y2018_lumi	+0.06, 0.99	+0.00, 0.99	-0.07
y2018_pu	+0.00, 1.00	-0.00, 0.99	+0.00
y2018_trig	+0.02, 0.99	+0.00, 0.99	-0.03

Table 63: Observed nuisance pulls for the cut-based analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.49, 1.02	+0.24, 1.00	-0.12
TTVV	+0.06, 1.00	+0.03, 1.00	-0.02
TTWSF	+1.06, 0.58	+1.02, 0.56	-0.05
TTZSF	+0.57, 0.42	+0.61, 0.42	+0.06
XG	+0.04, 1.00	+0.04, 1.00	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.62, 0.92	+0.10, 0.97	-0.27
fakes	+0.55, 1.02	+0.34, 1.00	-0.09
pdf	+0.15, 0.98	+0.06, 1.00	-0.04
rares	+0.06, 1.00	+0.07, 1.00	+0.01
scale	-0.09, 0.91	-0.02, 0.96	-0.00
y2016_btaghf	+0.07, 0.99	+0.04, 0.99	-0.03
y2016_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2016_fakes_EWK	+0.08, 1.00	+0.06, 0.99	-0.01
y2016_flips	+0.00, 0.99	+0.01, 1.00	+0.01
y2016_fsrvar	+0.00, 0.99	-0.01, 1.04	+0.01
y2016_isr	-0.13, 0.99	-0.04, 0.99	+0.05
y2016_isrvar	+0.00, 0.99	-0.04, 1.06	-0.03
y2016_jer	-0.07, 1.04	+0.01, 0.98	+0.04
y2016_jes	+0.39, 1.07	+0.26, 1.12	-0.08
y2016_lep	+0.04, 0.99	+0.03, 0.99	-0.01
y2016_lumi	+0.05, 1.00	+0.04, 1.00	-0.02
y2016_prefire	-0.01, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.03, 0.99	+0.01, 0.99	-0.01
y2016_trig	+0.04, 0.99	+0.04, 0.99	-0.01
y2017_btaghf	+0.10, 1.00	+0.06, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	+0.24, 0.98	+0.22, 0.97	-0.01
y2017_flips	+0.00, 0.99	+0.01, 1.00	+0.01
y2017_fsrvar	+0.00, 0.99	-0.01, 0.99	+0.00
y2017_isr	+0.08, 0.99	+0.02, 0.99	-0.04
y2017_isrvar	+0.00, 0.99	-0.01, 0.99	-0.01
y2017_jer	-0.07, 1.24	+0.01, 0.96	+0.03
y2017_jes	+0.14, 0.93	+0.09, 0.93	-0.05
y2017_lep	+0.03, 0.99	+0.03, 0.99	-0.01
y2017_lumi	+0.05, 1.00	+0.04, 1.00	-0.01
y2017_prefire	-0.02, 0.99	-0.02, 0.99	+0.01
y2017_pu	+0.02, 1.00	+0.01, 1.00	-0.00
y2017_trig	+0.05, 0.99	+0.04, 0.99	-0.01
y2018_btaghf	+0.15, 0.99	+0.08, 0.99	-0.05
y2018_btaglf	+0.12, 0.99	+0.03, 0.99	-0.05
y2018_fakes_EWK	+0.15, 1.01	+0.09, 0.99	-0.03
y2018_flips	+0.01, 0.99	+0.02, 1.00	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_isr	+0.08, 0.99	+0.02, 0.99	-0.04
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_jer	-0.19, 1.16	+0.06, 1.05	+0.09
y2018_jes	+0.48, 1.03	+0.30, 1.08	-0.13
y2018_lep	+0.06, 0.99	+0.05, 0.99	-0.02
y2018_lumi	+0.20, 0.99	+0.17, 0.99	-0.06
y2018_pu	+0.00, 1.00	-0.00, 0.99	+0.00
y2018_trig	+0.09, 0.99	+0.07, 0.99	-0.02

Table 64: Expected nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s</i> + <i>b</i> fit	$\rho(\theta, \mu)$
	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	$\Delta x/\sigma_{\text{in}}, \sigma_{\text{out}}/\sigma_{\text{in}}$	
TTH	+0.31, 0.98	-0.00, 0.96	-0.11
TTVV	+0.06, 1.00	-0.00, 0.99	-0.02
TTWSF	+0.07, 0.64	-0.00, 0.63	-0.06
TTZSF	-0.03, 0.50	+0.00, 0.48	+0.03
XG	-0.01, 0.99	+0.00, 0.99	+0.00
alphas	+0.00, 0.99	+0.00, 0.99	+0.00
bb	+0.89, 0.94	-0.00, 0.99	-0.26
fakes	+0.31, 0.90	+0.00, 0.90	-0.09
pdf	+0.12, 0.97	-0.00, 0.99	-0.03
rares	-0.02, 0.99	+0.00, 0.99	+0.00
scale	-0.25, 0.98	+0.00, 0.99	+0.01
y2016_btaghf	+0.04, 0.99	+0.00, 0.99	-0.02
y2016_btaglf	+0.03, 0.99	-0.00, 0.99	-0.01
y2016_fakes_EWK	+0.03, 0.99	-0.00, 0.99	-0.01
y2016_flips	-0.02, 0.99	+0.00, 0.99	+0.01
y2016_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.01
y2016_isr	-0.13, 0.99	+0.00, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	-0.00, 0.99	-0.01
y2016_jer	-0.04, 1.03	+0.00, 0.99	+0.01
y2016_jes	+0.07, 0.90	+0.00, 0.99	-0.04
y2016_lep	+0.01, 0.99	+0.00, 0.99	-0.01
y2016_lumi	+0.01, 0.99	+0.00, 0.99	-0.02
y2016_prefire	-0.00, 0.99	-0.00, 0.99	+0.00
y2016_pu	+0.04, 0.99	-0.00, 0.99	-0.01
y2016_trig	+0.01, 0.99	+0.00, 0.99	-0.02
y2017_btaghf	+0.03, 0.99	+0.00, 0.99	-0.03
y2017_btaglf	+0.03, 0.99	+0.00, 0.99	-0.01
y2017_fakes_EWK	+0.03, 0.98	+0.00, 0.96	-0.01
y2017_flips	-0.02, 0.99	+0.00, 0.99	+0.00
y2017_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2017_isr	+0.07, 0.99	-0.00, 0.99	-0.02
y2017_isrvar	+0.00, 0.99	-0.00, 0.99	-0.01
y2017_jer	-0.00, 1.18	+0.00, 0.99	+0.00
y2017_jes	+0.07, 0.90	+0.00, 0.99	-0.04
y2017_lep	+0.00, 0.99	+0.00, 0.99	-0.01
y2017_lumi	+0.00, 0.99	+0.00, 0.99	-0.02
y2017_prefire	-0.00, 0.99	-0.00, 0.99	+0.01
y2017_pu	+0.03, 1.00	-0.00, 0.99	-0.01
y2017_trig	+0.00, 0.99	+0.00, 0.99	-0.02
y2018_btaghf	+0.08, 0.99	+0.00, 0.99	-0.05
y2018_btaglf	+0.11, 0.99	-0.00, 0.99	-0.04
y2018_fakes_EWK	+0.06, 0.98	+0.00, 0.98	-0.01
y2018_flips	-0.03, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_isr	+0.04, 0.99	-0.00, 0.99	-0.02
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	+0.00
y2018_jer	+0.01, 1.49	+0.00, 0.95	-0.00
y2018_jes	+0.12, 0.85	+0.00, 0.97	-0.08
y2018_lep	+0.01, 0.99	+0.00, 0.99	-0.02
y2018_lumi	+0.05, 0.99	+0.00, 0.99	-0.07
y2018_pu	+0.02, 1.01	-0.00, 0.99	-0.01
y2018_trig	+0.02, 0.99	+0.00, 0.99	-0.03

Table 65: Observed nuisance pulls for the BDT analysis. The final column indicates the correlation between the nuisance and the signal strength

name	<i>b</i> -only fit	<i>s + b</i> fit	$\rho(\theta, \mu)$
	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	$\Delta x / \sigma_{\text{in}}, \sigma_{\text{out}} / \sigma_{\text{in}}$	
TTH	+0.59, 1.02	+0.34, 1.01	-0.09
TTVV	+0.08, 1.00	+0.04, 1.00	-0.02
TTWSF	+0.65, 0.67	+0.77, 0.60	-0.00
TTZSF	+0.62, 0.43	+0.66, 0.43	+0.05
XG	+0.02, 0.99	+0.04, 1.00	+0.01
alphas	+0.00, 0.99	+0.00, 1.00	+0.00
bb	+0.88, 0.95	+0.07, 0.98	-0.27
fakes	+0.95, 0.99	+0.34, 0.99	-0.13
pdf	+0.04, 1.00	-0.04, 1.01	-0.02
rares	+0.03, 0.99	+0.04, 0.99	+0.00
scale	-0.25, 0.94	-0.05, 0.94	+0.00
y2016_btaghf	+0.08, 0.99	+0.05, 0.99	-0.02
y2016_btaglf	+0.04, 1.00	+0.01, 0.99	-0.01
y2016_fakes_EWK	+0.17, 0.99	+0.08, 0.99	-0.02
y2016_flips	-0.01, 0.99	+0.00, 1.00	+0.00
y2016_fsrvar	+0.00, 0.99	+0.02, 1.07	+0.02
y2016_isr	-0.19, 0.99	-0.07, 0.99	+0.04
y2016_isrvar	+0.00, 0.99	+0.01, 1.00	-0.01
y2016_jer	-0.06, 0.89	-0.03, 0.87	+0.01
y2016_jes	+0.04, 0.99	-0.04, 1.11	-0.03
y2016_lep	+0.04, 0.99	+0.03, 0.99	-0.01
y2016_lumi	+0.06, 1.00	+0.05, 1.00	-0.02
y2016_prefire	-0.01, 0.99	-0.00, 0.99	+0.00
y2016_pu	-0.01, 1.00	-0.05, 1.00	-0.01
y2016_trig	+0.05, 0.99	+0.04, 0.99	-0.02
y2017_btaghf	+0.09, 0.99	+0.08, 0.99	-0.03
y2017_btaglf	+0.05, 0.99	+0.02, 0.99	-0.01
y2017_fakes_EWK	+0.44, 1.03	+0.26, 0.98	-0.04
y2017_flips	-0.01, 0.99	+0.01, 1.00	+0.01
y2017_fsrvar	+0.00, 0.99	+0.02, 1.00	+0.01
y2017_isr	+0.08, 0.99	+0.02, 0.99	-0.02
y2017_isrvar	+0.00, 0.99	+0.00, 1.00	-0.01
y2017_jer	+0.09, 0.88	+0.07, 0.84	-0.01
y2017_jes	+0.15, 0.90	+0.09, 1.00	-0.04
y2017_lep	+0.05, 0.99	+0.04, 0.99	-0.01
y2017_lumi	+0.05, 1.00	+0.05, 1.00	-0.02
y2017_prefire	-0.01, 0.99	-0.01, 0.99	+0.01
y2017_pu	+0.05, 1.00	+0.03, 1.00	-0.01
y2017_trig	+0.04, 0.99	+0.05, 0.99	-0.01
y2018_btaghf	+0.14, 0.99	+0.09, 0.99	-0.05
y2018_btaglf	+0.06, 0.99	-0.03, 0.99	-0.04
y2018_fakes_EWK	+0.10, 0.98	-0.01, 0.99	-0.03
y2018_flips	-0.02, 0.99	+0.00, 0.99	+0.01
y2018_fsrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_isr	+0.04, 0.99	+0.01, 0.99	-0.02
y2018_isrvar	+0.00, 0.99	+0.00, 0.99	-0.00
y2018_jer	+0.74, 0.77	+0.50, 0.97	-0.14
y2018_jes	+0.21, 0.67	+0.14, 0.81	-0.07
y2018_lep	+0.06, 0.99	+0.05, 0.99	-0.02
y2018_lumi	+0.16, 0.99	+0.15, 0.99	-0.06
y2018_pu	+0.09, 1.01	+0.06, 0.99	-0.02
y2018_trig	+0.06, 0.99	+0.05, 0.99	-0.03

1790 **H.4 Impacts**

1791 The leading 30 nuisance impacts for two sets of impacts, expected and observed for cut-based
 1792 and BDT analyses, are shown in Figure 117 (expected cut-based analysis), Figure 118 (observed
 1793 cut-based analysis), Figure 119 (expected BDT analysis), and Figure 120 (observed BDT anal-
 1794 ysis). The leading expected nuisance in both cases corresponds to the $\sigma(\text{ttbb})/\sigma(\text{ttjj})$ scaling.
 1795 Note that the “prop binSS” nuisances for MC statistics include (and are dominated by) tight-
 1796 loose sideband statistics.

1797 The observed pulls show the most constrained/pulled nuisances correspond to normalization
 1798 parameters for ttW and ttZ, as we would expect from the control regions. “TTWSF” is moved
 1799 by approximately 1σ (0.8σ) with respect to the input nuisance sizes for the cut-based (BDT)
 1800 analysis. “TTZSF” is moved up by approximately 0.6σ (0.7σ) for the cut-based (BDT) analysis.




Figure 117: Expected nuisance impacts for the cut-based analysis.




Figure 118: Observed nuisance impacts for the cut-based analysis.




Figure 119: Expected nuisance impacts for the BDT-based analysis.




Figure 120: Observed nuisance impacts for the BDT-based analysis.

1801 **H.5 Goodness of fits**

1802 The goodness of fit distributions (using the saturated, Kolmogorov-Smirnov, and Anderson-Darling test statistics) with the signal+background fit to data for the cut-based and BDT analyses are shown in Figure 121. We note that the observation is generally within the bulk of the expected distributions for both analyses and all three fit types.

1803

1804




Figure 121: GOF test for the cut-based analysis (left) and BDT-based analysis (right) using the saturated (top), KS (middle), AD (bottom) test statistics.

Table 66: Event kinematics for tail BDT events

runlumievent	SRnum	year	nleps	id1	id2	pT1	pT2	eta1	eta2	phi1	phi2	Nj	Nb	MET	HT
276776:1569:2615397660	17	2016	2	11	13	131.313	71.6004	1.54234	0.574022	-0.434035	-1.93809	7	3	256.921	1458.27
283050:58:92826095	17	2016	2	-13	-11	48.3626	39.4692	-0.217067	-2.34033	2.24725	-0.568915	7	3	168.629	844.845
305377:944:1751515966	17	2017	2	13	13	146.593	117.307	-1.86365	0.71062	2.92705	0.0105647	7	3	212.99	836.855
315644:87:107703005	17	2018	2	-13	-13	102.218	24.2424	-1.61099	-0.637295	1.11559	-1.64171	7	5	82.6389	754.944
316240:915:1292896376	17	2018	2	11	13	46.3985	30.2089	0.580867	-1.1041	0.0199213	3.09587	7	4	56.2346	632.974
317435:1373:1969850764	17	2018	3	-11	-11	62.5848	38.0043	0.175213	0.108995	2.11431	-0.932109	9	2	95.8658	1622.7
321051:870:1284078109	18	2018	2	13	13	55.4322	50.6932	1.42346	0.887908	2.50299	-1.42865	8	4	78.4541	558.675
321774:5088657761	18	2018	2	-11	-11	112.112	107.589	1.17477	0.895216	1.18255	-2.27924	9	4	245.434	1466.6
322356:79:153159025	18	2018	2	-11	-13	54.3049	31.4315	-0.017497	-0.924084	1.47584	1.63024	8	4	174.04	650.043
323727:421:771405322	17	2018	2	11	11	55.2933	42.2259	-0.0188495	-1.35181	2.1227	-1.93872	7	3	126.791	592.848

1806 H.6 Tail BDT events

1807 We included event information about the 7+3 events in the two tail BDT bins in Table 66. As
 1808 expected, the high number of jets and b-tag multiplicity contribute to a high BDT score. We
 1809 don't see any localization in the detector (for example, due to HEM).

DRAFT

I Checks from ARC review

I.1 Additional uncertainty from $t\bar{t}+b\bar{b}$

Figure 122 shows the fraction of events affected by the correction to account for the measured ratio of $t\bar{t}b\bar{b}/t\bar{t}j\bar{j}$ cross sections (1.7 ± 0.6), as discussed in Section 2. $t\bar{t}$ and $t\bar{t}Z$ show agreement in this fraction (approximately 1%), while $t\bar{t}W$ is low by 30%. To study possible effects due to differences in kinematics with respect to $t\bar{t}$, we tested a configuration where an additional 30% uncertainty was included in quadrature, so that the ratio and corresponding systematic uncertainty become 1.7 ± 0.78 . The effect on the analysis was to reduce the observed signal strength of the cut-based and BDT analyses by 0.5%, and the expected (observed) significance of both analyses by 1% (1.5%). This additional uncertainty was eventually not adopted in the analysis.




Figure 122: Fraction of events rewighted by measured ratio of $t\bar{t}b\bar{b}/t\bar{t}j\bar{j}$ for $t\bar{t}$, $t\bar{t}W$, and $t\bar{t}Z$, inclusively. Note that the same-sign requirement has been lifted from $t\bar{t}$, which requires opposite-sign dileptons.

1821 I.2 Distributions of BDT inputs for signal region events

1822 Figure 123 shows distributions of the 19 BDT input variables for the same events that enter
 1823 the BDT signal regions, except for CRZ. They are shown with “postfit yields”, meaning prefit
 1824 shapes are used, but normalization of processes is adjusted to match postfit/prefit scale factors.
 1825 Good agreement is observed.




Figure 123: Run2 Data to prediction comparisons for the BDT input variables. From left top to right bottom, \cancel{E}_T , N_{jets} , $N_{\text{b jets}}$, N_{leps} , H_T^b , N_{looseb} , N_{tightb} , $\Delta\phi(\ell_1, \ell_2)$, $\Delta\eta(\ell_1, \ell_2)$, $p_T(\ell_1)$, $p_T(\ell_2)$, $p_T(\ell_3)$, q_1 , $m(\ell_1, j_1)$, $\max(m(j)/p_T(j))$ and the p_T for jets 1, 6, 7, and 8, are shown for the BDT signal region events excluding CRZ, with scaled $t\bar{t}$ signal stacked when in solid red.

1826 I.3 More jets at higher η

1827 The analysis has a jet requirement of $|\eta| < 2.4$ to keep (b)jets almost completely within tracker
 1828 acceptance. Additionally, b-tag scale factor uncertainties are provided up to $|\eta|$ of 2.4 (2.5) for
 1829 2016 (2017, 2018) according to
 1830 <https://twiki.cern.ch/twiki/bin/view/CMS/BtagRecommendation>, and reflected
 1831 in the .csv files.

1832 While optimizing the Run 2 analysis, we considered relaxing both the p_T and $|\eta|$ requirements
 1833 for jets and b-jets. First, we studied the additional acceptance for signal jets when relaxing these
 1834 cuts. The additional jet acceptance from relaxing the $|\eta|$ requirement is small. Figure 124 shows
 1835 the $|\eta|$ distribution of $p_T > 30\text{GeV}$ generator-level jets which are matched to top decay prod-
 1836 ucts (light-flavor quarks and b quarks) in $t\bar{t}\ell\ell$ events with 2 leptonically-decaying W bosons.
 1837 The left plot shows an additional 3.8% of jets would be gained by completely relaxing the $|\eta|$
 1838 requirement up to approximately 3.6. The right plot shows up to 0.8% of jets would be gained
 1839 by relaxing $|\eta|$ to 2.5, in order to stay within an appropriate range of the b-tag scale factors and
 1840 allow for well-behaved JEC factors. Figure 125 shows the significantly more impressive gain
 1841 from relaxing p_T thresholds instead of η .

1842 As the gain from relaxing p_T thresholds was much larger, we put aside the η cut and focused on
 1843 the p_T cut. This resulted in the studies shown in Appendix C, where the whole analysis was re-
 1844 run with varying p_T cuts, signal and background counts were re-evaluated, but the significance
 1845 was not found to improve. For these reasons, we decided not to change our thresholds with
 1846 respect to the 2016 analysis.




Figure 124: $|\eta|$ distribution of generator-level jets which are matched to top decay products for 2 lepton $t\bar{t}\ell\ell$ events with coarse-binning (left) and fine-binning (right). The plots are normalized to unit area and bin values are superimposed.




Figure 125: p_T distribution of generator-level jets which are matched to top decay products for 2 lepton $t\bar{t}\ell\ell$ events.

1847 1.4 Effect of $t\bar{t}H$ and $t\bar{t}b\bar{b}/t\bar{t}j\bar{j}$ measurement updates

1848 In this section, we quantify the effects of potential updates to the $t\bar{t}H$ and $t\bar{t}b\bar{b}$ measurements
 1849 on this analysis.

1850 **I.4.1 $t\bar{t}H$**

1851 As we do not scale the $t\bar{t}H$ background, an updated measurement of $t\bar{t}H$ would only change the
 1852 25% relative uncertainty we place on this process. The effect of modifying this uncertainty can
 1853 be seen in the impact plots from Appendix A, which shows the $t\bar{t}H$ normalization uncertainty
 1854 nuisance is up to a 6% effect on the measured signal strength, approximately. Changes with
 1855 respect to the 25% normalization uncertainty can then be extrapolated from this 6%.

1856 **I.4.2 $t\bar{t}b\bar{b}/t\bar{t}j$**

1857 Conversely, an updated measurement of $t\bar{t}b\bar{b}/t\bar{t}j$ can change the central value of background
 1858 predictions. Currently, the $t\bar{t}b\bar{b}/t\bar{t}j$ ratio is 1.7 ± 0.6 . Based on TOP-18-002 (<http://cms.cern.ch/icMS/analysisadmin/cadilines?line=TOP-18-002>), the $t\bar{t}b\bar{b}/t\bar{t}j$ ratio in the
 1859 visible phase space is 1.17 ± 0.16 for the lepton+jets channels, and 1.38 ± 0.24 for the dilepton
 1860 channels. While a numerical combination is not provided, a crude estimate using the average
 1861 would be around 1.30 ± 0.20 . Thus, we use 1.3 ± 0.2 and re-evaluate the results of this analy-
 1862 sis. Numbers for both values/uncertainties of $t\bar{t}b\bar{b}/t\bar{t}j$ are tabulated in Table 67. The smaller
 1863 scale factor from $t\bar{t}b\bar{b}/t\bar{t}j$ results in 10% higher expected and observed significance values on
 1864 average.
 1865

BDT			
$t\bar{t}b\bar{b}/t\bar{t}j$	exp. σ	obs. σ	obs. μ
1.7 ± 0.6	2.699	2.561	$1.052^{+0.483}_{-0.438}$
1.3 ± 0.2	2.904	2.841	$1.108^{+0.470}_{-0.420}$
Cut-based			
$t\bar{t}b\bar{b}/t\bar{t}j$	exp. σ	obs. σ	obs. μ
1.7 ± 0.6	2.478	1.713	$0.784^{+0.514}_{-0.469}$
1.3 ± 0.2	2.629	1.968	$0.852^{+0.498}_{-0.450}$

Table 67